



# Assessment of Genetic Variability for Yield and Yield Contributing Characters in Rice (*Oryza sativa* L.)

Md. Nazmul Hasan Shahin <sup>a</sup>, Md. Harun-Or-Rashid <sup>b\*</sup>,  
Md. Atikuzzamman <sup>b</sup>, M. Raihan Uddin <sup>c</sup>  
and Md. Shahriar Kobir <sup>d</sup>

<sup>a</sup> Syngenta Bangladesh Limited, Bangladesh.

<sup>b</sup> International Maize and Wheat Improvement Center (CIMMYT), Bangladesh Office, Bangladesh.

<sup>c</sup> Department of Agricultural Extension, Ministry of Agriculture, Bangladesh.

<sup>d</sup> Bangladesh Agricultural Research Institute, Bangladesh.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AJAHR/2023/v10i4269

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104215>

Original Research Article

Received: 01/06/2023

Accepted: 04/08/2023

Published: 07/08/2023

## ABSTRACT

The study was conducted at the department of Genetics and Plant Breeding, Patuakhali Science and Technology University, Bangladesh during the period from April 2018 to August 2018 to study variability, heritability, genetic advance, interrelationship among 14 important characters among the 15 popular rice genotypes of Bangladesh. Significant variation in plant height of BRR1 dhan43, in 50% flowering BRR1 dhan42 and BRR1 dhan43, for days to maturity BRR1 dhan42 and BRR1 dhan43, for number of unfilled grain per panicle BRR1 dhan48, for flag leaf length Lakhi Lata, for flag leaf width Lohar Gura, for panicle plant<sup>-1</sup> BRR1 dhan27, for number of filled grain per panicle Kalo Jamvi, number of filled grain per panicle Lakhi Lata, for 100-grain weight BRR1 dhan48, for yield plant<sup>-1</sup> BRR1 dhan48 were found. Besides, yield per plant and 100-grain weight showed high genotypic and phenotypic co-efficient of variation while other characters showed moderate

\*Corresponding author: Email: harun.hmm1@gmail.com;

genotypic and phenotypic co-efficient of variation, except days to 50% flowering, days to maturity, panicle length, flag leaf length, flag leaf width, dry matter, and harvest index. All but, plant height and spikelet per panicle, all other characters studied in this experiment showed high heritability and high genetic advance in percent of mean. Panicle per plant, no. of filled grain, 100-seed wt., and harvest index are positively correlated. Yield per plant is also significantly and positively correlated with flag leaf length at the genotypic level but non-significant at phenotypic level. Grain yield showed non-significant positive association with flag leaf width and panicle length. Above information suggests that yield per plant would increase with increase of these characters. Path co-efficient analysis suggested that flag leaf length, panicle per plant, and flag leaf width showed moderate to low direct effect, but 100-seed weight showed highest positive direct effect toward yield per plant. Plant height showed a low negative direct effect on grain yield per plant while days to maturity, panicle length, no. of filled grain, no. of unfilled grain per panicle and dry matter had considerable negative direct effect on grain yield per plant and showed a negligible indirect effect. Thus, selection based on maturity period, number of panicles per plant, panicle length and higher number of filled grains per panicle might be effective for improving yield of rice.

**Keywords:** *Oryza sativa*; yield; South Asia; heritability; genetic variability.

## 1. INTRODUCTION

“Rice (*Oryza sativa* L.) is one of the most important cereal crops of the world meeting the dietary requirements of the people living in the tropics and sub-tropics. It is the main source of calories for almost 40% of the world population” [1]. It is mostly associated with the South, Southeast, and East Asian countries, extending from Pakistan to Japan. The region contains 18 of the 26 major rice producing countries, which account for 96% of global production. “Rice continues to be the major source of livelihood, especially in rural areas, and the main staple food of the population. In most Asian countries, therefore, government development agenda always been geared toward achieving self-sufficiency in rice. Among the major cereal crops, rice is unique, it can be grown in a wide range of agro-ecological conditions. Rice is very versatile tropical and sub-tropical or warm season temperate crop and is cultivated in 111 countries of all continents, from 53°N to 40°S latitude and from sea level to over 3000 m in the Himalayas” [2]. “Rice is a semi-aquatic plant and can grow successfully in standing water. Rice cultivation appears to be limited to those areas where the annual rainfall exceeds 1000 mm” [3]. “In Bangladesh, rice is grown in three overlapping seasons and farmers use a large number of varieties that fit various agro-ecological and climatic niches” [4]. And “farmers grow four diverse kinds of rice varieties: namely Aus, Deep-water Aman, Transplant Aman and Boro” [5].

“Rice is the leading cereal globally and a staple food crop in Bangladesh. Over 90% of Myanmar,

Sri Lanka, Vietnam, Cambodia, and Bangladesh rely on rice as their primary food source” [6]. For many farmers in Bangladesh, rice is the sole cash income [7]. In 1994-95, rice covered approximately 74.35% of the total cropped area [8]. Bangladesh imports 11,600 m tons of rice annually to meet its demand [9]. At the beginning of the 21st century, total rice production was about 590 million tons, a significant increase compared to the late 1970s [10]. Bangladesh ranks fourth in acreage and production [11] and 39th in yield among rice-growing countries [12]. The national average yield in Bangladesh is 2.3 tons ha<sup>-1</sup>, much lower than other countries like Japan, Korea, and Australia [13].

Rice plays a vital role in Bangladesh's agro-economy and national health. Per capita rice consumption is estimated at 153.4 kg year<sup>-1</sup> [14], covering 94% of cereals consumed in the country [5]. However, rice lacks beta-carotene, leading to vitamin A deficiency in populations dependent on rice as a staple food. Researchers have developed a modified grain with sufficient beta-carotene to address this issue (Swiss Federal Institute of Technology). Rice production in Bangladesh faces challenges due to variable monsoon rainfall. The country's average rice yield is 2852 kg ha<sup>-1</sup>, lower than the Asia average of 3906 kg ha<sup>-1</sup> [10]. Despite a growth rate of 1.8% in rice production from 1985 to 1995, population growth at 2% per year poses difficulties in overcoming food demands [4]. Therefore, the introduction and development of high-yielding rice varieties are crucial for increasing production [15].

Yield is the complex end product of many factors jointly or singly influence the seed yield [16].

“Rice yield is dependent on many important characters as well as on conservational influence” [17]. It is critical to understand the genetic variability of various features in order to optimize productivity. “Knowledge of nature and magnitude of genetic variation governing the inheritance of quantitative characters like yield and its components is essential for effecting genetic improvement. A critical analysis of genetic variability is a requirement for initiating any crop improvement program and for adopting appropriate selection techniques. The variability of a biological population is an outcome of genetic constitution of the individuals making up that population in relation to prevailing environment. A survey of genetic variability with the help of suitable parameters such as genotypic co-efficient of variation, habitability, and genetic advance necessary to start an efficient breeding program” [18].

About 5 tons ha<sup>-1</sup> of high yielding rice varieties are now being produced annually [19]. To get past that level, emphasis needs to be placed on characters who produce. As a result, character analysis is also required to ascertain how they contribute to yield. When choosing high yielding genotypes, both direct and indirect impacts of yield-contributing traits on yield are crucial. Path co-efficient analysis is used to detect characters having direct and indirect effects on yield. Several studies had been reported on variability [20-23] and characters association [24,25].

The present study was conducted to estimate and analyze the performance of 15 rice genotypes concerning yield and yield-contributing characters. Additionally, the study aimed to evaluate the interrelationship between yield and various yield-contributing characters. The ultimate goal was to identify the most promising genotypes for potential utilization in future agricultural endeavors.

## 2. MATERIALS AND METHODS

### 2.1 Experimentation

The experiment was situated at the rooftop of the Department of Genetics and Plant Breeding, Patuakhali Science and Technology University, Dumki, Patuakhali, during the period from April to August 2018 in ‘Chari’ (22cm) under irrigated condition. “The experimental site belongs to the sub-tropical monsoon climate and is characterized by high temperature and heavy rainfall during April to August and scant rainfall associated with moderate low temperature during

the rest period of the year. The soil of the experimental area was silty loam in texture belonging to the Gangas Flood Plain of AEZ 13 having non-calcareous dark grey plain soil. The selected area was medium highland. It was fertile, well drained, and slightly acidic with the pH varying from 5.5 to 6.8” [26]. The experiment was laid out in a Completely Randomized Design (CRD) with three replications. Seeds of fifteen rice genotypes were sown on March 10, 2018 followed by transplantation at the rate of 2 seedling hill<sup>-1</sup> in the main plot when they were 32 days old maintaining plant to plant distances 15 cm. The Chari were made by earthen, containing 2-3 aeration holes at the bottom for removal of excess water. Total 45 experimental pots were filled using soil & well decompose cow dung at the ratio of 2:1. Urea, Triple super phosphate (TSP), Muriate of potash (MP) and Gypsum were applied at the rate of 150, 100, 70 and 60 kg ha<sup>-1</sup>, respectively. 1/3 of the urea and entire TSP. MP and Gypsum were applied during the final land preparation. The remaining 2/3 of the urea was applied in two splits, one at 10 days after transplanting and the other before panicle initiation. Two weeding were done and only one irrigation was given. The other intercultural operations of exigent nature were done as and when needed. The crops were protected from the attack of pests by spraying diazinon @ 2ml per liter. Harvesting was started when 80% of the plant population of each “chari” reached maturity.

### 2.2 Data Recording

Data were recorded on individual hill basis from each hill from each ‘chari’ (pot). Data on days to 50% flowering, plant height and days to maturity (80%) were recorded in the field and the remaining namely length of flag leaf, width of flag leaf, number of panicles/plant, panicle length, number of filled grain/panicle, 100 grain weight and yield/plant were recorded in the laboratory after harvesting.

### 2.3 Statistical Analyses

All statistical analyses were performed using MSTAT-C. Analysis of variance was done for all the characters under study using the mean values [27]. Duncan's Multiple Range Test (DMRT) was used for all the characters to test the difference between the means of the genotypes following Steel and Torrie [28].

Genotypic and phenotypic variances were estimated according to the formula given by

Johnson et al. [29]. Genotypic and phenotypic co-efficient of variation were estimated according to Burton, [30]; Singh and Chaudhury, [31]. Heritability in broad sense was estimated using the formula suggested by Johnson et al. [29] and Hanson, et al. [32]. Expected genetic advance under selection was estimated using the formula suggested by Johnson et al. [29]. Genetic advance in percent of mean was calculated as proposed by Comstock and Robinson [33]. Genotypic and phenotypic correlation co-efficients were estimated using the formula suggested by Johnson et al. [29]; Miller et al. [34]; Singh and Chaudhury, [20]. The components of correlation co-efficient of different yield attributes with grain yield  $hill^{-1}$  were partitioned into components of direct and indirect effects by path co-efficient analysis. Path co-efficient analysis was done according to the procedure stated by Singh and Chaudhury [20] and Dabholkar [35] was originally suggested by Dewy and Lu [36]. In the present study, yield  $hill^{-1}$  was considered resulting characters (effect) and the yield attributes were considered as the causal factors.

The following sets of simulators equations were used depending the cause-and-effect relationship-

$$\begin{aligned}
 r_{1Y} &= P_{1y} + r_{1.2}P_{2y} + r_{1.3}P_{3y} + r_{1.4}P_{4y} + r_{1.5}P_{5y} + r_{1.6}P_{6y} + r_{1.7}P_{7y} + \dots + r_{1.14}P_{14y} \\
 r_{2Y} &= r_{2.1}P_{1y} + P_{2y} + r_{2.3}P_{3y} + r_{2.4}P_{4y} + r_{2.5}P_{5y} + r_{2.6}P_{6y} + r_{2.7}P_{7y} + \dots + r_{2.14}P_{14y} \\
 r_{3Y} &= r_{3.1}P_{1y} + r_{3.2}P_{2y} + P_{3y} + r_{3.4}P_{4y} + r_{3.5}P_{5y} + r_{3.6}P_{6y} + r_{3.7}P_{7y} + \dots + r_{3.14}P_{14y} \\
 r_{4Y} &= r_{4.1}P_{1y} + r_{4.2}P_{2y} + r_{4.3}P_{3y} + P_{4y} + r_{4.5}P_{5y} + r_{4.6}P_{6y} + r_{4.7}P_{7y} + \dots + r_{4.14}P_{14y} \\
 r_{5Y} &= r_{5.1}P_{1y} + r_{5.2}P_{2y} + r_{5.3}P_{3y} + r_{5.4}P_{4y} + P_{5y} + r_{5.6}P_{6y} + r_{5.7}P_{7y} + \dots + r_{5.14}P_{14y} \\
 r_{6Y} &= r_{6.1}P_{1y} + r_{6.2}P_{2y} + r_{6.3}P_{3y} + r_{6.4}P_{4y} + r_{6.5}P_{5y} + P_{6y} + r_{6.7}P_{7y} + \dots + r_{6.14}P_{14y} \\
 r_{7Y} &= r_{7.1}P_{1y} + r_{7.2}P_{2y} + r_{7.3}P_{3y} + r_{7.4}P_{4y} + r_{7.5}P_{5y} + r_{7.6}P_{6y} + P_{7y} + \dots + r_{7.14}P_{14y} \\
 &\dots\dots\dots \\
 R_{14Y} &= r_{14.1}P_{1y} + r_{14.2}P_{2y} + r_{14.3}P_{3y} + r_{14.4}P_{4y} + r_{14.5}P_{5y} + r_{14.6}P_{6y} + r_{14.7}P_{7y} + \dots + P_{14y}
 \end{aligned}$$

Where,

$r_{iy}$  = genotypic correlation co-efficient between  $y$  and  $i^{th}$  character ( $i = 1, 2, 3 \dots 14$ )  
 $P_{iy}$  = Path co-efficient due to  $i^{th}$  character ( $i = 1, 2, 3 \dots 14$ )

Total genotypic correlation say, between 1 and  $y$ , i.e.  $r_{1y}$  was thus partitioned as follows-

$$r_{1Y} = P_{1y} + r_{1.2}P_{2y} + r_{1.3}P_{3y} + r_{1.4}P_{4y} + r_{1.5}P_{5y} + r_{1.6}P_{6y} + r_{1.7}P_{7y} + \dots + r_{1.14}P_{14y}$$

$P_{1y}$  = the direct effect of 1 on  $y$   
 $r_{1.2}P_{2y}$  = the indirect effect of 1 via 2 on  $y$   
 $r_{1.3}P_{3y}$  = the indirect effect of 1 via 3 on  $y$   
 $r_{1.4}P_{4y}$  = the indirect effect of 1 via 4 on  $y$   
 $r_{1.5}P_{5y}$  = the indirect effect of 1 via 5 on  $y$   
 $r_{1.6}P_{6y}$  = the indirect effect of 1 via 6 on  $y$   
 $r_{1.7}P_{7y}$  = the indirect effect of 1 via 7 on  $y$   
 .....  
 $r_{1.14}P_{14y}$  = the indirect effect of 1 via 14 on  $y$

After calculating the direct and indirect effects of the characters, residual effect (R) was calculated using the following formula [20]:

$$P_{RY}^2 = 1 - \sum p_{iy}r_{iy}$$

Where,

$P_{RY}^2 = R_2$   
 $P_{iy}$  = direct effect of the characters on yield  
 $r_{iy}$  = correlation co-efficient of the characters with yield

Therefore,

$$\text{Residual effect} = \sqrt{P_{RY}^2} = \sqrt{1 - \sum p_{iy}r_{iy}}$$

### 3. RESULTS

#### 3.1 Genotypic Performance

Significant variations were observed among the rice genotypes for all studied traits. The genotypes were generally of medium to tall stature. BRR1 dhan 43 was the shortest, similar to BRR1 42. Lemma was the tallest, similar to Matiya and Ratul. Lemma took the longest time for 50% flowering, similar to Kalo Jamvi, Boilla, and Mari Dhan. BRR1 42 and BRR1 43 flowered earlier than others. Kalo Sathi, Lohar Gura, Munsu Murali, and Ratul had a medium flowering duration. Flag leaf length showed a wide range, with Lakhi Lata having the longest, and Lemma and BRR1 42 having the shortest. Flag leaf width didn't vary much, with Lohar Gura being the broadest and Lemma the narrowest. Days to maturity also varied widely, with BRR1 42 and BRR1 43 maturing earliest. Mari Dhan was late-maturing, similar to Lemma, Boilla, Matiya, Lakhi Lata, and Kalo Jamvi. Panicle length didn't show significant variation, with Lakhi Lata and Kalo Jamvi having the longest, and BRR1 42 and Lemma the shortest, similar to Parija. Kalo Jamvi had the highest number of Spikelet panicle<sup>-1</sup>,

while Lemma had the least. BRR1 27 had the highest Panicle plant<sup>-1</sup>, while Lakhi Lata had the least. Kalo Jamvi produced the most filled grain panicle<sup>-1</sup>, similar to Lakhi Lata. Lemma produced the fewest filled grain panicle<sup>-1</sup>. Lakhi Lata had the highest number of unfilled grain panicle<sup>-1</sup>, significantly different from others. BRR1 48 had the lowest unfilled grain panicle<sup>-1</sup>. Highest 1000-grain weight was observed in BRR1 48, similar to Lakhi Lata, Lohar Gura, Kalo Jamvi, Munsu Murali, and BRR1 42. Mari Dhan had the lowest 1000-grain weight, similar to BRR1 48, Lemma, Parija, and Ratul. BRR1 48 had the highest yield plant<sup>-1</sup>, while Mari Dhan had the lowest, similar to Lemma, Lohar Gura, Munsu Murali, Parija, and Ratul. Lakhi Lata and Parija had the highest dry matter plant<sup>-1</sup>, similar to most genotypes. BRR1 42 had the lowest dry matter plant<sup>-1</sup>, similar to Boilla. BRR1 42 had the highest harvest index, not significantly different from BRR1 48. Mari Dhan had the lowest harvest index, followed by Parija, Ratul, Lohar Gura, Munsu Murali, Lakhi Lata, and Matiya, not significantly different from each other. All of these are presented in Table 1.

### 3.2 Variability and Character Association

#### 3.2.1 Variability, heritability and genetic advance

The genotypic ( $\sigma^2g$ ), phenotypic ( $\sigma^2p$ ) and environmental variance ( $\sigma^2e$ ), genotypic coefficient of variation (GCV), phenotypic coefficient of variation (PCV), heritability in broad sense ( $h^2b$  %), genetic advance (GA) and genetic advance in percent of mean (GA %) for all the quantitative characters under study are presented in Table 4. The PCV were higher than their corresponding GCV for all the characters studied indicating that they all interacted with the environment beside some extent. Results revealed that highest GCV and PCV found in unfilled grain per panicle while lowest GCV and PCV were showed by 1000 grain weight and dry matter, respectively. The value of GA ranged from 12.19% to 76.41% where minimum and maximum values represented by the character of dry matter and unfilled grain per panicle. The studied characters showed medium to high heritability. However, flag leaf width showed high heritability (.588) and low genetic advance (28.40%) in percent of mean, indicating non-additive genetic control of those characters.

#### 3.2.2 Correlation study

Table 3 displays genotypic and phenotypic correlation coefficients for yield and yield-

contributing characters in rice. Genotypic coefficients were generally higher, indicating a strong inherent relationship among the characters, while lower phenotypic coefficients suggested environmental influences. Grain yield plant<sup>-1</sup> significantly correlated with panicle plant<sup>-1</sup>, filled grains, 1000-seed weight, and harvest index. Days to 50% flowering showed a significant correlation with days to maturity but not with plant height, unfilled grains panicle<sup>-1</sup>, or dry matter. Harvest index, yield plant<sup>-1</sup>, and panicle plant<sup>-1</sup> had non-significant negative correlations with the number of panicle plant<sup>-1</sup>. At the genotypic level, days to maturity positively correlated with plant height, days to 50% flowering, and unfilled grains panicle<sup>-1</sup> but negatively correlated with harvest index, yield plant<sup>-1</sup>, and panicle plant<sup>-1</sup>. Phenotypically, plant height, days to 50% flowering, and unfilled grains panicle<sup>-1</sup> showed positive correlations, while harvest index, yield plant<sup>-1</sup>, and panicle plant<sup>-1</sup> showed negative correlations with days to maturity. Plant height significantly correlated with days to 50% flowering, days to maturity, and dry matter. Genotypically, harvest index, yield (g plant<sup>-1</sup>), filled grains, and 1000-seed weight had negative correlations. Number of panicle plant<sup>-1</sup> significantly and negatively correlated with days to 50% flowering, flag leaf length, flag leaf width, days to maturity, panicle length, spikelet panicle<sup>-1</sup>, filled grains, unfilled grains panicle<sup>-1</sup>, and dry matter. Panicle length significantly correlated with flag leaf length, flag leaf width, spikelet panicle<sup>-1</sup>, filled grains, unfilled grains panicle<sup>-1</sup>, 1000-seed weight, and dry matter. Previous studies [37,38] found significant positive correlations between panicle length and filled grains panicle<sup>-1</sup>. Spikelet panicle<sup>-1</sup> significantly correlated with flag leaf length, flag leaf width, panicle length, filled grains, unfilled grains panicle<sup>-1</sup>, 1000-seed weight, and dry matter, while panicle plant<sup>-1</sup> showed a negative genotypic correlation. Unfilled grains panicle<sup>-1</sup> exhibited significant negative correlations with panicle plant<sup>-1</sup>, while days to 50% flowering, flag leaf length, days to maturity, panicle length, spikelet panicle<sup>-1</sup>, filled grains, and dry matter had positive correlations. Moreover, 1000-seed weight significantly correlated with flag leaf length, panicle length, spikelet panicle<sup>-1</sup>, filled grains, harvest index, and yield (g plant<sup>-1</sup>). Plant height and dry matter had negative genotypic correlations. Dry matter significantly and negatively correlated with panicle plant<sup>-1</sup>, 1000-seed weight, harvest index, and yield (g plant<sup>-1</sup>). However, plant height, days to 50% flowering, days to maturity, and unfilled grains panicle<sup>-1</sup>

**Table 1. Mean performance of 15 rice genotypes considering 14 important characters**

Variety	Plant height (cm)	Days to 50% flowering	Flag leaf length (cm)	Flag leaf width (cm)	Days to maturity	Panicle length (cm)	Spikelet panicle <sup>-1</sup>	Panicle plant <sup>-1</sup>	Number of filled grain panicle <sup>-1</sup>	Number of unfilled grain panicle <sup>-1</sup>	1000 seed wt. (g)	Yield plant <sup>-1</sup> (g)	Dry matter plant <sup>-1</sup> (g)	Harvest index
BRR1 27	141.3 b	83.00 f	28.78 bc	1.111 b	114.7 d	22.56 b	67.56 b-d	13.33 a	52.33 b-d	15.22 cd	18.48 b-e	12.81 b	21.47 ab	0.5975 bc
BRR1 42	102.0 g	66.00 h	17.89 d	1.000 b	100.3 f	16.44 f	64.78 cd	11.00 c	51.00 c-e	13.78 cd	20.76 a-c	11.62 bc	15.45 c	0.7523 a
BRR1 43	101.7 g	68.67 h	29.56 bc	1.222 ab	101.3 f	21.78 bc	76.78 b	11.33 c	60.67 b	16.11 cd	16.25 ef	11.09 cd	25.04 ab	0.4463 de
BRR1 48	108.7 f	79.67 g	30.78 b	1.000 b	109.7 e	21.11 b-d	69.67 b-d	12.33 b	60.67 b	9.000 d	22.03 a	16.41 a	25.25 ab	0.6502 ab
Boilla	117.7 e	114.3 a-c	30.33 b	1.316 ab	145.3 ab	20.22 c-e	76.44 b	8.333 e-g	58.33 bc	18.11 c	19.43 bc	9.407 d-f	19.86 a-c	0.4978 cd
Kalo Jamvi	123.5 de	115.7 ab	29.78 bc	1.333 ab	144.0 a-c	26.00 a	108.3 a	7.333 gh	76.67 a	31.67 b	19.59 a-c	10.99 cd	23.29 ab	0.4798 d
Kalo Sathi	129.3 cd	102.0 e	24.56 bc	1.111 b	141.7 bc	20.44 b-e	68.67 b-d	9.000 e	54.67 b-d	14.00 cd	18.91 b-d	9.334 d-f	22.40 ab	0.4177 de
Lakhi Lata	131.1 c	112.0 c	36.78 a	1.167 ab	143.7 a-c	26.00 a	114.3 a	6.667 h	74.00 a	40.33 a	20.91 ab	10.22 cd	26.32 a	0.3910 d-f
Lemma	150.7 a	116.0 a	13.33 d	0.500 c	145.7 ab	16.89 f	59.33 d	11.33 c	42.33 e	17.00 cd	16.18 ef	7.751 fg	19.18 bc	0.4136 de
Lohar Gura	141.7 b	103.3 e	25.44 bc	1.444 a	141.7 bc	21.00 b-d	67.22 b-d	7.667 fg	56.33 b-d	10.89 cd	19.55 a-c	8.397 e-g	24.89 ab	0.3357 ef
Mari Dhan	139.8 b	113.7 a-c	26.33 bc	1.333 ab	147.0 a	21.00 b-d	73.11 bc	8.333 e-g	56.00 b-d	17.11 cd	15.43 f	7.178 g	25.40 ab	0.2888 f
Matiya	149.4 a	112.3 bc	23.89 c	1.316 ab	143.7 a-c	19.22 de	76.56 b	8.667 ef	61.00 b	15.56 cd	18.25 c-e	9.630 de	24.04 ab	0.4006 d-f
Munsi Murali	150.9 a	106.7 d	26.33 bc	1.039 b	142.3 bc	21.28 b-d	66.22 b-d	8.333 e-g	48.67 de	17.56 c	20.29 a-c	7.997 e-g	22.80 ab	0.3569 ef
Parija	141.7 b	112.3 bc	24.89 bc	1.000 b	144.7 a-c	18.44 ef	63.33 cd	10.00 d	49.33 c-e	14.00 cd	15.68 f	7.665 fg	26.34 a	0.2927 f
Ratul	149.0 a	108.7 d	26.11 bc	1.111 b	141.0 c	20.22 c-e	69.44 b-d	9.333 de	52.67 b-d	16.78 cd	1649 d-f	8.010 e-g	24.08 ab	0.3348 ef
LSD Value	6.079	3.231	5.491	0.2848	3.521	1.997	9.331	0.9520	8.145	7.103	2.244	1.564	5.484	0.1058

\*\* , \* , ns denotes significant at 1% level of significance, significant at 5% level of significance and non-significant respectively. Values followed by same letter(s) are statistically similar as per DMRT at 5%

**Table 2. Maximum, minimum, mean, standard deviation and variance of fifteen rice genotype for different characters (Singh et al., 2011)**

Parameters	Plant height(cm)	Days to 50% flowering	Flag leaf length (cm)	Flag leaf width(cm)	Days to maturity	Panicle length(cm)	Spikelet/panicle	Panicle/plant	Number of filled grain	Number of unfilled grain/panicle	1000 seed wt. (g)	Dry matter(g)	Harvest index	Yield (g/plant)
Maximum	154.60	118.00	40.33	1.50	149.00	28.00	120.67	15.00	81.00	47.67	23.0	29.19	0.79	17.39
Minimum	98.00	64.00	11.67	0.50	98.00	14.67	54.33	5.00	37.00	8.00	13.8	14.49	0.24	6.59
Mean	131.89	100.96	26.32	1.13	133.78	20.84	74.79	9.53	56.98	17.81	18.5	23.05	0.44	9.90
Standard deviation	17.15	17.03	5.95	0.26	16.91	2.78	15.79	2.11	9.57	8.40	0.23	3.94	0.14	2.51
Variance	300.75	296.45	36.18	0.07	292.49	7.92	254.99	4.57	93.70	72.10	0.06	15.88	0.02	6.47
% CV	2.76%	1.57%	12.47%	15.05%	1.91%	5.73%	7.46%	12.07%	8.55%	23.85%	7.16%	14.22%	14.24%	9.44%

**Table 3. Analysis of variance (mean squares) for different characters of fifteen genotypes of rice**

Parameters	df	Panicle/plant	Number of filled grain	1000 seed wt. (g)	Yield (g/plant)	Spikelet/panicle	Number of unfilled grain/panicle	Panicle length(cm)	Flag leaf length (cm)	Flag leaf width(cm)	Dry matter(g)	Harvest index	Plant height(cm)	Days to 50% flowering	Days to maturity
Replication	2	2.467	8.622	0.59	4.472	3.262	2.921	1.864	6.965	0.078	8.027	0.004	11.603	18.956	15.089
Genotype	14	11.371**	245.832	1.32	17.932	738.685	190.103	21.771	91.164	0.154	27.256*	0.054	917.129	907.698	922.089
Error	28	1.324	23.717	0.18	0.874	31.124	18.037	1.426	10.778	0.029	10.753	0.004	13.211	4.432	3.732
% CV		12.07%	8.55%	7.16%	9.44%	7.46%	23.85%	5.73%	12.47%	15.05%	14.22%	14.24%	2.76%	1.57%	1.91%

**Table 4. Heritability, genetic advance and genetic advance in percentage of mean in respect of 14 yield and yield contributing characters of rice**

Characters	S <sup>2</sup> G	S <sup>2</sup> E	S <sup>2</sup> P	PCV	GCV	GA	% GA
Plant height(cm)	301.306	13.211	314.517	13.446	13.161	34.999	26.536
Days to 50% flowering	306.119	3.732	309.851	17.436	17.331	35.825	35.486
Flag leaf length(cm)	26.795	10.778	37.573	23.290	19.668	9.005	34.216
Flag leaf width(cm)	0.042	0.029	0.071	23.449	17.982	0.322	28.407
Days to maturity	301.089	4.432	305.521	13.066	12.971	35.485	26.525
Panicle length(cm)	6.782	1.426	8.208	13.747	12.496	4.876	23.399
Spikelet panicle <sup>-1</sup>	235.854	31.124	266.978	21.849	20.536	29.735	39.761
Panicle plant <sup>-1</sup>	3.349	1.324	4.673	22.675	19.197	3.192	33.478
No. of filled grain	74.038	23.717	97.756	17.353	15.102	15.426	27.074
No. of unfilled grain panicle <sup>-1</sup>	57.355	18.037	75.393	48.760	42.529	13.607	76.414
1000-seed wt.(g)	0.038	0.018	0.056	12.748	10.544	0.333	17.966
Dry matter(g)	5.501	10.753	16.254	17.487	10.173	2.811	12.192
Harvest index	0.017	0.004	0.021	32.274	28.961	0.238	53.537
Yield (g plant <sup>-1</sup> )	5.686	0.874	6.560	25.869	24.084	4.573	46.189

$\sigma^2_g$  = Genotypic Variation,  $\sigma^2_e$  = Environmental Variation,  $\sigma^2_p$  = Phenotypic Variation, GCV = Genotypic Coefficient of Variance, GCV = Phenotypic Coefficient of Variance, GA =Genetic Advance, % GA= Genetic Advance in Percent of Mean



**Table 5. Genotypic and phenotypic correlations between fifteen characters of 15 genotypes of rice**

Characters	Plant height(cm)	Days to 50% flowering	Flag leaf length (cm)	Flag leaf width(cm)	Days to maturity	Panicle length(cm)	Spikelet panicle <sup>-1</sup>	Panicle plant <sup>-1</sup>	No. of filled grain	No. of unfilled grain panicle <sup>-1</sup>	1000 seed wt.(g)	Dry matter(g)	Harvest Index	yield (g plant <sup>-1</sup> )
Plant height(cm)	<b><u>0.958</u></b>	0.700**	-0.271	-0.158	0.724**	-0.077	-0.196	-0.254	-0.368*	0.021	-0.412**	0.310*	-0.701**	-0.648**
Days to 50% flowering	0.682**	<b><u>0.988</u></b>	0.020	0.087	0.986**	0.169	0.265	-0.724**	0.114	0.408**	-0.290	0.359*	-0.749**	-0.689**
Flag leaf length (cm)	-0.199	0.019	<b><u>0.713</u></b>	0.668**	-0.005	0.918**	0.758**	-0.383**	0.854**	0.568**	0.424**	0.832**	-0.059	0.387**
Flag leaf width(cm)	-0.109	0.068	0.487**	<b><u>0.588</u></b>	0.133	0.576**	0.481**	-0.601**	0.660**	0.225	0.225	0.554**	-0.191	0.023
Days to maturity	0.704**	0.982**	-0.009	0.096	<b><u>0.985</u></b>	0.138	0.199	-0.768**	0.052	0.344*	-0.276	0.366*	-0.799**	-0.753**
Panicle length(cm)	-0.073	0.145	0.717**	0.400**	0.112	<b><u>0.826</u></b>	0.893**	-0.442**	0.895**	0.794**	0.338*	0.669**	-0.109	0.263
Spikelet panicle <sup>-1</sup>	-0.188	0.249	0.546**	0.328	0.193	0.737**	<b><u>0.883</u></b>	-0.601**	0.956**	0.942**	0.353*	0.456**	-0.068	0.153
Panicle plant <sup>-1</sup>	-0.245	-0.602**	-0.204	-0.450**	-0.643**	-0.371	-0.536**	<b><u>0.717</u></b>	-0.488**	-0.665**	-0.105	-0.468**	0.602**	0.531**
No. of filled grain	-0.325*	0.089	0.569**	0.510**	0.045	0.708**	0.896**	-0.506**	<b><u>0.757</u></b>	0.802**	0.471**	0.540**	0.061	0.372*
No. of unfilled grain panicle <sup>-1</sup>	0.016	0.368**	0.379*	0.037	0.312*	0.581**	0.862**	-0.432**	0.546**	<b><u>0.761</u></b>	0.180	0.312*	-0.207	-0.112
1000 seed wt.(g)	-0.338*	-0.226	0.230	0.013	-0.221	0.259	0.264	-0.143	0.283	0.174	<b><u>0.684</u></b>	-0.323*	0.655**	0.657**
Dry matter(g)	0.155	0.221	0.407**	0.256	0.216	0.391**	0.213	-0.169	0.289	0.072	-0.077	<b><u>0.338</u></b>	-0.668**	-0.156
Harvest index	-0.640**	-0.670**	-0.064	-0.180	-0.709**	-0.139	-0.010	0.516**	0.054	-0.081	0.482**	-0.611*	<b><u>0.805</u></b>	0.839**
Yield (g plant <sup>-1</sup> )	-0.630**	-0.634**	0.280	-0.060	-0.691**	0.199	0.158	0.541**	0.304*	-0.049	0.541**	0.010	0.763**	<b><u>0.867</u></b>

*Diagonal bold and underline values showed heritability of respective character. Above diagonal values are genotypic correlation and below diagonal values are phenotypic correlation*

**Table 6. Partitioning of genotypic correlations into direct and indirect effects of the studied morphological, physiological and yield contributing characters as influenced by fifteen genotypes by path analysis**

Characters	Plant height(cm)	Days to 50% flowering	Flag leaf length (cm)	Flag leaf width (cm)	Days to maturity	Panicle length (cm)	Spikelet panicle-1	Panicle plant-1	No. of filled grain	No. of unfilled grain panicle-1	1000 seed wt. (g)	Dry matter(g)	Harvest index	Yield (g plant-1)
Plant height(cm)	<b><u>-0.041</u></b>	0.279	-0.058	-0.007	-0.310	0.026	-0.472	-0.289	0.496	-0.318	-0.290	-0.049	0.385	-0.648
Days to 50% flowering	-0.029	<b><u>0.398</u></b>	0.004	0.004	-0.423	-0.058	0.711	-0.825	-0.132	-0.484	-0.204	-0.057	0.411	-0.689
Flag leaf length(cm)	0.011	0.008	<b><u>0.215</u></b>	0.199	0.602	-0.314	0.410	-0.437	-0.115	-0.396	0.298	-0.131	0.033	0.387
Flag leaf width(cm)	0.501	0.035	0.150	<b><u>0.043</u></b>	-0.057	-0.197	0.572	-0.685	-0.289	-0.267	0.159	-0.087	0.146	0.023
Days to maturity	-0.030	0.392	-0.001	0.006	<b><u>-0.429</u></b>	-0.047	0.479	-0.875	-0.030	-0.408	-0.195	-0.058	0.438	-0.753
Panicle length(cm)	0.400	0.673	0.197	0.025	-0.059	<b><u>-0.342</u></b>	0.215	-0.504	-0.421	-0.642	0.238	-0.106	0.586	0.263
Spikelet panicle <sup>-1</sup>	0.008	0.105	0.150	0.021	-0.205	-0.306	<b><u>0.241</u></b>	-0.168	-0.129	-0.112	0.248	-0.072	0.373	0.153
Panicle plant <sup>-1</sup>	0.010	-0.288	-0.082	-0.026	0.329	0.151	-0.145	<b><u>0.139</u></b>	0.085	0.689	-0.074	0.074	-0.330	0.531
No.of filled grain	0.015	0.075	0.183	0.428	-0.022	-0.306	0.630	-0.555	<b><u>-0.135</u></b>	-0.151	0.331	-0.085	-0.034	0.372
No. of unfilled grain panicle <sup>-1</sup>	-0.001	0.162	0.122	0.010	-0.148	-0.272	0.227	-0.184	-0.108	<b><u>-0.119</u></b>	0.127	-0.049	0.114	-0.112
1000-seed wt.(g)	0.017	-0.116	0.091	0.010	0.119	-0.116	0.849	-0.120	-0.347	-0.214	<b><u>0.704</u></b>	0.142	-0.359	0.657
Dry matter(g)	-0.013	0.143	0.179	0.024	-0.157	-0.229	0.110	-0.153	-0.023	-0.037	-0.208	<b><u>-0.158</u></b>	0.367	-0.156
Harvest index	0.029	-0.298	-0.013	-0.008	0.428	0.037	-0.163	0.657	-0.086	0.236	0.461	0.105	<b><u>-0.549</u></b>	0.839

*The bold and underlined values are direct effect toward yield per plant.  
Residual effect = 0.49*

showed positive correlations at the genotypic level. Lastly, harvest index significantly correlated with panicle plant<sup>-1</sup>, 1000-seed weight (g), and yield (g plant<sup>-1</sup>). Genotypically, plant height, days to 50% flowering, days to maturity, and dry matter showed negative correlations with harvest index.

### 3.2.3 Path co-efficient

From the results of path analysis (Table 4), it was evident that direct effects contributed by 1000 seed wt. (0.704), Days to 50% flowering (0.398) spikelet panicle<sup>-1</sup> (0.241) were high indicating that among the component traits, these three characters contributed maximum for yield plant<sup>-1</sup>. The direct effects of 1000 seed wt., Days to 50% flowering, spikelet panicle<sup>-1</sup> were higher than their respective correlation co-efficient but lower for number of filled grains panicle<sup>-1</sup>. The indirect effect of the above three characters not important. Thus, direct selection for these traits will be effective. Although the character, harvest index was positively correlated with yield and its direct effect on grain yield plant<sup>-1</sup> was negative (-0.549). It indicated that this character influenced grain yield by its indirect positive effects through days to maturity, 1000-grain weight, and panicle plant<sup>-1</sup>. Plant height showed a low negative direct effect (-0.041) on grain yield plant<sup>-1</sup> but the correlation co-efficient was significantly negative. Flag leaf length (0.215), panicle plant<sup>-1</sup> (0.139) and flag leaf width (0.043) showed moderate to low direct effect as the correlation coefficient was moderate to low for these characters. Days to maturity (-0.429), panicle length (-0.342), no. of filled grain (-0.135), no. of unfilled grain panicle<sup>-1</sup> (-0.119), dry matter (-0.158) had considerable negative direct effect on grain yield plant<sup>-1</sup> and showed negligible indirect effect. Therefore, the positive correlation between panicle length, no. of filled grain and grain yield plant<sup>-1</sup> may be explained by positive indirect effect via days to 50% flowering, harvest index and flag leaf width (cm) and spikelet panicle<sup>-1</sup> respectively.

The residual effect was 0.496 (Table 6), indicating that the eight characters contributed 51% percent of the variability in yield hill<sup>-1</sup> studied in path analysis. The residual effect towards grain yield in this study may be due to a variety of reasons such as causal factors (characters) not studied and sampling errors etc. Das et al. (1992) observed a residual effect of 0.364 indicating 64 percent of the variability in the yield plant<sup>-1</sup> contributed by 8 characters studied in path analysis. The correlation and path co-

efficient studies revealed that number of filled grains panicle<sup>-1</sup>, 1000-grain weight, days to maturity and days to 50% flowering are the most important yield components. The present study suggested that prolonged maturity period, higher number of panicles plant<sup>-1</sup>, longer panicle length and higher number of filled grains panicle<sup>-1</sup> should be given the importance in selection of genotypes for higher grain yield in rice.

## 4. DISCUSSION

### 4.1 Performance of the Genotypes

The genotypes varied in stature, with BRR1 43 being the shortest, statistically similar to BRR1 42. Lemma was the tallest and not significantly different from Matiya and Ratul. Lemma took the longest time to reach 50% flowering, similar to Kalo Jamvi, Boilla, and Mari Dhan. BRR1 42 and BRR1 43 reached 50% flowering earlier, not significantly different from each other. Kalo Sathi, Lohar Gura, Munsu Murali, and Ratul had a medium flowering duration. Flag leaf length showed a wide range, with Lakhi Lata having the longest, and Lemma and BRR1 42 having the shortest. Flag leaf width did not vary significantly, except for Lohar Gura, which produced the broadest, and Lemma, the narrowest. Days to maturity had a wide range, with BRR1 42 and BRR1 43 being the earliest, and Mari Dhan being late maturing, similar to Lemma, Boilla, Matiya, Lakhi Lata, and Kalo Jamvi. Panicle length did not vary significantly, but Lakhi Lata and Kalo Jamvi had the longest, while BRR1 42 and Lemma had the shortest, similar to Parija. Kalo Jamvi produced the highest number of spikelet panicle<sup>-1</sup>, while Lemma had the least. BRR1 27 produced the highest number of Panicle plant<sup>-1</sup>, while Lakhi Lata had the least, similar to Kalo Jamvi. Kalo Jamvi also produced the highest number of filled grain panicle<sup>-1</sup>, similar to Lakhi Lata. Lemma produced the lowest number of filled grain panicle<sup>-1</sup>. Lakhi Lata had the highest number of unfilled grain panicle<sup>-1</sup>, significantly different from others. BRR1 48 had the lowest number of unfilled grain panicle<sup>-1</sup>. BRR1 48 had the highest 1000-grain weight, similar to Lakhi Lata, Lohar Gura, Kalo Jamvi, Munsu Murali, and BRR1 42. Mari Dhan had the lowest 1000-grain weight, not significantly different from BRR1 48, Lemma, Parija, and Ratul. BRR1 48 had the highest yield plant<sup>-1</sup>, while Mari Dhan had the lowest, similar to Lemma, Lohar Gura, Munsu Murali, Parija, and Ratul. Lakhi Lata and Parija produced the highest dry matter plant<sup>-1</sup>, not significantly different from most genotypes. BRR1

42 produced the lowest dry matter plant<sup>-1</sup>, similar to Boilla. Among the genotypes, BRRI 42 had the highest harvest index, not significantly different from BRRI 48. Mari Dhan had the lowest harvest index, followed by Parija, Ratul, Lohar Gura, Munsu Murali, Lakhi Lata, and Matiya, which were not significantly different from each other.

## 4.2 Studies on Variability and Character Association

### 4.2.1 Variability, heritability, and genetic advance

The study focused on the assessment of different agronomic traits in rice genotypes and their variation, heritability, and genetic advance. The results indicated that some traits were influenced more by genetic factors, while others showed considerable environmental influence. Yield plant<sup>-1</sup> and 1000-grain weight exhibited high genotypic and phenotypic coefficients of variation, suggesting minimal environmental influence on these characters, thus providing a better scope for selection. The study concurred with Chauhan et al. [39], who found similar estimates for phenotypic (11.4%) and genotypic (11.1%) coefficients of variation for 1000-grain weight. For plant height, panicle plant<sup>-1</sup>, number of filled grains panicle<sup>-1</sup>, spikelet panicle<sup>-1</sup>, and number of unfilled grain panicle<sup>-1</sup>, moderate genotypic (13.161%) and phenotypic (13.446%) coefficients of variation were observed, indicating a considerable impact of the environment on these traits. Maurya et al. [21] reported comparable results for genotypic (10.4%) and phenotypic (11.7%) coefficients of variation. Similarly, Borbora and Hazarika found moderate genetic and phenotypic coefficients of variation for the number of filled grains panicle<sup>-1</sup> in rice. Maurya et al. [21] also reported moderate genotypic (10.4%) and phenotypic (11.7%) coefficients of variation for spikelet panicle<sup>-1</sup>. Conversely, days to 50% flowering, days to maturity, panicle length, flag leaf length, flag leaf width, dry matter, and harvest index showed low genotypic and phenotypic coefficients of variation, indicating minor environmental influence on these traits. Similar observations were made by Sarma et al. [40] for days to flowering, and by Rao and Shrivastav [41] for panicle length and dry matter. High heritability and genetic advance were found for days to maturity, the number of panicles plant<sup>-1</sup>, number of unfilled grain panicle<sup>-1</sup>, yield plant<sup>-1</sup>, flag leaf length, flag leaf width, dry matter, and harvest index in rice, suggesting a substantial non-

additive genetic control for these traits. Yadav [42] reported high heritability for yield plant<sup>-1</sup>, while Gupta et al. [43] found similar results for days to flowering. The 1000-grain weight exhibited high heritability (97.77%) and genetic advance (22.9%) in percent of mean, consistent with the findings of Chauhan et al. [44]. Sawant and Patil [45] also reported high coefficients of variation for yield plant<sup>-1</sup> in rice, as did Li et al. [46] and Das et al. [47]. Medium heritability (.958) and moderate genetic advance (26.53%) in percent of mean were observed for medium plant height and spikelet panicle<sup>-1</sup>, suggesting a balanced contribution of additive and non-additive gene actions in these traits. This finding was in line with Kumar et al. [48] who reported similar genetic advance for plant height and spikelet panicle<sup>-1</sup>. In conclusion, the study highlighted varying degrees of genetic and environmental influences on different agronomic traits in rice. Traits like 1000-grain weight and yield plant<sup>-1</sup> exhibited high heritability and low environmental influence, making them favorable targets for selection in breeding programs, while traits such as days to 50% flowering and panicle length showed low heritability and were more affected by environmental factors. Understanding these variations is crucial for effective trait improvement and breeding strategies in rice cultivation.

### 4.2.2 Correlation study

Genotypic correlation coefficients were generally higher than phenotypic correlation coefficients, indicating a strong inherent relationship among the characters [49,39]. Grain yield per plant showed significant positive correlations with panicle per plant, number of filled grains, 1000-seed weight, and harvest index [50,44]. Days to 50% flowering showed significant positive correlation with days to maturity, but non-significant correlation with plant height, number of unfilled grain per panicle, and dry matter [47,39]. On the other hand, harvest index, yield per plant, and panicles per plant showed non-significant negative correlations with the number of panicles per plant [41]. Plant height exhibited significant positive associations with days to 50% flowering, days to maturity, and dry matter, while harvest index, yield per plant, number of filled grains, and 1000-seed weight showed negative correlations at the genotypic level [49,50]. The number of panicles per plant showed significant negative associations with days to 50% flowering, flag leaf length, flag leaf width, days to maturity, panicle length, spikelet per panicle,

panicles per plant, number of filled grains, and number of unfilled grain per panicle [51]. Flag leaf length, flag leaf width, panicle length, spikelet per panicle, number of filled grains, 1000-seed weight, dry matter, and yield per plant showed significant positive correlations with each other [37,38]. Spikelet per panicle exhibited significant positive associations with flag leaf length, flag leaf width, panicle length, number of filled grains, 1000-seed weight, and dry matter, while panicles per plant showed negative correlation at the genotypic level [49,50]. Number of unfilled grain per panicle showed negative and significant associations with panicles per plant. On the other hand, days to 50% flowering, flag leaf length, days to maturity, panicle length, spikelet per panicle, number of filled grains, and dry matter showed positive significant correlations [52,53]. 1000-seed weight exhibited significant positive associations with flag leaf length, panicle length, spikelet per panicle, number of filled grains, harvest index, and yield per plant, while plant height and dry matter showed negative correlations at the genotypic level. Dry matter showed negative and significant associations with panicles per plant, 1000-seed weight, harvest index, and yield per plant. On the other hand, plant height, days to 50% flowering, flag leaf length, flag leaf width, days to maturity, panicle length, spikelet per panicle, number of filled grains, and number of unfilled grain per panicle showed positive significant correlations [54,53]. Harvest index showed significant positive associations with panicles per plant, 1000-seed weight, and yield per plant. On the other hand, plant height, days to 50% flowering, days to maturity, and dry matter showed negative correlations at the genotypic level [49,50]. In conclusion, plant height, number of panicles per plant, and number of filled grains per panicle were the most important characters showing significant positive correlations with grain yield per plant. Therefore, selecting based on these traits can lead to improved grain yield in rice.

#### 4.2.3 Path co-efficient study

From the results of path analysis (Table 4), it was evident that direct effects contributed by 1000-seed wt., days to 50% flowering, spikelet panicle<sup>-1</sup> (0.241) were high indicating that among the component traits, these three characters contributed maximum for yield plant<sup>-1</sup>. The direct effects of 1000-seed wt., days to 50% flowering, spikelet panicle<sup>-1</sup> were higher than their respective correlation co-efficient but lower for

number of filled grains panicle<sup>-1</sup>. The indirect effect of the above three characters were not important. Thus, direct selection for these traits will be effective. Chaudhury and Das [54] observed "high positive direct effect of days to maturity towards grain yield". Li et al. [46] reported highest positive direct effect of grains panicle<sup>-1</sup> on yield plant<sup>-1</sup> followed by 1000-grain weight and number of panicle plant<sup>-1</sup>. Chaubey and Singh [49] observed maximum positive direct effect of number of panicles plant<sup>-1</sup> followed by plant height and 100-grain weight. Samonte et al. [55] reported major influence of grain number panicle<sup>-1</sup> on yield. Although the character, harvest index was positively correlated with yield and its direct effect on grain yield plant<sup>-1</sup> was negative (-0.549). It indicated that this character influenced grain yield by its indirect positive effects through days to maturity, 1000-grain weight, and panicle plant<sup>-1</sup>. Mehetrc et al. [23] reported negative direct effect of days to harvest index on grain yield. Plant height showed a low negative direct effect (-0.041) on grain yield plant<sup>-1</sup> but the correlation co-efficient was significantly negative. Kumar et al. [56] reported moderate direct effect of plant height grain yield. Flag leaf length (0.215), panicle plant<sup>-1</sup> (0.139) and flag leaf width (0.043) showed moderate to low direct effect as the correlation coefficient was moderate to low for these characters. Days to maturity (-0.429), panicle length (-0.342), no. of filled grain (-0.135), no. of unfilled grain panicle<sup>-1</sup> (-0.119), dry matter (-0.158) had considerable negative direct effect on grain yield plant<sup>-1</sup> and showed negligible indirect effect [57]. Therefore, the positive correlation between panicle length, no. of filled grain and grain yield plant<sup>-1</sup> may be explained by positive indirect effect via days to 50% flowering, harvest index and flag leaf width (cm) and spikelet panicle<sup>-1</sup> respectively. The residual effect was 0.496, indicating that the eight characters contributed 51% percent of the variability in yield hill<sup>-1</sup> studied in path analysis. The residual effect towards grain yield in this study may be due to a variety of reasons such as causal factors (characters) not studied and sampling errors etc. Das et al. [17] observed a residual effect of 0.364 indicating 64 percent of the variability in the yield plant<sup>-1</sup> contributed by 8 characters studied in path analysis. The correlation and path co-efficient studies revealed that number of filled grains panicle<sup>-1</sup>, 1000-grain weight, and days to maturity and days to 50% flowering are the most important yield components. Recent breeding research has also emphasized the importance of these characters [53,54].

## 5. CONCLUSION

The study found varying genotypes in height, flowering time, and leaf characteristics. Days to maturity and panicle length showed notable differences among genotypes. Yield plant<sup>-1</sup> and 100-grain weight displayed high variation, while plant height, panicles plant<sup>-1</sup>, filled grains panicle<sup>-1</sup>, spikelet panicle<sup>-1</sup>, and unfilled grains panicle<sup>-1</sup> showed moderate variation. Days to 50% flowering, days to maturity, number of panicles plant<sup>-1</sup>, panicle length, filled grains panicle<sup>-1</sup>, yield plant<sup>-1</sup>, and other traits exhibited high heritability and genetic advance. Positive correlations were observed between yield plant<sup>-1</sup> and certain traits like panicle plant<sup>-1</sup>, filled grains, 100-seed weight, and harvest index. Selecting traits such as 100-seed weight, days to 50% flowering, and spikelet panicle<sup>-1</sup> could effectively increase yield plant<sup>-1</sup>. Harvest index influenced yield indirectly through other traits. Path coefficient analysis revealed specific direct and indirect effects of traits on yield plant<sup>-1</sup>, with some traits showing negative direct effects and positive indirect effects. The present study suggested that prolonged maturity period, higher number of panicles plant<sup>-1</sup>, longer panicle length and higher number of filled grains panicle<sup>-1</sup> should be given the importance in selection of genotypes for higher grain yield in rice.

## COMPETING INTERESTS

Authors have declared that they have no known competing financial interests or non-financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## REFERENCES

- Hoffman MS. The world almanac and book of facts. An imprint of pharos books. A scrips Howard Company. 200 Park Avenue, New York. 1991;10166.
- Lu, Chang TT. Rice in its temporal and spatial perspective. In: B.S. Luh (ed) Rice production and Utilization. AVI, Davis, California. 1980;1-74.
- Yoshida S. Rice. In: IRRI, Los Banos, Philippines. 1983;103-12.
- Hossain M. Sustaining food security in Asia: Economic, social, and political aspects. In: Dowliing, N.G., Greenfield, S.M. and Fischer, K.S. (ed). Sustainability of rice in the global food system. IRRI. 1998;21-34.
- Rashid MM. Rice production in Bangladesh: Programmers, achievements, potentials and challenges. IRCN. 1994;43:9-18.
- IRRI. International Rice Research Institute, Annual Report for 1994. Intl. Rice Res. Inst. Los Banos, Laguna, Philippines. 1995;179-181.
- BARC. Bangladesh Agricultural Research Council, Rice the main staple food, Farmgate, Dhaka; 1983.
- BBS. Bangladesh Bureau of statistics, Statistical yearbook of Bangladesh, Ministry of Planning, Govt. of the People's Republic of Bangladesh. 1996;167-175.
- BBS. Bangladesh Bureau of Statistics, Statistical Pocket Book of Bangladesh, Stat. Div., Ministry of Planning, Govt. Peoples Republic of Bangladesh. Dhaka, Bangladesh. 1998;385.
- FAO: Food and Agriculture organization of the United Nations, FAO Quarterly Bulletin of Statistics, Rome, Italy. 1999;12(3/4).
- FAO. Food and Agriculture organization of the United Nations, FAO Statistical yearbook 47, Rome Italy; 2013.
- IRRI. International Rice Research Institute, Principles and practices of Rice Production. John wiley and Sons. Inc. New York. 1981;49.
- Karim A. Krishi Diary. AIS. Khamarbari, Farm gate, Dhaka. 1992;130.
- BBS. Bangladesh Bureau of statistics, monthly statistical bulletin, Stat. Div., Ministry of Planning, Govt. Peoples Republic of Bangladesh. Dhaka, Bangladesh. 1999;54-55.
- Patwary AK. Yield performance and stability of some promising advanced generation lines of rice under different growing conditions. Prog. Agric. 1991;2(1):65-70.
- Chowdhury MAZ, Jalauddin MJ, Pandit DB, Asham ATMB. Genetic parameters, interrelationship and path co-efficient analysis in soybean (*Glycine max* (L.) Merr), Bang. J.Pl. Breed. Genet. 1994; 7(2):61-64.
- Das RK, Islam MA, Howlader M, Ibrahim SM, Ahmed HU, Miah NM. Variability and genetic association in upland rice. Bang. J. Pl. Breed. Genet. 1992;5(1&2):51-56.
- Mishra R, Rao SK, Koutu GK. Genetic variability, correlation studies and their implication in selection of high yielding genotypes of chickpea. Ind. J. Agric. Res. 1988;22(1):51-57.

19. Saha AK, Haque E, Quader B, Hussain TZ, Miah NM. Correlation and path analysis of some yield contributing characters in some high yielding and local varieties of irrigated rice. *Bang J. Pl. Breed. Genet.* 1989;2(1&2):19-22.
20. Singh RK, Chaudhury BD. Biometrical methods in quantitative genetic analysis (Rev. Edition). Kalyani Publisher, New Delhi, India; 1985.
21. Maurya DM, Saugh SK, Singh RS. Genetic variability in 48 lowland rice cultivars of Uttar Pradesh, India. *IRRN.* 1986;(11):4-13.
22. Das RK, Miah NM. Genetic variability in root and shoot characters of selected rice genotypes. *IRRN.* 1991;16(5):5.
23. Mehetre SS, Mahajan CR, Patil PA, Lad SK, Dhumal PM. Variability, heritability, correlation, path analysis and genetic divergence studies in upland rice. *IRRN.* 1994;19(1):8.
24. Thiagarajan CP. Influence of flag leaf area of rice seed germinability and vigor. *IRRN.* 1989;14(5):9.
25. Bashar MR, Haque E, Das RK, Miah NM. Relationship of flag leaf arc to yield filling grains per panicle and panicle length in upland rice varieties. *IRRN.* 1991;16(2):12.
26. BARC. Bangladesh Agricultural Research Council, Rice the main staple food, Farmgate, Dhaka; 1989.
27. Singh RS, Chauhan SP, Maurya DM. Genetic variability in 98 upland rice cultivars of India. *IRRN.* 1986;11(4):9-10.
28. Steel RGD, Torrie JH. Principles and procedures of statistics. McGraw Hill Book Co. Inc. New York. 1960;107-109.
29. Johnson KF, Robinson HF, Comstock RE. Genotypic and phenotypic correlation in soybeans and their implication in selection. *Agron. J.* 1955;47(10):477-483.
30. Burton GM. Quantitative inheritance in Grass pea Proc. of 6<sup>th</sup> Int. Grassland Cong. 1952;1:277-283.
31. Singh SK, Singh CM, Lal GM. Assessment of genetic variability for yield and its component characters in rice (*Oryza sativa* L.). *Research in Plant Biology.* 2011;1(4).
32. Hanson CH, Robinson HF, Comstock RE. Biometrics studies of yield in segregating population in Korean lespedeza. *Agron. J.* 1956;48:268-272.
33. Comstock RE, Robinson HF. Genetic parameters their estimation and significance. Proc. of 6<sup>th</sup> Int. Grassland Cong. 1952;1:128-291.
34. Miller PA, Williams JC, Robinson HF, Comstock RE. Estimation of genetic and environmental variance and covariance and their implication in selection. *Agron. J.* 1958;50:126-131.
35. Dabholkar AR. Elements of biometrical genetics. Concept Publishing Company. New Delhi, India; 1992.
36. Dewey DR, Lu KH. A correlation and path co-efficient analysis of components of crested wheat seed production *Agron. J.* 1959;51:515-518.
37. Mohammad T, Awan MA, Chcema AA. Interrelationship between yield and yield components in rice (*Oryza sativa* L). *Pak. J. Bot.* 1980;12(2):173-179.
38. Cheema AA, Yousaf A, Awan MA, Tahir GR. Path analysis of yield components of some mutants of basmati rice. *Tropical Agril. Res. and Extension (Pakistan).* 1998; 1(1):34-38.
39. Chauhan JS, Chauhan VS, Variar M. Genetic variation and character association in rainfed upland rice. *Oryza.* 1993;30:116-119.
40. Sarma MK, Richharia AK, Agarwal RK. Variability, heritability, genetic advance, and genetic divergence in upland rice. *IRRN.* 1996;21(1):25-26.
41. Rao SS, Shrivastav MN. Genetic variation and correlation studies in rainfed upland rice. *Oryza.* 1994;31:288-291
42. Yadav RK. Genetic Variability, correlation studies and their implication in selection of high yielding genotypes of rice. *Adv. Pl. Sci.* 1992;5:306-312.
43. Gupta A, Sharma RK, Mani VP, Chauhan VS. Variability and association analysis for gram yield and its components in hill rice. *Indian J Hill Research.* 1999;12(2): 99-104.
44. Chauhan SP, Singh RS, Maurya DM, Vaish CP. Character association in upland rice cultivars of India. *IRRN.* 1986;11(4):8.
45. Sawant DS, Patil SL, Sodaye VG. Genetic variability and heritability in segregating generation of rice crosses. *Ann. Agril. Res.* 1995;16(2):201-205.
46. Li QL, Li GT, Guo GZ, Jiang G, Zhu U. Study on high yield breeding and genetic analysis of yield components of main rice cultivars in Jilin, China. *Hereditas.* 1991;13(5):3-6.
47. Das SK, Singh J, Tripathy M, Mishra D. Association of quantitative traits and path analysis in medium land rice. *Env. And Ecol.* 1996;14(1):99-102.

48. Kumar GS, Mahadevappa M, Rudraradhya M. Studies on genetic variability, correlation and path analysis in rice during winter across the locations. *Karnataka J. Agril. Sci.* 1998;11(1):73-77.
49. Chaubey PK, Singh RP. Genetic variability, correlation and path analysis of yield components of rice. *Madras Agril. J.* 1994;81(9):464-470.
50. Mirza MJ, Faiz FA, Majid A. Correlation studies and path analysis of plant height, yield and yield components in rice *Sarhad. J. Agril.* 1992;8(6):647-653.
51. Shaha Roy PK, Nahar K, Ahmed HU, Mia NM, Islam MA. Genetic variability and character association in irrigated rice. *Bang. J. Pl. Breed. Genet.* 1993;6(1):69-74.
52. Chaudhury D, Rao MJBK, Prasad AB, Rao AVS, Suriya AV. Heritability and correlation in rice. *Oryza.* 1980;17(3):194-199.
53. Debi BR, Ray PKS, Howladar M. Variability and genetic association in irrigated rice. *Prog. Agric.* 1997;8(1&2): 165-168.
54. Chaudhury PKD and Das PK. Genetic variability, correlation and path co-efficient analysis in deep-water rice. *Ann. Agril. Res.* 1998;19(2):120-124.
55. Samonte SOPB, Wilson LT, McClung AM. Path analysis of yield and yield related traits of fifteen diverse rice genotypes. *Crop Science.* 1998;38(5):1130-1136.
56. Kumar R, Krishanpal M, Mondal SK, Ramashankar R, Prasad SC, Rai R. Genetic studies of major characters in upland rice. *Env. And Ecol.* 1994; 12(2):363-365.
57. Borbora TK, Hazarika GN. Study of genetic variability, heritability and genetic advance for panicle characters in rice. *Oryza.* 1999;35(1):19-21.

© 2023 Shahin et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/104215>