



Effect of Hormones on Rooting of Apical Cuttings for Propagation of Guava Cv. Lucknow-49

S. Yogesh ^{a++}, P. Harikanth ^{b#*}, T. Suresh Kumar ^{bt}
and B. Madhavi ^{b#}

^a Department of Fruit Science, College of Horticulture, Rajendranagar, Sri Konda Laxman Telangana State Horticultural University, Mulugu, Telangana- 502279, India.

^b Fruit Research Station, Sangareddy, Sri Konda Laxman Telangana State Horticultural University, Telangana, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i113447

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108923>

Original Research Article

Received: 24/08/2023

Accepted: 31/10/2023

Published: 11/11/2023

ABSTRACT

An attempt was made on the propagation of Guava cv. Lucknow-49 during the period of 2022-23 from apical cuttings under mist chamber conditions was undertaken. The apical cuttings were collected from shoot tips measuring about 10 to 15 cm treated with different hormones of concentrations viz., IAA @ 2000, 3000 and 4000 ppm; NAA @ 2000, 3000 and 4000 ppm along with IBA @ 3000, 4000 and 5000 ppm and these were compared with control, the treated apical cuttings were planted in 40 cavity trays and kept till 30 days for root induction under mist conditions. The different growth hormonal applications and its varied concentrations exerted a significant effect on root development in cuttings. Apical cuttings treated with IBA @ 5000 ppm was

⁺⁺ PG Scholar;

[#] Scientist;

[†] Principal Scientist (Hort.) & Head;

*Corresponding author: E-mail: harikanthporika@gmail.com;

found better for rooting percentage (69.90 %), survival percentage (67.50 %), time taken to root (23.75 days), number of roots per cutting (15.15, 16.25, 17.53 and 24.64) and root length (7.33, 9.78, 13.85 and 18.33 cm) respectively at 30, 60, 90 and 120 DAP, fresh root weight (3.93 and 10.15 g), dry root weight (1.13 and 2.18 g) at 60 and 120 DAP, respectively. The rooting hormone IBA performed better in formation of rooting in apical cuttings when compared to IAA, NAA and untreated control. It can be revealed from the current study that propagation of guava through apical cuttings when treated with IBA @ 5000 ppm was proved as a better propagation method for rooting of cuttings.

Keywords: Guava; propagation; cuttings; IBA; growth rate; mist chamber.

1. INTRODUCTION

Guava is a nutritious fruit that belongs to the family Myrtaceae. It is one of the most important fruit crop after Mango, Banana and Citrus. In India, it is cultivated in an area of 3.59 Lakh Ha with annual production of 55.90 Lakh MT [1]. In Telangana state, Guava stands as a major fruit crop with an area of 0.14 lakh acres and production of 1.17 Lakh MT [2]. There is a great demand for true to type and quality planting material [3] from the growers where nursery people cannot meet the requirement. Vegetative propagation by ground layering or stooling is a commercial practice to ensure true to type and early bearing of fruits but, in asexual method of propagation, soil media may carry the nematodes along with plant materials. Hence, the production of nematode free planting material is essential for the sustainable production of guava. Commercial methods of propagation such as air layering, grafting or stooling cannot meet the requirement of planting stock. Therefore, there is an urgent need to develop low cost protocol; fast and can provide high quality genetically true to type planting material. In this regard, a rapid method of propagation becomes an important when planting material supply is limited. Apical/terminal cutting is an easy, quick, novel and economical method of propagation.

Production of any crop is influenced mainly by the use of quality planting material [4]. External and internal factors affect the rooting of cuttings where the hormonal treatment is an exogenous factor to enhance the initiation of root primordia and growth through cell multiplication [5]. Hormones support mobilization of sugars and nutrients by the hydrolysis of starch to the base of cuttings [6]. Most of the time auxins like Indole acetic acid (IAA), Indole Butyric Acid (IBA), Naphthalene Acetic Acid (NAA) are used for rooting of cuttings. The present experiment was formulated to standardize the technique of propagation method in Guava cv. Lucknow-49 to

multiply a good and healthy root system through these cuttings under controlled mist chamber with the use of different hormonal and its varied concentrations [7].

2. MATERIALS AND METHODS

The present experimentation entitled "Effect of hormones on rooting of cuttings for propagation of guava cv. Lucknow-49" was carried out under controlled mist chamber during the period of 2022-23. The experiment was laid out in Completely Randomized Design with 10 treatments replicated four times. The apical cuttings were collected during the month of January and around 40 terminal cuttings per treatment were used for the trial. The tender, terminal portion of immature cuttings were taken from 8 years old mother plants of uniform size measuring a length of 10-12 cm, consisting of 2-3 nodes. These terminal cuttings were treated with different concentrations of IAA @ 2000, 3000 and 4000 ppm and NAA @ 2000, 3000 and 4000 ppm; IBA @ 3000, 4000 and 5000 ppm by quick dip method in their respective hormonal concentrations and compared with untreated control and later these cuttings were immediately planted in 40 celled cavity of portray measuring 3.5 cm top diameter, 3 cm bottom diameter and 8 cm depth and were transferred to mist chamber by maintaining a temperature of 25°C and 90% RH till first 30 days of rooting of cuttings (Plate 1). Irrigation applied with the help of foggers in the form of mist at an interval of every 20 minutes for 30 seconds till 30 days to maintain moist for rooting of cuttings. Before planting, these cuttings were immersed in 1% Bavistin solution to avoid any fungal attack. The root formation was observed during the mist chamber and shoot formation in the shade net conditions. After 30 days of planting in the mist chamber conditions, these portrays with rooted cuttings were transferred for further study under partial shade net (50% shade net) for 15-20 day more for shoot initiation. The data recorded were

statistically analysed by adopting procedure as suggested by Yates et al. [8].

For recording the observations about 4 cuttings were selected randomly in each replication per each treatment. The rooting percentage in each treatment was recorded at every 30 days after planting (DAP) and it was determined by counting the number of the rooted cuttings per replication to the total number of cuttings per replication and expressed in percentage.

Rooting percentage (%) = Number of the rooted cuttings per replication / Total number of for replication *100

Survival percentage of rooted cuttings was calculated at 60, 90 and 120 days after planting (DAP) by following formulae. Survival percentage of cuttings = $T-Mp / T \times 100$ were, T: Total number of rooted cuttings, Mp: Mortal plants

The apical cuttings were picked randomly within the treatment from each protray after 20 DAP under mist chamber conditions to observe the days taken to root till transferring to 50% shade net and the mean value was calculated and expressed in days. The number of roots per cutting was recorded from randomly selected 4 cuttings per treatment at 30, 60 and 90 days and average values were drawn. The root length from collar region to the tip of the root was measured for 4 randomly uprooted plants in each treatment in a centimetre scale at 30, 60 and 90 DAP and then average was calculated. The fresh root weight of cutting was recorded at 30, 60 and 90 DAP with the help of electronic weighing balance and the average value was computed. The same samples were kept in hot air oven at 60°C for 24 hrs to arrive a constant dry weight with the help of electronic weighing balance and the average value was computed and expressed in grams.

3. RESULTS AND DISCUSSION

3.1 Rooting Percentage

The method of propagation of guava through terminal cuttings, and its varied hormonal concentrations significantly influenced rooting of cuttings under mist chamber and the data obtained are presented in Table 1 and Fig. 1. The rooting percentage resulted significantly with their respective hormones and its concentrations. The maximum rooting percentage was recorded in IBA @ 5000 ppm (69.90 %) and it was

followed by IBA @ 4000 ppm (65.85 %), while the minimum registered (46.00 %) in untreated control. The exogenous application of IBA can synthesise auxins and cytokinins for the initiation of root and shoot in the selected softwood cuttings. Auxins are synthesized in apical shoots and then translocated downwards from cell to cell in a polar fashion, with a basipetal polarity in stems [9]. The possible reason for the lower success rate in untreated control might be due to the presence of higher polyphenols which acts as auxin transport inhibitors [10].

3.2 Survival Percentage

The apical softwood cuttings treated with IBA @ 5000 ppm significantly influenced the survival percentage of 67.50% and it was followed by IBA @ 4000 ppm (64.73 %), while the minimum percentage (47.50 %) was observed in untreated control (Table 1) and (Fig. 1). The highest survival percentage might be due to the fact that higher number of roots and root length influenced the better uptake of water and nutrients [11]. The translocation of photo assimilates synthesized in a newly formed shoots towards the root system helps in better survival of apical rooted cuttings. These results are in agreement with the earlier findings of Shukla and Bist [12] in Pear, Lakhani and Gajapara [13] in Guava Srivastava et al. [14] in Kiwi fruit, Saed [15] in Pomegranate and Rahad et al. [16]; Ali et al. [17] in Dragon fruit.

3.3 Days taken for Root Initiation

The cuttings treated with different hormones and its varied concentrations significantly differed days taken for root initiation, root length and no. of roots per cutting and obtained mean data are represented in Table 1.

Among the different growth hormones used, IBA decreased days taken for rooting of cuttings. The terminal cuttings dipped in 5000 ppm IBA resulted early rooting (23.75 days) and it was followed by IBA @ 4000 ppm (24.53 days), while the maximum days (30.32) observed in untreated control. The better utilization of reserve carbohydrates, nitrogen and other factors stored in stem cuttings activated hydrolytic enzymes with IBA application [18]. The rooting in terminal cuttings might be due to availability of auxins and cytokinins with IBA and these apical cuttings are one of the sites of natural auxin synthesis [19]. The minimum days to root (30.32 days) was observed in untreated control which might be due to the over absorption of auxins, causing

inhibitory action on rooting. Temperature, relative humidity and media played a significant role in realizing better success rate [20]. Similar findings were reported earlier by Srivastava et al. [14] in Kiwi fruit and Muhammad et al. [21] in Ber.

3.4 Number of Roots Per Cuttings

The data presented in the Table 2, indicated the number of roots per cutting significantly influenced with different hormones and its concentrations at 30, 60, 90 and 120 DAP. The maximum number of roots per cutting (15.15, 16.25, 17.53 and 24.64) was recorded in IBA @ 5000 ppm and it was followed by 4000 ppm IBA (13.75, 15.25, 16.51 and 22.04) whereas, the minimum number of roots per cutting was recorded in untreated control (6.10, 8.57, 10.25 and 14.74) at 30, 60, 90 and 120 DAP, respectively. The maximum number of roots might be due to hormonal effect leading to accumulation of internal substances and their downward movement as well as cell division, enhanced better expression of root primordial that resulted in significant development of root system. These findings are in accordance with the results reported earlier by Noor et al. [22] in Litchi, Athani et al. [23] in Guava, Kumar and Syamal (2005) in Guava, Maurya et al. [24] in Guava and Bhosale et al. [25] in Pomegranate.

3.5 Root Length (cm)

The root length parameter significantly differed with different hormones and its concentrations at 30, 60, 90 and 120 DAP (Table 2). Among the different treatments evaluated, the maximum root length (7.33, 9.78, 13.85 and 18.33 cm) was

observed in 5000 ppm IBA at 30, 60, 90 and 120 DAP, respectively, while the minimum root length was recorded in control. The IBA @ 5000 ppm initiated better root length which could be due to hydrolysis of polysaccharides stored in these cuttings into physiologically active sugars, provides energy through respiratory activity to the root primordia that helped in rapid elongation of meristematic cells to initiate better root length [26]. Similar results were reported by Kumar et al. [27] in Guava, Reddy et al. [28] in Fig, Gohil [29] in Cashewnut and Tomar and Tomar [30] in Pomegranate.

3.6 Fresh and Dry Root Weight (g)

The fresh root weight (3.93 and 10.15 g) and dry root weight (1.13 and 2.18 g) resulted significant with different growth hormones and its varied concentrations at 60 and 120 DAP in IBA @ 5000 ppm while, the minimum fresh root weight (1.79 and 4.85 g) and dry root weight (0.48 and 0.85 g) was recorded in untreated control at 60 and 120 DAP, respectively (Table 1). The higher fresh and dry root weight could be attributed to the rapid hydrolysis of polysaccharides stored in the cuttings into physiologically active sugars by activation of hydrolytic enzymes. These sugars provide energy for the meristematic tissue and root primordial through respiratory activity leads to initiate a greater number of adventitious roots and cell elongation which helps in establishment of higher root weight per cutting. The present findings are in agreement with the results of Singh et al. [31] in Lemon, Porghorban et al. [32] in Olive, Rahad et al. [16]; Ali et al. [17] in Dragon fruit.




Fig. 1. Effect of hormones on rooting and survival percentage of rooted cuttings in Guava cv. Lucknow-49



Plate 1. Procedure and steps in preparation of apical cuttings for propagation of Guava cv. Lucknow-49

a) Guava mother block; b) Selected terminal shoots for preparation of softwood cutting; c) Prepared softwood cuttings d) Cuttings treated with 1% Carbendazim to prevent any fungal disease; e) Treated cuttings planted in portrays; f) Pro-trays kept under mist chamber for rooting; g) Rooted cuttings; h) Clones produced by apical softwood cuttings after 120 DAP

Table 1. Effect of hormones on rooting percentage, days taken to root, survival percentage, fresh and dry root weight in propagation of guava apical cuttings cv. Lucknow-49

Treatments	Rooting percentage	Days taken for root initiation	Survival percentage	Fresh root weight (g)		Dry root weight (g)	
				60 DAP	120 DAP	60 DAP	120 DAP
IAA @ 2000 ppm	52.40	28.79	52.13	2.46	6.74	0.68	1.12
IAA @ 3000 ppm	55.65	26.77	56.26	2.74	8.16	0.72	1.29
IAA @ 4000 ppm	57.50	25.42	59.53	2.82	8.68	0.78	1.42
IBA @ 3000 ppm	58.65	24.79	61.62	2.89	9.25	0.85	1.57
IBA @ 4000 ppm	65.85	24.53	64.73	3.03	9.47	0.97	1.87
IBA @ 5000 ppm	69.90	23.75	67.50	3.93	10.15	1.13	2.18
NAA @ 2000 ppm	49.40	29.25	49.34	2.29	5.98	0.56	1.05
NAA @ 3000 ppm	51.28	28.91	51.22	2.41	6.34	0.65	1.13
NAA @ 4000 ppm	53.53	27.25	53.48	2.68	7.57	0.70	1.19
Control	46.00	30.32	47.50	1.79	4.85	0.48	0.85
S.Em ±	0.811	0.471	1.170	0.041	0.162	0.009	0.021
CD at 5%	2.351	1.372	3.391	0.133	0.471	0.026	0.062

Table 2. Effect of hormones on number of roots/cutting and root length at 30, 60, 90 and 120 DAP in propagation of guava apical cuttings cv. Lucknow-49

Treatments	No. of roots/cutting				Root Length (cm)			
	30 DAP	60 DAP	90 DAP	120 DAP	30 DAP	60 DAP	90 DAP	120 DAP
IAA @ 2000 ppm	8.50	10.53	12.56	18.58	5.25	6.49	10.74	15.88
IAA @ 3000 ppm	9.60	10.75	13.53	20.52	5.45	6.88	11.36	16.53
IAA @ 4000 ppm	10.13	12.69	14.65	21.58	5.65	7.08	11.55	16.78
IBA @ 3000 ppm	10.91	13.35	15.65	21.87	5.83	7.15	11.85	16.92
IBA @ 4000 ppm	13.75	15.25	16.51	22.04	6.93	8.18	12.35	17.58
IBA @ 5000 ppm	15.15	16.25	17.53	24.64	7.33	9.78	13.85	18.33
NAA @ 2000 ppm	8.03	10.12	11.75	16.96	5.11	6.27	10.42	15.18
NAA @ 3000 ppm	8.35	10.25	12.38	17.81	5.15	6.35	10.68	15.52
NAA @ 4000 ppm	9.25	10.69	12.82	19.65	5.33	6.63	10.89	16.38
Control	6.10	8.57	10.25	14.74	4.73	5.88	9.17	12.13
S.Em ±	0.149	0.161	0.296	0.309	0.092	0.121	0.218	0.233
CD at 5%	0.430	0.465	0.857	0.893	0.266	0.351	0.631	0.673

4. CONCLUSION

It could be finally concluded from the present investigation that the effect of growth hormones on propagation of guava cv. Lucknow 49 through apical cuttings, the different rooting hormones and its concentrations exerted a significant effect on root development in apical cuttings. Among the various treatments IBA @ 5000 ppm could found better when compared to other treatmental effect in respect to various attributes viz., rooting and survival percentage, time taken for root formation, number of roots and its root length per cutting at 30, 60, 90 and 120 DAP, respectively, similarly fresh and dry root weight at 60 and 120 DAP, respectively recorded higher when treated with 5000 ppm IBA. The hormonal treatment with IBA @ 5000 ppm performed better in propagation of apical cuttings when compared to IAA, NAA and untreated control. This experiment can reveal the potentiality of guava propagation through terminal or apical cuttings with IBA @ 5000 ppm and it will be a novel, economical method to propagate for mass production.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Area and production of horticulture crops for 2022-23 in India. Department of agriculture and farmers welfare, New Delhi; 2023. Available: <https://agricoop.gov.in/en/StatHortEst>
2. Anonymous. Area and production of horticulture crops for 2021-22 in Telangana state. Department of horticulture, Telangana, India; 2022. Available: <https://horticulture.tg.nic.in>
3. Amit V, Singh JN, Surendra P Singh. Standardization of wedge grafting in Guava under 11. North Indian plains. Indian Journal of Horticulture. 2010;67(Special issue):111-114.
4. Ambebe TF, Agbor AEW, Siohdjie CS. Effect of different growth media on sprouting and early growth of cutting propagated *Cordia Africana* (Lam.). International Journal of Forest, Animal and Fisheries Research. 2018;2(1):28-33.
5. Fogasa CM, Fett-Neto AG. Role of auxin and its modulators in the adventitious rooting of Eucalyptus species differing in recalcitrance. Plant Growth Regulation. 2005;4(1):1–10.
6. Das P, Basak U, DAP A. Metabolic changes during rooting in pre-girdled stem cuttings and air layers of Heritiera. Botanical Bulletin of Academia Sinica. 1997;38:91–95.
7. Davies FTJ, Hartman HT. The physiological basis of adventitious root formation. Acta Horticulturae. 1988;227: 113-120.
8. Yates F, Mather K. Ronald Aylmer Fisher, 1890-1962; 1963.
9. Lomax TL, Muday GK, Rubery P. Auxin transport. In PJ Davies, ed, Plant hormones: Physiology, biochemistry, and molecular biology. Kluwer Academy Press, Norwell, The Netherlands. 1995;509-530.
10. Brown DE, Rashotte AM, Murphy AS, Normanly J, Tague BW, Peer WA, Taiz L, Muday GK. Flavonoids act as negative regulators of auxin transport *in vivo* in Arabidopsis. Plant Physiology. 2001;126:524–535.
11. Reddy RKV, Pulla RC, Goud VR. Role of auxins synergists in the rooting of hardwood and semi hardwood cuttings of Fig (*Ficus carica* L.). Indian Journal of Agricultural Research. 2008;42(1):75-78.
12. Shukla GS, Bist LD. Studies on the efficacy of IBA and NAA on clonal propagation by cutting in low chilling Pear rootstocks. Indian Journal of Horticulture. 1994;51(4):351-357.
13. Lakhani RN, Gajapara NN. Effect of plant growth substances on rooting and survival of stem cuttings of Pomegranate (*Punica granatum*) var. Ganesh and Dholka. Journal of Applied Horticulture. 1998;4(1-2):52.
14. Srivastava K, Biswajit DK, Bhatt KM. Effect of Indole Butyric Acid and variety on rooting of leafless cutting of Kiwi fruit under zero energy-humidity-chamber. ENVIS Bulletin: Himalayan Ecology. 2005;14(1): 31-34.
15. Saed JO. Rooting response of five pomegranate varieties to IBA concentrations and cutting age. Pakistan Journal of Biological Sciences. 2010; 13(2):51-58.
16. Rahad MK, Islam MA, Rahim MA, Monira S. Effects of rooting media and varieties on rooting performance of Dragon fruit

- cuttings (*Hylocereus undatus* Haw.). Agriculture, Livestock and Fisheries an open Access Peer Reviewed Journal. 2016;3(1):67-77.
17. Ali SI, Kumar TS, Kumar AK, Joshi V, Kumar BN. Studies on effect of different concentrations of IBA and length of cuttings on rooting and shoot growth performance in Dragon fruit *Hylocereus* spp.-red flesh with pink skin under Telangana conditions. The Pharma Innovation Journal. 2022;11(3):738-743.
 18. Chandramouli H. Influence of growth regulators on the rooting of different types of cuttings in *Bursera pennisiliata*. M.Sc., Thesis. University of Agricultural Sciences, Bangalore, India; 2001.
 19. Wahab F, Ghulam Nabi, Nawab Ali, Muslim Shah. Rooting Response of semi hardwood cuttings of Guava (*Psidium guajava* L.) to various concentration of different Auxins. Journal of Biological Sciences. 2001;1(4):184-187.
 20. Rymbai H, Reddy GS. Effect of IBA, time of layering and rooting media on air layers and plantlets survival under different growing nursery conditions in guava. Indian Journal of Horticulture. 2010;67:99-104.
 21. Muhammad AB, Muhammad AA, Zubeda C, Hamid R. Response of Jojoba (*Simmondsia chinensis*) cuttings to various concentrations of auxins. Pakistan Journal of Botany. 2009;41(6):2831-2840.
 22. Noor R, Ijaz H, Imran Md, Taslim Jan, Azmat AA. Effect of different concentration of IBA on rooting of Litchi (*Litchi chinensis*) in Air layering. Pakistan Journal of Biological Sciences. 2000;3(2):330-331.
 23. Athani SI, Swamy GSK, Patil PB. Effect of different pre-treatments on rooting of air layers in guava cv. Sardar. Karnataka Journal of Agricultural Science. 2001;14(1):199-200.
 24. Maurya RK, Ray NR, Chawda JC, Chauhan VB, Patil AK. Evaluation of different organic media and water holding materials with IBA on rooting and survival of air layering in guava (*Psidium guajava* L.) cv. Allahabad safeda. The Asian Journal of Horticulture. 2012;7(1):44-47.
 25. Bhosale VP, Shinde SM, Turkhade PD, Dekhmukh SB, Sawant SN. Studied the response of different media and PGR'S on rooting and survival of air layers in Pomegranate cv. Sindhuri. Annals of Horticulture. 2014;7(1):73-77.
 26. Singh KK, Choudhary T, Kumar A. Effect of various concentrations of IBA and NAA on the rooting of stem cuttings of mulberry (*Morus alba* L.) under mist house condition in Garhwal hill region. Indian Journal of Hill Farming. 2014;27(1):74-77.
 27. Kumar K, Aulakh PS, Baidwan RPS. Standardization of time of budding in guava (*Psidium guajava* L.) under lower Shivaliks conditions of Punjab. Haryana Journal of Horticultural Sciences. 2007;36(1-2):61-62.
 28. Reddy PPN, Ray R, Patel AD, Patel JS. Effect of rooting media and IBA (Indole butyric acid) levels on rooting and survival of Air layering in Fig (*Ficus carica* L.) cv. Poona under middle Gujarat agro-climatic conditions. The Asian Journal of Horticulture. 2014;9(1):1-5.
 29. Gohil JH. Effect of Different Concentrations of IBA and NAA on Air Layering of Cashew nut cv. Vengurla 4. Trends in Biosciences. 2014;7(5):351-354.
 30. Tomar KS, Tomar SS. Effect different dosage of growth regulators on rooting and survival percentage (*Punica granatum* L.) of Pomegranate air layers. Green Farming. 2012;3(6):701-703.
 31. Singh KK, Choudhary T, Kumar P. Effect of IBA concentrations on growth and rooting of Citrus Limon cv. Pant lemon cuttings. Hort Flora Research Spectrum. 2013; 2(3):268-270.
 32. Porghorban M, Moghadam EG, Asgharzadeh A. Effect of media and Indole Butyric Acid (IBA) concentrations on rooting of Russian Olive (*Elaeagnus angustifolia* L.) semi-hard wood cuttings. Indian Journal of Fundamental and Applied Life Sciences. 2014;4(3):517-522.

© 2023 Yogesh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/108923>