



# **Developing Biocontrol Strategies for the Management of Set rot Disease of Sugarcane under Wider Row Planting**

**K. Yamunarani <sup>a</sup>, S. Sangeetha <sup>a\*</sup>, K. Chitra <sup>b</sup>,  
K. Dhanalakshmi <sup>a</sup> and V. Dhanushkodi <sup>c</sup>**

<sup>a</sup> Horticultural College and Research Institute for Women, Tiruchirappalli, Tamil Nadu, India.

<sup>b</sup> ICAR-Krishi Vigyan Kendra, Virinchipuram, Vellore, India.

<sup>c</sup> ICAR-Krishi Vigyan Kendra, Needamangalam, Thiruvavur, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/JSRR/2023/v29i121828

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112081>

**Original Research Article**

**Received: 25/10/2023**

**Accepted: 29/12/2023**

**Published: 30/12/2023**

## **ABSTRACT**

Sugarcane (*Saccharum officinarum* L.) is one of the economically valuable agricultural crops grown worldwide in tropical and subtropical areas mainly for their sugar source. Among the 110 sugarcane cultivated countries, India and Brazil contribute half of global production [1]. During the period of last five decades the production, productivity and sugarcane recovery has also shown remarkable resilience in productivity growth rate. One of the reasons for the low productivity and recovery is increasing incidence of insect pests and diseases. An ecologically friendly alternative to these problems is biological control using rhizobacteria and their metabolic products [2]. In order to develop a technology capsule Surveys and surveillance were conducted in sugarcane fields both in plant and ratoon crops distributed in Orthanadu and Vallam block in New Cauvery delta area of Thanjavur district during three consecutive years 2019- 22 with an objective to assess the incidence of sett rot, red rot and smut in New Cauvery delta area of Thanjavur district. Fortnightly fixed plot

\*Corresponding author: E-mail: [sangeetha.s@tnau.ac.in](mailto:sangeetha.s@tnau.ac.in);

observation was taken. The Maximum and minimum temperature, RH and quantity of rain and rainy days also recorded daily and consolidated on monthly basis and weather parameters were correlated with the incidence of disease. The results revealed that the maximum incidence sett rot negatively correlated with temperature. Similarly the number of rainy days and rainfall is positively correlated ( $r = 0.7523$ ) with incidence of sett rot. For managing sett rot disease of sugarcane through biological approach, studies were conducted on biocontrol agents viz., *Trichoderma asperellum* (TNAU), *Bacillus subtilis* (TNAU), *Chaetomium globosum* (Cg6) with fungicide check propiconazole 0.1%. Among the treatment, highest millable cane (91,080/ha) and yield (91.39 tonnes / ha) and sugar yield ( 10.63 tonnes/ha) were recorded in M1- (Single bud treated with *Trichoderma asperellum* (TNAU)@ 4 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>) followed by M<sub>2</sub>. (Single bud treated with *Bacillus subtilis* (TNAU)@ 10 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>) in which, the cane yield of 88.54 tonnes/ha and sugar yield of 10.68 tonnes/ha were recorded. This study provides a technology capsule for the management of important disease of sugarcane.

**Keywords:** Sugarcane; sett rot; biological control.

## 1. INTRODUCTION

Sugarcane is one of the important commercial crops in tropics and serves as the main source of sugar in the world. In Tamil nadu, sugarcane is cultivated in 3.20 lakhs hectares with a production of about 350 lakhs tonnes of canes. Productivity of Tamil Nadu is higher (108 t/ha) than the national average [3,4]. Sugarcane red rot caused by *Colletotrichum falcatum* and smut disease caused by *Sporisorium scitamineum* is a devastating disease in sugarcane growing areas globally, which results in considerable loss of sugar yield [5]. Currently, control of sugarcane disease mainly relies on the breeding of resistant cultivars [6], which is constrained by long breeding processes, high cost, and availability of resistant parental lines. Rhizobacteria inhibit plant pathogens by numerous mechanisms related to: (i) secretion of metabolites like siderophores, which suppress pathogens by sequestering iron [7] (ii) hydrolytic enzymes, which degrade the cell wall of many pathogens [8]; (iii) antibiotics. Which induce systemic resistance in plants. Among the different diseases Sett rot or pineapple disease caused by *Ceratocystis paradoxa* (de Seynes) Moreau is both sett and soil borne disease of sugarcane affecting the sett germination at the early stages of planting. Three different biocontrol agents were evaluated in field condition for the management sett rot.

Introduction of wider row planting in sugarcane cultivation in Tamil Nadu has aimed to increase the cane yield in unit area and same time it provides best tool for applying plant protection chemicals to the targeted sites in sugarcane. Under wider row condition, the fertilizers are

being applied in heavy dose in multiple split till harvest which may alter the pests and disease occurrence and favour the pests and diseases population build up. As the wider row planting of sugarcane cultivation is being in advocacy to the farming community, it now becomes imperative that the pest and disease pattern in the changed scenario has to be studied in detail as the moisture pattern is known to influence the pest and disease behavior significantly in many crops. Hence the detailed studies on pests and diseases occurrence pattern under wider row sugarcane cultivation are warranted. In the present study periodical pest and disease surveillance was conducted to assess the incidence of different pests and diseases of sugarcane along with natural enemies in New Cauvery Delta region of Thanjavur District.

## 2. METHODOLOGY

The popular and high sugar sugarcane variety viz Co 86032 was raised under wider row with row spacing of 120 cm in red loam soil condition at Agricultural college and Research institute, Eachangkottai. The crop was raised with single bud setts in paired /wider rows spacing of 120 cm between rows following the recommended wider row package of practices. The treatments were imposed as per the technical programme.

**M<sub>1</sub>**.Single bud treated with *Trichoderma asperellum* (TNAU)@ 4 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>

**M<sub>2</sub>**.Single bud treated with *Bacillus subtilis* (TNAU)@ 10 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>

**M<sub>3</sub>**.Single bud treated with *Chaetomium globosum* (Cg6)@ 10 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>

**M<sub>4</sub>**.Single bud treated with Propiconazole @ 1 g l<sup>-1</sup> + soil drenching @ 0.1 %

**M<sub>5</sub>**.Untreated control

Observations viz., Germination percent on 30<sup>th</sup> day, Set rot incidence on 30<sup>th</sup> day, tiller populations were recorded on 90 days after planting and economic shoot populations were recorded on 210 days after planting.

### 3. RESULTS AND DISCUSSION

Fortnightly fixed plot observation was taken in cane crop planted under wider row planting in Orthanadu and Vallam block in New Cauvery delta area of Thanjavur district to assess the incidence of diseases viz., sett rot, red rot and smut. The Maximum and minimum temperature, RH and rainfall and rainy days also recorded daily and consolidated on monthly basis and

weather parameters were correlated and presented in Tables 1,2, 3 and 4.

#### 3.1 Effect of Biocontrol Treatments on Sett rot Disease and Yield Characters

The popular and high sugar sugarcane variety viz Co 86032 was raised under wider row at Agricultural college and Research institute, Eachangkottai during june 2019 and harvested during may 2020. The treatments were imposed and observations viz., Germination percent on 30<sup>th</sup> day, Set rot incidence on 30<sup>th</sup> day, Smut and red rot incidence at monthly intervals, Tillers population counts on 90<sup>th</sup> day, economic population counts on 210<sup>th</sup> day were recorded as per the technical programme. The observation viz., Set rot incidence, Smut and red rot incidence at monthly intervals were presented along with percent reduction and yield contributing characters in Tables 5 to 8.

**Table 1. Incidence of diseases of sugarcane in new cauvery delta of Thanjavur District. 2019 – 2020**

S.No.	Month	Maximum temp.	Minimum temp.	Average	RH %	No. of rainy days	Rain fall (mm)	Sett rot Incidence (%)
1	Jun 19	32.00	21.50	27.00	79.00	3	60.0	-
2	July 19	33.00	23.50	28.00	85.60	11	300.0	-
3	Aug 19	33.40	23.50	29.00	85.00	8	154.0	-
4	Sep 19	33.60	25.00	28.00	88.00	11	116.0	-
5	Oct 19	33.50	22.70	28.80	83.00	5	150.0	5.80
6	Nov 19	33.53	23.00	28.50	89.00	10	160.0	5.90
7	Dec 19	31.50	21.00	26.50	81.00	10	620.5	7.25
8	Jan 20	31.70	21.50	26.00	82.00	14	501.0	7.25
9	Feb 20	31.00	22.50	27.00	82.00	1	0	-
10	Mar 20	32.00	23.60	28.00	84.00	0	0	-
11	April 20	31.00	23.50	27.50	84.00	0	7.0	-
12	May 20	35.50	23.50	29.50	79.00	1	8.0	-
Correlation Coefficient (r) with disease intensity			Max temp			-0.3146		
			Min temp			-0.6512		
			RH			-0.4611		
			No of rainy days			0.7412		
			Rain fall			0.6546		

**Table 2. Incidence of diseases of sugarcane in new cauvery delta of Thanjavur District. 2020 – 2021**

S.No.	Month	Maximum temp.	Minimum temp.	Average	RH%	No. of rainy days	Rain fall (mm)	Sett rot Incidence (%)
1	Mar 20	33.10	23.60	28.35	84.70	-	-	-
2	April 20	32.00	23.60	27.80	84.30	1	5.0	-
3	May 20	36.10	23.70	29.95	79.40	1	4.0	-
4	Jun 20	32.10	23.80	27.95	78.70	4	69.0	-
5	July 20	33.10	24.50	28.8	85.60	10	313.6	-
6	Aug 20	34.40	23.80	29.10	85.80	7	150.0	-
7	Sep 20	33.80	25.10	28.80	88.50	10	106.0	-
8	Oct 20	33.90	23.70	28.90	83.50	6	152.0	4.86
9	Nov 20	33.4	23.3	28.35	89.8	13	162.0	5.45
10	Dec 20	31.2	21.5	26.35	80.0	12	625.5	6.66
11	Jan 21	30.7	21.4	26.05	83.0	11	506.0	6.62
12	Feb 21	31.9	22.7	27.3	82.2	-	0	-
Correlation Coefficient (r) with disease intensity				Max temp		-0.4136		
				Min temp		-0.7105		
				RH		-0.5709		
				No of rainy days		0.7503		
				Rain fall		0.7523		

**Table 3. Incidence of diseases of sugarcane in New Cauvery Delta of Thanjavur District .2021 – 2022**

S.No.	Month	Maximum temp.	Minimum temp.	Average	RH %	No. of rainy days	Rain fall (mm)	Sett rot incidence (%)
1	Mar 21	34.00	24.00	29.00	80.29	0	0	-
2	April 21	36.51	25.08	30.79	75.43	1	23	-
3	May 21	36.45	25.12	30.78	71.87	2	45	-
4	Jun 21	36.40	25.35	30.87	73.03	2	11	-
5	July 21	35.70	24.50	30.25	78.54	2	28.2	-
6	Aug 21	34.41	23.85	29.13	82.29	7	238.0	-
7	Sep 21	34.36	24.31	29.33	83.96	10	258.0	-
8	Oct 21	33.67	24.32	28.99	84.54	6	371.0	10.70
9	Nov 21	34.43	23.73	29.08	83.56	13	628.5	12.20
10	Dec 21	34.56	23.67	23.61	88.58	12	12.5	7.20
11	Jan 22	34.01	24.01	29.01	87.80	11	62	5.62
12	Feb 22	33.91	23.83	28.87	85.78	0	31	-
13	Mar 22	34.50	23.33	28.91	79.08	0	0	-
					Max temp	--0.0485		
					Min temp	-0.6912		
					RH	-0.5171		
					No of rainy days	0.7123		
					Rain fall	0.6425		

**Table 4. Pooled mean analysis for Incidence of diseases of sugarcane in new cauvery delta region of Thanjavur District**

S.No.	Month	Maximum temp.	Minimum temp.	Average	RH %	No. of rainy days	Rain fall (mm)	Sett rot Incidence (%)
1	Jan	Jan	32.1	22.3	27.0	84.3	12.0	6.49
2	Feb	Feb	32.3	23.0	27.7	83.3	0.3	-
3	Mar	Mar	33.0	23.7	28.5	83.0	0.0	-
4	April	April	33.2	24.1	28.7	81.2	0.7	-
5	May	May	36.0	24.1	30.1	76.8	1.3	-
6	Jun	Jun	33.5	23.6	28.6	76.9	3.0	-
7	July	July	33.9	24.2	30.3	83.2	7.7	-
8	Aug	Aug	34.1	23.7	29.1	84.4	7.3	-
9	Sep	Sep	33.9	24.8	28.7	86.8	10.3	-
10	Oct	Oct	33.67	24.32	28.99	84.54	6.0	7.12
11	Nov	Nov	33.8	23.3	28.6	87.5	12.0	7.85
12	Dec	Dec	34.56	23.67	23.61	88.58	12	7.03
Correlation Coefficient (r) with disease intensity				Max temp		-0.0447		
				Min temp		-0.6843		
				RH		-0.5164		
				No of rainy days		0.7346		
				Rain fall		0.6831		

*The minimum temperature and higher relative humidity prevailing during October to January might have favoured the disease incidence which ultimately hindered the sett germination. Shanmugam et al. [14].*

**Table 5. Effects of Main treatment on disease and yield contributing parameters**

S.No.	Treatments	Germination %	Set rot %	Percent reduction over control	Millable cane (000 /ha)	Yield (t/ha)	Percent increase over control	CCS (%)	Sugar Yield (t/ha)
1	<b>M<sub>1</sub></b>	83.12 (66.87)	3.0 (11.44)	66.70	<b>91.08</b>	<b>91.39</b>	<b>10.41</b>	<b>12.2</b>	<b>10.63</b>
2	<b>M<sub>2</sub></b>	75.62 (60.47)	4.5 (12.22)	50.00	87.10	86.40	5.24	12.2	10.36
3	<b>M<sub>3</sub></b>	77.19 (61.67)	4.5 (12.22)	50.00	88.03	87.74	7.16	12.3	10.52
4	<b>M<sub>4</sub></b>	71.00 (57.45)	4.0 (11.15)	55.55	85.99	86.09	4.09	12.3	10.32
5	<b>M<sub>5</sub></b>	64.47 (53.54)	9.0 (17.43)	-	83.19	81.87	-	12.2	9.82
CD (P=0.05)		1.67	1.28	-	<b>3.88</b>	<b>1.97</b>		<b>NS</b>	

Among treatments, highest millable cane (91,080/ha) and yield (91.39 tonnes / ha) and sugar yield ( 10.63 tonnes/ha) were recorded in **M<sub>1</sub>**- (Single bud treated with *Trichoderma asperellum*(TNAU)@ 4 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>) followed by **M<sub>3</sub>**- (Single bud treated with *Chaetomium globosum* (Cg6)@ 10 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>) in which, the cane yield of 87.74 tonnes/ha and sugar yield of 10.52 tonnes/ha were recorded. There is no significant difference in CCS percent among the treatments.

**Table 6. Effect of biocontrol on disease and yield character (2020-2021)**

S.No.	Treatments	Germination %	Set rot %	Percent reduction over control	Millable cane (000 /ha)	Yield (t/ha)	Percent increase over control	CCS (%)	Sugar Yield (t/ha)
1	<b>M<sub>1</sub></b>	80.12 (63.52)	3.2 (10.30)	62.79	<b>92.70</b>	<b>92.86</b>	<b>10.70</b>	<b>12.31</b>	<b>10.74</b>
2	<b>M<sub>2</sub></b>	73.23 (58.84)	4.8 (12.65)	44.18	89.05	87.91	6.03	12.76	10.85
3	<b>M<sub>3</sub></b>	78.28 (62.22)	5.1 (13.05)	40.69	88.20	86.96	4.88	12.82	10.83
4	<b>M<sub>4</sub></b>	72.11 (58.12)	5.0 (12.92)	41.86	86.10	86.90	4.09	12.80	10.89
5	<b>M<sub>5</sub></b>	68.36 (55.77)	8.6 (17.05)	-	84.28	82.91	-	12.56	9.95
CD (P=0.05)		1.65	1.23	-	<b>3.78</b>	<b>1.95</b>		<b>NS</b>	

In Main treatment, highest millable cane (92,700 /ha) and yield (92.86 tonnes / ha) and sugar yield ( 10.70 tonnes/ha) were recorded in **M<sub>1</sub>**- (Single bud treated with *Trichoderma asperellum*(TNAU)@ 4 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>) followed by **M<sub>2</sub>**- (Single bud treated with *Bacillus subtilis* (TNAU)@ 10 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>) in which, the cane yield of 89.05 tonnes/ha and sugar yield of 10.85 tonnes/ha were recorded. The CCS percent among the treatment is also significantly different.

**Table 7. Effects of Main treatment on disease and yield contributing parameters**

S.No.	Treatments	Germination %	Set rot %	Percent reduction over control	Millable cane (000 /ha)	Yield (t/ha)	Percent increase over control	CCS (%)	Sugar Yield (t/ha)
1	<b>M<sub>1</sub></b>	85.12 (66.87)	6.5 (14.77)	<b>35</b>	<b>92.09</b>	<b>98.76</b>	<b>18.80</b>	<b>12.35 (20.58)</b>	<b>10.67</b>
2	<b>M<sub>2</sub></b>	83.62 (60.47)	7.0 (15.89)	30	89.23	96.53	16.70	12.33 (20.56)	10.23
3	<b>M<sub>3</sub></b>	81.19 (1.67)	8.0 (16.43)	20	87.19	88.64	9.43	12.18 (20.42)	9.66
4	<b>M<sub>4</sub></b>	81.00 (57.45)	7.2 (15.34)	28	85.71	86.09	5.90	12.20 (20.44)	9.85
5	<b>M<sub>5</sub></b>	74.47 (53.54)	10.0 (18.43)	-	84.19	81.70	-	12.17 (20.41)	9.61
CD (P=0.05)		2.15	2.09		1.91	1.75		2.07	0.19

*In Main treatment, highest millable cane (92,090 /ha) and yield (98.76 tonnes / ha) and sugar yield ( 10.67 tonnes/ha) were recorded in **M<sub>1</sub>**- (Single bud treated with *Trichoderma asperellum*(TNAU)@ 4 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>) followed by **M<sub>2</sub>**- (Single bud treated with *Bacillus subtilis* (TNAU)@ 10 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>) in which, the cane yield of 89.23 tonnes/ha and sugar yield of 10.23 tonnes/ha were recorded. The CCS percent among the treatment is also significantly different.*

**Table 8. Effect of main treatments on disease incidence and yield characters**

S.No.	Treatments	Germination %	Set rot %	Percent reduction over control	Millable cane (000 /ha)	Yield (t/ha)	Percent increase over control	CCS (%)	Sugar Yield (t/ha)
1	<b>M<sub>1</sub></b>	81.62 (65.19)	3.1 (10.85)	64.74	<b>91.89</b>	<b>92.125</b>	<b>10.555</b>	<b>12.295</b>	<b>10.685</b>
2	<b>M<sub>2</sub></b>	77.42 (59.65)	4.65 (12.43)	47.09	88.54	87.155	5.635	12.515	10.685
3	<b>M<sub>3</sub></b>	77.73 (61.94)	4.8 (12.6)	45.34	87.65	87.35	6.02	12.565	10.595
4	<b>M<sub>4</sub></b>	71.55 (57.78)	4.5 (12.03)	48.70	86.045	86.495	4.09	12.55	10.605
5	<b>M<sub>5</sub></b>	66.41 (54.65)	8.8 (17.24)	-	83.735	82.39	-	12.415	9.885
CD (P=0.05)		1.66	1.25	-	<b>3.83</b>	<b>1.96</b>		<b>NS</b>	

*In pooled mean analysis, highest millable cane (91089/ha) and yield (92.12 tonnes / ha) and sugar yield ( 10.67 tonnes/ha) were recorded in **M<sub>1</sub>**- (Single bud treated with *Trichoderma asperellum* (TNAU)@ 4 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>) followed by **M<sub>2</sub>**- (Single bud treated with *Bacillus subtilis* (TNAU)@ 10 g l<sup>-1</sup> + soil application @ 2.5 kg ha<sup>-1</sup>) in which, the cane yield of 88.54 tonnes/ha and sugar yield of 10.68 tonnes/ha were recorded. The CCS percent among the treatment is also significantly different.*



### 3.2 Pooled Mean Analysis

Among the biocontrol agents *P. fluorescens*, *Bacillus subtilis* and *Chaetomium globosum* are highly potential antagonists [9]. Antagonistic strains of the genus *Bacillus* are advantageous over other biocontrol agents in numerous ways, as they are ubiquitous in soils, sporulate excessively, have prolonged shelf life and enhance plant nutrition[10]. Their efficacy in controlling many plant diseases has repeatedly been proven. Three strains of the genus *Bacillus* reduced disease incidence by 45-49% in sugarcane plants challenged by pathogen inoculation in the stem and by 48-56% in the plants inoculated in the soil near the roots. *Bacillus* strains significantly controlled red rot disease on sugarcane varieties Co-1148 and SPF-234 under field conditions[11]. The antagonistic strains suppressed the primary infection (Soil inoculation) as well as secondary infection (Challenge inoculation) of red rot which is caused by certain vectors like the sugarcane borer (*Diatraea saccharalis*) and tools used in various cultural operations. These *Bacillus* strains also produced the broad-spectrum antifungal metabolite surfactin [12] which further characterize them as systemic resistance inducers, since surfactin is a good inducer of systemic resistance in [13,14].

### 4. CONCLUSION

This study provides a technology capsule for the management of Set rot of sugarcane. Weather parameter studies provide an theoretical and practical knowledge about the correlation between sett rot, maximum temperature, minimum temperature, RH, rainfall and rainy day which is helpful for the development of disease forecasting model.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Fischer D, Pfitzner B, Schmid M, Simões-Araújo JL, Reis VM, Pereira W, Ormeño-Orrillo E, Hai B, Hofmann A, Schloter M. Molecular characterisation of the diazotrophic bacterial community in uninoculated and inoculated field-grown sugarcane (*Saccharum* sp.) Plant Soil. 2012;356:83–99.
2. Vanitha SC, Sharan Umesha. *Pseudomonas fluorescens* mediated systemic resistance in tomato is driven through an elevated synthesis of defense enzymes. *Biologia plantarum*. 2011;55(2): 317-322
3. Thirumurugan A, Koodalingam K. Management of borer complex in sugarcane through companion cropping under drought condition of Palar river basin area. *Sugar Tech*. 2005;7:163-164
4. Thirumurugan A, Joseph M, Sudhagar R, Meenakshi Ganesan N. Improving efficacy of *Trichogramma chilonis* against shoot borer, *Chilo infuscatellus* (Snellen) in sugarcane ecosystem of tropical India. *Sugar Tech*, Volume 2006;8:155-159.
5. Comstock JC, Glaz B, Edme S, Davidson R, Gilbert R, Glynn N, Miller J, Tai P. Registration of 'CP 00-1446' Sugarcane. *J. Plant Regist*. 2009;3:28–34.
6. Shen B, Zhang W, Zhang J, Zhou J, Wang J, Chen L, Wang L, Hodgkins A, Iyer V, Huang X, Skarnes WC. Efficient genome modification by CRISPR-Cas9 nickase with minimal off-target effects. *Nat Methods*. 2014 Apr;11(4):399-402.
7. Matthijs S, Tehrani KA, Laus G, Jackson RW, Cooper RM, Cornelis P.. Thioquinolobactin, a *Pseudomonas* siderophore with antifungal and anti-*Pythium* activity. *Environ. Microbiol*. 2007; 9:425–434.
8. El Tarabily A, Sivasithamparam K. Potential of yeasts as biocontrol agents of soil-borne fungal plant pathogens and as plant growth promoters. *Mycoscience*. 2006;47(1):25-35.
9. Henry M, Picard N, Carlo T, Manlay J, Valentini R, Bernoux M, Laurent André. Estimating tree biomass of sub-Saharan African forests: a review of available allometric equations. *Silvia Fennica*. 2011; 45(38).
10. Hassan SM, Haq AU, Byrd JA, Berhow MA, Cartwright AL, Bailey CA. Haemolytic and antimicrobial activities of saponin-rich extracts from guar meal. *Food Chem*. 2010;119:600–605
11. Viswanathan R. Recent advances in sugarcane disease management. In: *Role of bio-control agents for disease management in sustainable agriculture*. (Ed. P. Ponmurugan and M.A. Deepa),

- Research India Publications. New Delhi. 2009;399-430.
12. Jayakumar V, Ramesh Sundar A, Viswanathan R. Biological suppression of sugarcane smut with endophytic bacteria. Sugar Tech. 2018;21(3):123
  13. Muhammad Nadeem Hassan, Shahid Afghan, Fauzia Y. Hafeez. Biological control of red rot in sugarcane by native pyoluteorin-producing *Pseudomonas putida* strain NH-50 under field conditions and its potential modes of action. Pest Management Science. 2011;67(9).
  14. Shanmugam PS, Sangeetha M, Saravanan NA, Tamil selvan N. Management of sett rot (*Ceratocystis paradoxa* (de Seynes) moreau) in sustainable sugarcane initiative (ssi) Nurseries. The Bioscan. 2016;11(3):1381-1384.

© 2023 Yamunarani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/112081>