



# Efficacy of Acupuncture in Chronic Low Back Pain Treatment: A Literature Review

Lucca Tamara Alves Carretta <sup>a\*</sup>,  
Pedro Rodrigues Teixeira <sup>a</sup>, Ana Clara Pirovani Dias <sup>a</sup>,  
Carolina Mesquita Devens <sup>a</sup>, Luiza Norbim Rones <sup>a</sup>,  
Camila Kafuri Bonacossa <sup>b</sup>  
and Cássio Newton Bomfim Barbosa <sup>c</sup>

<sup>a</sup> Faculty of Medicine, Escola Superior de Ciências da Santa Casa de Misericórdia de Vitória, Vitória, Brasil.

<sup>b</sup> Faculty of Medicine, Universidade de Vila Velha, Vila Velha, Brasil.

<sup>c</sup> Department of Family and Community Medicine, Faculty of Medicine, Escola Superior de Ciências da Santa Casa de Misericórdia de Vitória, Vitória, Brasil.

## Authors' contributions

This work was carried out in collaboration among all authors. Authors LTAC, PRT, ACPD, CMD, LNR and CKB elaboration and design of the study method, bibliographical research, interpretation and technical analysis of results and discussion with data from the literature. Author CNBB guidance and critical review of the manuscript. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/JOCAMR/2024/v25i2515

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112463>

**Minireview Article**

**Received: 07/12/2023**  
**Accepted: 10/02/2024**  
**Published: 15/02/2024**

\*Corresponding author: E-mail: [lucattamara12@gmail.com](mailto:lucattamara12@gmail.com);

## ABSTRACT

**Aims:** To evaluate the effects of the use of acupuncture as a treatment of chronic low back pain.

**Methodology:** This study follows the integrative methodology of literature review, seeking articles published on PubMed, using keywords from Medical Subject Headings (MeSH) to form the search

**Formula:** Acupuncture AND "low back pain". Inclusion criteria were randomized controlled trials published between 2018 and 2023, with free full text available in english or portuguese. The exclusion criteria was not answering this review's orienting question.

**Results:** Several studies analyzed the efficacy of different acupuncture and electroacupuncture approaches in the treatment of chronic low back pain. One study highlighted invasive laser acupuncture with different wavelengths, revealing improvements in pain and disability scales after 4 weeks. Another research compared electroacupuncture at various frequencies, observing similar reductions in pain between the groups. Combined acupuncture (hands and ear) was more effective than usual medical treatment, showing greater reduction in pain and cure rate. Both electroacupuncture and manual acupuncture also demonstrated efficacy in reducing pain intensity and quality. Acupuncture with embedded wires stood out, showing positive and rapid results in chronic low back pain. Another study revealed that both electroacupuncture and a placebo had positive impacts on pain intensity.

**Conclusion:** Even though there are multiple forms of acupuncture used throughout the studies, and even those that appear on multiple studies are done differently on those appearances, all are used as a way to reduce CLBP. In that sense, the positive effects from using traditional acupuncture or some different form of acupuncture as a treatment for CLBP is highlighted by the results from the studies that were a part of this review.

*Keywords: Acupuncture; low back pain; complementary therapies; therapeutics; chronic pain.*

## 1. INTRODUCTION

Chronic low back pain (CLBP) is characterized by dysfunction in lumbar biomechanics, manifesting as painful episodes lasting more than 12 weeks [1]. It can be classified as specific when the etiology is known, such as hernias or fractures, or nonspecific when the cause is unknown [2]. Furthermore, CLBP is considered a public health issue, being one of the primary causes of work absenteeism worldwide [3]. CLBP has a global prevalence of 25.4% in the elderly population ( $\geq 60$  years) and 3-20% in adults ( $< 60$  years) [4,5].

Various therapeutic interventions have been proposed as alternatives for CLBP treatment, given that this condition affects approximately 80% of the population at some stage in life, primarily impacting the elderly population, which is more susceptible to the side effects of analgesics and anti-inflammatories. Additionally, certain factors are associated with the presence of CLBP, such as male gender, age over 30 years, smoking, alcohol consumption, obesity, and improper posture [6].

Some of the non-traditional curative and preventive measures for CLBP include pilates, kinesiotherapy, heat therapy, hydrotherapeutic

treatment, and acupuncture [7]. Currently, acupuncture is part of Integrative and Complementary Practices (PICS) offered by the Unified Health System (SUS) and is used as an alternative treatment for CLBP, potentially effective in reducing symptoms and consequently improving the patient's quality of life by alleviating pain [8,9].

Acupuncture is an ancient therapeutic practice that involves the insertion of needles into specific points on the body for pain relief [10]. There are variations of the technique that incorporate technological advancements, such as electrical stimulation, the use of lasers in conjunction with needles, and the insertion of biodegradable threads to enhance the effectiveness of acupuncture [11]. As an intervention that avoids the use of pharmaceuticals, acupuncture is an important alternative for patients affected by CLBP [12,13].

Therefore, this mini review aims to evaluate the effects from the use of acupuncture as a treatment of chronic low back pain.

## 2. METHODOLOGY

This study consists of a review following the integrative methodology for bibliographical

research [14]. In order to clarify the purpose of this review, the following question was determined: "Does the practice of different forms of acupuncture have any effects in treating chronic low back pain?". The search for articles was conducted on the PubMed database during July and August of 2023, utilizing keywords from the Medical Subject Headings (MeSH) in order to form the following search formula: Acupuncture AND "low back pain.". The selection criteria were randomized controlled trials published in English or Portuguese between 2018 and 2023. Two different researchers reviewed the reference list, excluding those not related to the theme of this review by first reading the title and abstract of the study, then analyzing the remaining studies' full text. Studies whose full text could not be found were excluded.

### 3. RESULTS

A total of 30 articles were found on PubMed using the search strategy described above. Upon reviewing the titles and abstracts, 21 articles were excluded, leaving 9 for full-article reading. Of these, 3 articles were excluded as they did not meet the objective of this review. Therefore, a total of 6 articles were included in this review, with the main findings described below.

Kim et al. examined the efficacy of Invasive Laser Acupuncture (ILA) with lasers of different wavelengths for treating CLBP. Participants were allocated into 3 groups: control, which received acupuncture without laser activation; a group with laser activation at a wavelength of 650 nanometers (nm); and a group with laser

activation at a wavelength of 830 nm. Following a bilateral acupuncture treatment lasting 10 minutes with ILA and subsequent electroacupuncture, also for 10 minutes, over 4 weeks, different patient responses regarding pain scales were observed. In terms of the Visual Analog Scale for pain (VAS), there was an average drop from 50.47 mm to 36.20 mm in the control group; from 55.07 mm to 23.13 mm in the 650 nm group; and from 49.60 mm to 28.60 mm in the 830 nm group after the 4 weeks of treatment. Regarding the Oswestry Disability Index (ODI), there was a decrease from 9 to 8 points in the control group, from 11 to 7 points in the 650 nm group, and from 11 to 5 points in the 830 nm group [15].

Over a 5-week period, Torres et al. conducted electroacupuncture (EA) therapy at different frequencies: high (100 Hz), low (2 Hz), alternate (2 and 100 Hz), control (without electrical stimulation), and placebo (with an adhesive moxa, mimicking needle penetration). EA sessions were held for 30 minutes, twice a week for 5 weeks. A significant difference between all groups from pre-treatment to post-treatment was observed, with an average decrease from 6.8 to 1.9 on the Numeric Pain Rating Scale (NPRS). No significant differences were found between the three groups with electrical stimulation at different frequencies. The study also showed that manual (control) and electrical (intervention) stimulation did not differ statistically and that a placebo effect occurred, with the placebo group achieving the greatest reduction in NPRS (from 7.5 to 0.9) [16].

**Table 1. Articles included as results from this review's search.**

Title	Authors	Year	Country
Efficacy of invasive laser acupuncture in treating chronic non-specific low back pain: A randomized controlled trial	Kim et al.	2022	South Korea
Effect of Different Frequencies of Electroacupuncture on Chronic Low Back Pain in Older Adults: A Triple-blind, Placebo-controlled, Randomized Clinical Trial	Torres et al.	2023	Brazil
Effect of hand-ear acupuncture on chronic low-back pain: a randomized controlled trial	Luo et al.	2019	China
Effectiveness of Manual and Electrical Acupuncture for Chronic Non-specific Low Back Pain: A Randomized Controlled Trial	Comachio et al.	2020	Brazil
Efficacy of Electronic Acupuncture Shoes for Chronic Low Back Pain: Double-Blinded Randomized Controlled Trial	Yeh et al.	2020	Taiwan
Efficacy and safety of thread embedding acupuncture combined with acupuncture for chronic low back pain: A randomized, controlled, assessor-blinded, multicenter clinical trial	Sung et al.	2020	South Korea

Luo and colleagues investigated, after 7 weeks of treatment, the superiority of acupuncture, whether hand acupuncture combined with auriculotherapy or standard acupuncture, compared to usual medical treatment for CLBP. Three groups were used for comparison of therapy results with VAS. After 6 months of treatment, a decrease from 6.7 to 3.02 on the VAS (0 to 10) was observed in the combined hand acupuncture and auriculotherapy group, from 6.92 to 4.16 in the standard acupuncture group, and 1.42 points in the usual medical treatment group. The research found that after 6 months of treatment, there was a cure rate of 70.4% and a pain reduction of 18.5% in the combined acupuncture group. In contrast, a cure rate of 46% and a 22% pain reduction were observed in the standard acupuncture group, and a 22.9% cure rate and a 22.9% pain reduction in the usual medical treatment group [17].

Comachio et al. examined the effectiveness of EA and manual acupuncture (MA) in treating patients with CLBP over 6-week, with a frequency of 2 sessions per week and 40 minutes per session. EA received 30 minutes of manual acupuncture, followed by 10 minutes of electroacupuncture at a frequency of 10 Hz. Pain intensity was quantified using the Numeric Rating Scale (NRS), and pain quality was assessed using the McGill Pain Questionnaire (MPQ). The treatment results indicated a reduction from 7.9 to 3.8 and from 7.8 to 4.2 in the NRS for the MA and EA groups, respectively, between the pre-treatment and post-treatment periods. In the sensory component of the MPQ, there was a decrease from 17.3 to 8 for MA and from 20.1 to 9.4 for EA during the same period [18].

In the study led by Yeh and colleagues, 60 participants were divided into a group receiving nonsteroidal anti-inflammatory drugs (NSAIDs) and fake electronic acupuncture shoes and another group with electronic acupuncture shoes (EAS) and a placebo for NSAIDs to compare the effectiveness of both over 6 weeks. The Visual Analog Scale (VAS) was used to analyze pain intensity during this period. The study found, after 2 weeks of treatment, an effectiveness of 68% in the EAS group and 59% in the NSAIDs group, based on a decrease in VAS. In contrast, after 1 week of treatment, there was 78% effectiveness in EAS and 59% in NSAIDs, while after the last treatment session, these numbers were 73% and 62%, respectively. The average VAS score before starting treatment was 4.4, and at the last visit, it was 3.1 [19].

The Korean study conducted by Sung and others evaluated the effects of acupuncture and Thread Embedding Acupuncture (TEA) as a treatment for CLBP. Both treatments resulted in a clinically relevant reduction in VAS after 8 weeks of the study. However, the TEA group showed a higher average reduction of 33.7, compared to the traditional acupuncture group, which had an average reduction of 15.6. The reduction in pain was observed in the long term for both treatments, increasing over time and with more sessions. The reduction in the TEA-treated group occurred more rapidly, within the first month of the study, while the acupuncture group experienced it from one and a half months onwards [20].

#### 4. DISCUSSION

The use of ILA was shown by Kim et al. and Lin et al. to reduce pain levels in patients with CLBP. Kim et al. evaluated different frequencies of ILA, with 650 nm having the most improvements on the pain scale. Lin et al. combined ILA with Chinese Cupping and had better results on the ILA group when compared to sham ILA group [15,21]. Similar effects of ILA were demonstrated by Huang et al., as a treatment for temporomandibular joint disorder, and by Manente et al., as a way to neuralgia on facial neuropathies [22,23]. These findings suggest that ILA is an effective way to reduce pain in multiple conditions and has further potential to be yet explored, with studies to compare its efficacy with regular forms of acupuncture and the best therapeutic protocols for its frequency being needed.

Comachio et al. and Torres et al. both found the use of EA to be effective as a treatment for CLBP, even with different charges being used (2 Hz, 10 Hz, 100 Hz). While Torres et al. focused on elders and only EA was utilized as a form of treatment, Comachio et al. had a mean patient age of 46 years in the EA group and tested a combination of manual acupuncture and EA [16,18]. Both studies, although they did not show any superiority of EA when compared to normal, manual acupuncture, presented with improvement on the pain scale. EA may not be more effective than manual acupuncture, but it is shown to be as effective regarding pain relief.

Yeh et al. also studied the effect EA has on CLBP, implementing it on therapeutic shoes, which resulted in analgesic effects and improvements in range of motion [19]. As elucidated by Zheng et al. and Wan et al., EA

also has an analgesic role on knee osteoarthritis and dry eye pain, reducing the expression of pain receptors and inflammation on rat models [24,25] and was shown to help during the rehabilitation of meniscal injuries on athletes [26]. Thus, EA could be an effective alternative treatment for many forms of not only CLBP but several chronic pain illnesses. Further studies exploring EA's effects on pain signaling in humans and anti-inflammatory effects may lead to discoveries regarding its benefits to a variety of diseases.

Auriculotherapy (AT) is a variation from traditional acupuncture, specifically performed on the patient's outer ear. Luo et al. combined it with acupuncture performed on specific points of the hand to analyze the treatment of CLBP with hand-ear acupuncture, finding better long-term results on patients treated with hand-ear than those treated by regular acupuncture [17]. Ushinohama et al. only evaluated the use of AT in a single session of treatment, but even with the methodological differences both found AT to have a great effect on reducing CLBP [27]. AT was also effective as a complementary treatment for chronic spinal pain, temporomandibular joint disorder pain, premenstrual syndrome symptoms and orthodontic pain [28-31]. AT is another form of acupuncture with many helpful effects that should keep being studied and explored to keep evolving it as a complementary to pain management.

Sung et al. combined TEA with acupuncture as a treatment for CLBP, with positive results on pain relief, but did not find any statistically significant difference between the use of TEA and the use of acupuncture [20]. Lee et al. found similar results, showing that TEA can be effective against CLBP, but without any superiority when compared to normal acupuncture [32]. Even though TEA isn't superior to regular acupuncture and may even have more risks attached as it consists of a more invasive treatment than AC, different studies have shown TEA to have as broad of a range as other acupuncture variations. Woo et al., Li et al., and Goo et al. studied TEA concerning knee osteoarthritis, anorectal pain, and herniated lumbar intervertebral disc, respectively, all with improvements in pain levels. Goo et al. also showed that TEA can have a greater long-term effect, even after treatment was already been done [33-35].

All five forms of acupuncture, traditional, ILA, EA, AT, and TEA were all found to be effective in relieving pain, not only CLBP but also knee

osteoarthritis, temporomandibular joint disorder, etc. Some studies even show acupuncture and some variations to have psychological effects, being complementary to medications when treating depression and anxiety [30,36,37]. And, as elucidated by Kong et al., the psychological effect of acupuncture may also be an effective way to help relieve pain in patients with CLBP [38]. This comes to show that acupuncture, in its many forms, has a wide range of beneficial effects that should be further explored. It is not an ancient technique without reason, and the combination with more recent developments shown in EA and ILA for example, may be the key to unlocking its full potential. More research with a bigger number of participants should help establish the most efficient and safe protocols for acupuncture and its varieties to follow when treating CLBP.

## 5. CONCLUSION

Even though there are multiple forms of acupuncture used throughout the studies, and even those that appear in multiple studies are done differently in those appearances, all are used as a way to reduce CLBP. In that sense, the positive effects of using traditional acupuncture or some different form of acupuncture as a treatment for CLBP are highlighted by the results from the studies that were a part of this review.

## CONCENT AND ETHICAL APPROVAL

It is not applicable.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Andersson GB. Epidemiological features of chronic low-back pain. *Lancet*. 1999;354(9178):581-5.
2. Bottamedi X, Ramos JS, Arins MR, Murara N, Woellner SS, Soares AV. Treatment program for chronic low back pain based on the principles of Segmental Stabilization and Back School. *Rev Bras Med Trab*. 2016;14(3):206-213.
3. Hoy D, Bain C, Williams G, March L, Brooks P, Blyth F et al. A systematic review of the global prevalence of low back pain. *Arthritis Rheum*. 2012;64(6):2028-37.

4. Oliveira RAA, Stump PRNAG, Rezende MC, Vieira LF, Silva MA, Elorza PM. Jornada do paciente com lombalgia crônica no Brasil: Uma revisão semi-sistemática sobre a melhor abordagem. *Rev Med.* 2022;101(5).
5. Hoy D, Brooks P, Blyth F, Buchbinder R. The Epidemiology of low back pain. *Best Pract Res Clin Rheumatol.* 2010;24(6):769-81.
6. Rigo JC, Pires APM, Oliveira LA, Harres SS, Dalacorte RR. Evidence of acupuncture in treatment of low back pain. *Geriatr Gerontol.* 2011;5(3):175-179.
7. Almeida DC, Kraychete DC. Low back pain - a diagnostic approach. *Rev Dor.* 2017;18(2):173-177.
8. Chou R, Qaseem A, Snow V, Casey D, Cross JT Jr, Shekelle P et al. Diagnosis and treatment of low back pain: A joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med.* 2007;147(7):478-91.
9. Política Nacional de Práticas Integrativas e Complementares no SUS (PNPIC). Ministério da Saúde; 2006. Accessed 15 January 2024. Available:<https://bvsmis.saude.gov.br/bvs/publicacoes/pnpic.pdf>
10. Martins ES, Costa N, Holanda SM, Castro RCMB, Aquino PS, Pinheiro AKB. Enfermagem e a prática avançada da acupuntura para alívio da lombalgia gestacional. *Acta Paul Enferm.* 2019;32(5):477-484.
11. Mao JJ, Liou KT, Baser RE, Bao T, Panageas KS, Romero SAD et al. Effectiveness of Electroacupuncture or Auricular Acupuncture vs Usual Care for Chronic Musculoskeletal Pain Among Cancer Survivors: The PEACE Randomized Clinical Trial. *JAMA Oncol.* 2021;7(5):720-727.
12. Patil S, Sen S, Bral M, Reddy S, Bradley KK, Cornett EM, Fox CJ, Kaye AD. The Role of Acupuncture in Pain Management. *Curr Pain Headache Rep.* 2016;20(4):22.
13. Mu J, Furlan AD, Lam WY, Hsu MY, Ning Z, Lao L. Acupuncture for chronic nonspecific low back pain. *Cochrane Database Syst Rev.* 2020;12(12).
14. Souza MT, Silva MD, Carvalho R. Integrative review: what is it? How to do it? *Einstein (Sao Paulo).* 2010;8(1):102-6.
15. Kim JH, Na CS, Cho MR, Park GC, Lee JS. Efficacy of invasive laser acupuncture in treating chronic non-specific low back pain: A randomized controlled trial. *Plos One.* 2022;17(5):e0269282
16. Torres SF, de Macedo ACB, Sakai RY, Bressan GCS, Dos Santos MBR, Marques AP. Effect of Different Frequencies of Electroacupuncture on Chronic Low Back Pain in Older Adults: A Triple-blind, Placebo-controlled, Randomized Clinical Trial. *Pain Physician.* 2023 Mar;26(2):161-173.
17. Luo Y, Yang M, Liu T, Zhong X, Tang W, Guo M, Hu Y. Effect of hand-ear acupuncture on chronic low-back pain: A randomized controlled trial. *J Tradit Chin Med.* 2019;39(4):587-598.
18. Comachio J, Oliveira CC, Silva IFR, Magalhães MO, Marques AP. Effectiveness of Manual and Electrical Acupuncture for Chronic Non-specific Low Back Pain: A Randomized Controlled Trial. *J Acupunct Meridian Stud.* 2020;13(3):87-93.
19. Yeh BY, Liu GH, Lee TY, Wong AM, Chang HH, Chen YS. Efficacy of Electronic Acupuncture Shoes for Chronic Low Back Pain: Double-Blinded Randomized Controlled Trial. *J Med Internet Res.* 2020;22(10).
20. Sung WS, Hong Y, Jeon SR, Yoon J, Chung EK, Jo HG et al. Efficacy and safety of thread embedding acupuncture combined with acupuncture for chronic low back pain: A randomized, controlled, assessor-blinded, multicenter clinical trial. *Medicine (Baltimore).* 2020;99(49).
21. Lin ML, Wu JH, Lin CW, Su CT, Wu HC, Shih YS, Chiu IT, Chen CY, Chang WD. Clinical Effects of Laser Acupuncture plus Chinese Cupping on the Pain and Plasma Cortisol Levels in Patients with Chronic Nonspecific Lower Back Pain: A Randomized Controlled Trial. *Evid Based Complement Alternat Med.* 2017;2017:3140403.
22. Huang YF, Lin JC, Yang HW, Lee YH, Yu CH. Clinical effectiveness of laser acupuncture in the treatment of temporomandibular joint disorder. *J Formos Med Assoc.* 2014;113(8):535-9.
23. Manente R, Pedroso GL, Gomes E Moura AP, Borsatto MC, Corona SAM. Laser acupuncture in the treatment of neuropathies in dentistry: A systematic review. *Lasers Med Sci.* 2023. 25;38(1):92.

24. Zheng X, Lin J, Wang Z, Zeng Z, Chen H. Research of the analgesic effects and central nervous system impact of electroacupuncture therapy in rats with knee osteoarthritis. *Heliyon*. 2024;10(1):e21825.
25. Wan MM, Jin T, Fu ZY, Lai SH, Gao WP. Electroacupuncture Alleviates Dry Eye Ocular Pain Through TNF- $\alpha$  Mediated ERK1/2/P2X3R Signaling Pathway in SD Rats. *J Pain Res*. 2023;16:4241-4252.
26. Li Y. Sports rehabilitation in athletes with meniscal lesions based on electroacupuncture associated with sports therapy. *Rev Bras Med Esporte*. 2023;29:e20220150.
27. Ushinohama A, Cunha BP, Costa LOP, Barela AMF, Freitas PB de. Effect of a single session of ear acupuncture on pain intensity and postural control in individuals with chronic low back pain: a randomized controlled trial. *Braz J Phys Ther*. 2016;20(4):328–35.
28. Morais BX, Munhoz OL, Moreira CHC, Kurebayashi LFS, Lopes LFD, Magnago TSBS. Auriculotherapy for reducing chronic spinal pain in health workers: a clinical trial. *Rev Lat Am Enfermagem*. 2023;31:e3953.
29. Ferreira LA, Grossmann E, Januzzi E, Gonçalves RT, Mares FA, de Paula MV, Carvalho AC. Ear Acupuncture Therapy for Masticatory Myofascial and Temporomandibular Pain: A Controlled Clinical Trial. *Evid Based Complement Alternat Med*. 2015;2015:342507.
30. Korelo RIG, Moreira NB, Miguel BAC, Cruz CGD, Souza NSP, Macedo RMB, Gallo RBS. Effects of Auriculotherapy on treatment of women with premenstrual syndrome symptoms: A randomized, placebo-controlled clinical trial. *Complement Ther Med*. 2022;66:102816.
31. Serritella E, Impellizzeri A, Liguori A, Galluccio G. Auriculotherapy used to manage orthodontic pain: a randomized controlled pilot study. *Dental Press J Orthod*. 2021;26(6):e2119381.
32. Lee HJ, Choi BI, Jun S, Park MS, Oh SJ, Lee JH, Gong HM, Kim JS, Lee YJ, Jung SY, Han CH. Efficacy and safety of thread embedding acupuncture for chronic low back pain: a randomized controlled pilot trial. *Trials*. 2018;19(1):680.
33. Woo SH, Lee HJ, Park YK, Han J, Kim JS, Lee JH et al. Efficacy and safety of thread embedding acupuncture for knee osteoarthritis: A randomized controlled pilot trial. *Medicine (Baltimore)*. 2022;101(31).
34. Li J, Sun Y, Zhang C, Thomas K, Lin W, Cheng C, Li H, Zhu Q, Ma S, Hua Q, Shi Q, Zheng X. A randomized, controlled clinical trial of acupoint catgut embedding as an effective control of functional anorectal pain. *Medicine (Baltimore)*. 2023;102(44).
35. Goo B, Kim JH, Kim EJ, Lee HJ, Kim JS, Nam D, Park YC, Kim TH, Baek YH, Nam SS, Seo BK. Thread embedding acupuncture for herniated intervertebral disc of the lumbar spine: A multicenter, randomized, patient-assessor-blinded, controlled, parallel, clinical trial. *Complement Ther Clin Pract*. 2022;46.
36. de Oliveira Rodrigues DM, Menezes PR, Machado Ribeiro Silotto AE, Heps A, Pereira Sanches NM, Schweitzer MC, Faisal-Cury A. Efficacy and Safety of Auricular Acupuncture for Depression: A Randomized Clinical Trial. *JAMA Netw Open*. 2023;6(11):e2345138.
37. Amorim D, Brito I, Caseiro A, Figueiredo JP, Pinto A, Macedo I, Machado J. Electroacupuncture and acupuncture in the treatment of anxiety - A double blinded randomized parallel clinical trial. *Complement Ther Clin Pract*. 2022 Feb;46:101541.
38. Kong JT, Puetz C, Tian L, Haynes I, Lee E, Stafford RS et al. Effect of Electroacupuncture vs Sham Treatment on Change in Pain Severity Among Adults With Chronic Low Back Pain: A Randomized Clinical Trial. *JAMA Netw Open*. 2020;3(10).

© 2024 Carretta et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/112463>