



Feasibility Tests of a Fish Farm Integrating the Breeding of Poultry and a Fly

Lawal W.S^{a,b*}, Alu S.O^c and Yusuf R.T^c

^a *Department of Agricultural Technology, Microbiology Unit, Kwara State Polytechnic, Nigeria.*

^b *Department of Science Laboratory Technology, Microbiology Unit, Kwara State Polytechnic, Nigeria.*

^c *Department of Biology, Microbiology Unit, Kwara State Polytechnic, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JALSI/2024/v27i2640

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112530>

Original Research Article

Received: 08/12/2023

Accepted: 12/02/2024

Published: 08/03/2024

ABSTRACT

A locally improvised poultry house built with pure local material called bamboo and covered with dried treated grasses (roof) known as sana which housed 300 laying birds was built near 20m X 20m earthen fish pond with an underground spring water as source of intake with a flow through system and stocked with an improved catfish *clarias geripinus* juveniles 2000 in number. The floor of the poultry house was cemented to prevent leakages of water and permanently covered with thick nylon white rubber and layer were placed at high level above the floor on top of bamboo tree having gaps in between them. The usage of these gaps created was to allow layers (chicken) dropping directly on the rubber which eventually served as breeding places for maggot. The maggot was collected washed thoroughly and processed to serve as feed to the fishes. The processed maggot reduces about 40% of total cost of feed in raising the fishes. The birds produced 70 pieces of eggs per day 7½ crates and a crate is sold for N2,700 per crate. The fishes were harvested after six (6) months of stocking with a total piece of 1,680 with an average of 800g (1344kg) and is sold at N1500 per kg. The revenue realized took care of both capital and recurrent

*Corresponding author: E-mail: awsl2004@gmail.com;

cost with a working force of three (3) people. The experiment contribute significantly ($P < 0.05$) in reducing the level of poverty and also contribute to food security, transformation agenda of Federal government of Nigeria.

Keywords: Maggot; bamboo; Clarias geripinus; juvenile and treated grasses.

1. INTRODUCTION

Poultry is the foremost developing segment of the Agricultural sector in Nigeria, Africa and indeed globally., The increase in poultry production has enhanced the per capita availability of eggs and per kilogram of poultry meat per annum [1].

In Nigeria, the average daily consumption of 54g protein with 6.5g from animal source falls below the recommended daily protein of 86g and 8.4g of protein from animal source [2] the reason being the exorbitant cost of fish, chicken and its products owing to high cost of feed which accounts for 60% - 80% total cost of production [2].

The integrated poultry-fish based farming provides enormous opportunities for improvisation of some of the necessary inputs especially feeds so that cost is saved and profit is improved as supplementary income for men and women empowerment and livelihood security to rural families.

The farming system also offers opportunities to these resource-poor families at meagre financial investments and addresses the issues of women empowerment and livelihood security at the grass root level. To encourage poultry entrepreneurship among women in rural households so as to support backyard or intensive broiler production and hatchery/small poultry feed mills in the rural areas and villages.

The basic reasons behind poor growth of poultry production in most of our villages and rural areas is the accessibility to basic poultry needs like good feedmills, feed ingredients and veterinary clinics and most importantly movement of youths from rural area to cities [3,4,5].

This type of system will bring a kind of recycling in Agriculture between fishery and poultry will create jobs for the rural people with minimum capital requirement to take off there by reducing poverty and also contribute to food security, transformation agenda of Federal government of Nigeria.

1.1 Justification

i-Poultry feaces is produced in large quantity in every poultry farm and this can cause nuisance
ii-Majority of the rural people are jobless.
iii-one of the major inputes in poultry production is land, which is readily available in the rural area.

1.2 Objective of the Study

i-Build both the poultry and fish pond on each other.
ii-Fed the fishes with poultry dropping.

2. MATERIALS AND METHODS

Site of experiment: The experiment was carried out in the Agricultural garden if the department of Agricultural Technology.

Construction: A locally improvised poultry house built with pure local material called bamboo and covered with dried treated grasses (roof) known as sana which housed 100 laying birds was built near 20m X 20m earthen fish pond with an underground spring water as source of intake with a flow through system and stocked with an improved catfish clarias geripinus juveniles 2000 in number. The floor of the poultry house was cemented to prevent leakages of water and permanently coveted with thick nylon white rubber and layer were placed at high level above the floor on top of bamboo tree having gaps in between them. The usage of these gaps created was to allow layers (chicken) dropping directly on the rubber which eventually served as breeding places for maggot [6,7].

Sources of the juvenile: The juvenile was purchased from Ministry of Agriculture, fisheries section along yidi road in Ilorin.

Sources of the layers: The point of layer was purchased from, Affcom farm Kulende junction , sango in Ilorin, the point of lay was allowed to start laying properly.

Sources of feed ingredients: The few feed ingredients that was used was purchased from Gbemisola feedmill house, Offa garage, Ilorin

Table 1. Performance characteristics of poultry and Fish combined production

Months	1	2	3	4	5	6
Feed intake	7.5	7.9	7.5	7.0	7.0	7.1
Egg prod	2130	2100	2070	2130	2190	2130
No of crates	2.3	2.3	2.3	2.3	2.4	2.3
Equiv. amt of money (N)	106,500	105,500	103,500	106,500	1680	106,500
SEM	0.1	0.4	0.1	0.1	0.2	0.1

Egg collection: egg were collected everyday into crates directly around 4pm, no sorting was done and the eggs was sold the way it is picked.

Statistics analysis: Data like feed intake, number of eggs collected, equivalent amount of money for the eggs collected, all these was subjected to complete randomized design.

3. RESULTS AND DISCUSSION

Every three hundred laying (200) birds will be fed by a bag of feed which is 25kg but the 300 birds that was used in the experiment, 1½ a bag will be fed to them on a daily basis, the table 1 above shows the amount of feed consumed when the maggot is supplemented and this accounts for about 40% reduction of the feed they were supposed to consumed a day and this has drastically reduced the amount of money for keeping the laying birds while the production rate is well above 65% or 70% in some situation, The fishes were harvested after six (6) months of stocking with a total piece of 1,680 with an average of 500g (1344kg) and is sold at N1500 per kg. The revenue realized took care of both capital and recurrent cost with a working force of three (3) people [8,9].

4. CONCLUSION AND RECOMMENDATION

It is concluded that the cost of feed has been reduced by 40% and egg production rate is about 70%. The sales of fishes is another source of income for sustainability. It is recommended that this system of farming could be adopted even in large scale production. The maggot could also be used in poultry feed production because it is a good source of methionine.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ajad MI, Lawal WS, Banjoko IK, Ahmed SA. Comparative proximate and mineral analysis of smoked dried and solar-dried African Mud Catfish; 2019.
2. Abu AO, Onifade AA, Abanikanda AA, Obineye RI. Status and promotional strategies for rabbit production in Nigeria.9th World Rabbits Congress Varona, Italy; 2008.
3. Lawal WS, Oladunjoye IO, Banjoko IK, Ahmed SA. Performance characteristics, sensory evaluation of broiler birds fed *Jatropha caucas*. A tertiary education trust fund sponsored project of Kwara state Polytechnic, Ilorin Nigeria; 2013.
4. Adewoye SO, Omotosho JS. Nutrient composition of some freshwater fishes in Nigeria Biosci. Res. Commun. 1997;11(4):333-336.
5. Akinneye JO, Amoo IA, Arannilewa ST. Effect of drying method on the nutritional composition of three species of (*Bonga spp*, *Sardinella spp*. And *Heterotis niloticus*. J. fish. Int; 2; 99-103 biology journal of North America. 2007;5(3):109-117.
6. Bolawa et al. Proximate composition properties of different fish species obtained from Lagos, Nigeria. Internet Journal of Food Safety. 2011;13:342-344.
7. Burgess GHO. Increasing the direct consumption of fish. In: WW Pirie (Edu). Food Protein Sources. International Biological Programme 4. Cambridge University Press, Cambridge, pp 187-200. Calabar. 1975;183.
8. Cui Y, Wootton RJ. Effects of ration, temperature and body size on the body composition, energy content and condition of the minnow, *Phoxinus*

- phoxinus* (L). Journal of Fish Biology. 2011;32:749-764.
9. Dawson AS, Griman AS. Quantitative seasonal changes in the protein, lipid and energy content of carcass, ovaries and liver of adult plaice (*Pleuronectes platena*) Fisheries. 1980;48: 297-30.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/112530>