



Evaluation of Biocontrol-based Formulations against Late Leaf Spot and Rust in Groundnut

Karuppiah Manikandan ^a, Raman Gopalakrishnan ^b
and Kaliyamoorthy Manikandan ^{c*}

^a Krishna College of Agriculture and Technology, Madurai, Tamil Nadu – 625532, India.

^b ICAR- Indian Agricultural Research Institute, New Delhi-110012, India.

^c Faculty of Agriculture, Annamalai University, Tamil Nadu-608002, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author Karuppiah Manikandan carried out the experiments and data analysis, author RG edited the manuscript and author Kaliyamoorthy Manikandan conceptualized and wrote the paper. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JABB/2024/v27i5838

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116170>

Original Research Article

Received: 18/02/2024

Accepted: 22/04/2024

Published: 25/04/2024

ABSTRACT

This study investigated the efficacy of various biocontrol-based formulations against late leaf spot (LLS) (*Phaeoisariopsis personata*) and rust (*Puccinia arachidis*) in groundnut. Pot culture and field experiments were conducted during the Kharif season (June-September) of 2022. Disease incidence of both late leaf spot and rust was significantly lower in groundnut plants treated with a combination of biocontrol-based formulations compared to control. The most effective eco-friendly treatment included a combination of *Trichoderma asperellum*, *Pseudomonas fluorescens*, fortified lignite fly ash and Annamalai mixture. This combination significantly reduced lesion frequency and Percent Disease Index (PDI) for both diseases at 50, 70 and 90 DAS, compared to other treatments. The combination of biocontrol-based formulations also significantly increased the pod yield. Furthermore, the same treatment increased the induction of defense enzymes peroxidase

*Corresponding author: E-mail: manipatho23194@gmail.com;

(2.22 fold), polyphenol oxidase (1.28 fold) and phenylalanine ammonia-lyase (6.25 fold). While Carbendazim effectively reduced disease incidence, it was surpassed by the biocontrol-based module in terms of both disease control and yield. These findings suggest that biocontrol-based formulations can be a sustainable approach for managing LLS and rust diseases in groundnut cultivation.

Keywords: Leaf spot; rust; defense enzyme; biocontrol-based module; groundnut.

1. INTRODUCTION

Groundnut (*Arachis hypogaea* L.), is often hailed as the "king of oilseeds." Globally, India has the largest cultivation area, ranking second in global production. In India, groundnut production in the *kharif* season is 68.57 lakh tonnes with an average productivity of 1,562 kg/ha (IOPEPC, 2023). Groundnut fosters soil health through symbiotic nitrogen fixation with Rhizobia bacteria, promoting sustainable agricultural practices. Groundnut cultivation faces a formidable challenge from fungal diseases like late leaf spot (LLS), rust, Alternaria blight, stem rot, dry root rot and collar rot [1,2]. Among these, LLS disease caused by *Phaeoisariopsis personata* and rust disease caused by *Puccinia arachidis* pose the most significant threats. These widespread fungal pathogens inflict severe damage on groundnut crops. In most cases, these two pathogens occur together and hamper groundnut production [3]. In India, these diseases have been documented to cause yield losses exceeding 70%, significantly impacting groundnut productivity [4]. Leaf rust alone could cause yield reduction reaching up to 65%, especially in areas with high rainfall [5]. The detrimental effects extend beyond yield reduction, as the quality of groundnut seeds is also compromised by these fungal infections.

In groundnut cultivation, conventional disease management practices for controlling LLS and rust often rely on chemical fungicides. Their indiscriminate use has led to concerns about environmental and human health risks [1]. Environmental contamination, disruption of ecological balance due to harm to beneficial soil microbes, and potential health risks to humans and non-target organisms are some of the concerning consequences associated with chemical fungicide application [6,7]. The emergence of eco-friendly alternatives presents a promising solution for sustainable groundnut production [8]. Antagonistic microorganisms offer a particularly effective approach for managing these fungal diseases. These beneficial microbes can act against pathogens through multiple

mechanisms, including competition for resources, production of antibiosis compounds, predation and parasitism [9]. Additionally, some antagonists have the potential to induce systemic resistance within the plant itself [10]. These mechanisms offer a multifaceted approach to disease control, potentially mitigating the need for harmful chemical fungicides.

This work explores use of antagonist microorganisms, as a sustainable and effective disease management strategy against the detrimental fungal pathogens, *P. personata* and *P. arachidis* in groundnut.

2. MATERIALS AND METHODS

2.1 Survey of Late Leaf Spot and Rust Disease Incidence in Groundnut

Field surveys were employed to assess the prevalence and severity of groundnut LLS and rust diseases in Cuddalore district. Villages with a history of groundnut cultivation were targeted to capture representative data. Ten locations were chosen for fixed-plot surveys. Within each plot, the number of groundnut plants exhibiting symptoms of LLS and rust disease were recorded alongside the total number of plants observed.

2.2 Inoculum Culture of *P. personata* and *P. arachidis*

Naturally infected leaves with LLS and rust disease symptoms were collected from fields within the Cuddalore district. These diseased leaves served as the inoculum source for *P. personata* and *P. arachidis*. The inoculum was maintained through pot culture containing groundnut plants.

2.3 Assessment of Disease Severity

Groundnut seeds (cv. VRI-2) were grown in pots (20 cm dia, 5 plants/pot) for 30 days within a polyhouse. The potting medium consisted of a 2:1:1 ratio of field soil, farmyard manure (FYM)

and sand. The plants were then inoculated with a spore suspension mixture of *P. personata* and *P. arachidis*. The spore suspension was prepared by following the protocol of Subrahmanyam et al. [11]. To achieve optimal infection conditions, an alternating wet and dry period was implemented following inoculation. The inoculated plants were maintained at $24 \pm 2^\circ\text{C}$. Each treatment group within the glasshouse experiments comprised 15 plants with three replicates. Disease severity of both *P. personata* and *P. arachidis* were assessed using disease incidence and lesion frequency following the methods established by Subrahmanyam et al. [11].

Quantification of lesion incidence in response to various treatments for combined LLS and rust disease [3].

Lesion frequency (LF) = Number of lesion or pustules/cm² leaf area

Disease scoring was done with modified 9-point (1-9) scale [11].

Percent diseases index (PDI)% = Sum of individual rating / Total no. of leaves observed x 100 / Maximum grade

2.4 Preparation of Biocontrol Based Formulations

Aliquots of bovine urine, bovine dung, ovine dung, poultry litter, and neem cake were obtained at a conc. of 100% (w/w) and homogenously mixed in a 1:1:1:1:1 ratio to prepare Annamalai mixture. The resulting organic amendment was filtered and subsequently diluted to 20%, 40%, 60%, and 80% (v/v) using distilled water at the time of application [12].

The antagonists, *T. asperellum* and *P. fluorescens*, were cultivated in Potato Dextrose broth and King's B broth respectively. Class F lignite fly ash (LFA), obtained from Neyveli Lignite Corporation, Neyveli, India, was used as the carrier material. One kilogram of LFA was mixed thoroughly with 10 grams of carboxymethyl cellulose (CMC). The LFA-CMC mixture was then autoclaved for two consecutive days to ensure proper sterilization. To 1000 g of sterilized LFA-CMC carrier, 400 ml of bacterial suspension, containing approximately 8×10^9 CFU/mL, were separately added. 20 ml of molasses was then incorporated into the mixture, followed by thorough mixing. The fortified LFA was shade-dried for 2 h under sterile conditions

to remove excess moisture. The final product, designated as fortified lignite fly ash, was packaged in polyethylene bags and stored at $28^\circ\text{C} \pm 2^\circ\text{C}$. To monitor the viability, the population of antagonistic strains within the fortified LFA was periodically evaluated using the serial dilution technique [13].

2.5 Evaluation of Antagonists Based Formulations against LLS and Rust under Pot culture Condition

The experiment included 10 treatments with three replications, conducted from June to September 2022 at the Department of Plant Pathology, Annamalai University. Previously isolated and available biocontrol agents (BCAs), specifically *T. asperellum* (PP348038) and *P. fluorescens* (PP348041) were evaluated for their ability to suppress the targeted pathogens. Fifteen kilograms of sterilized topsoil from a groundnut field were filled in 45 x 30 cm earthen pots. Groundnut seeds of a susceptible cultivar (VRI-2) were surface sterilized using 3% NaOCl solution for 30 sec, followed by two washes with sterile water. Five seeds were sown per pot. Biocontrol-based components including these BCAs, fortified lignite fly ash (FL-ash) and Annamalai mixture were tested against LLS and rust. The treatments include talc formulation of *T. asperellum* and *P. fluorescens* (2×10^8 CFU g⁻¹) as seed treatments (10 g/kg) and soil drench after sowing, FL-ash as seed treatment (10 g/kg) and soil drench (40 kg/ha equivalent), Annamalai mixture as a seed treatment (10 ml/kg) and foliar spray (20 L/ha equivalent). Spore suspensions of *P. personata* and *P. arachidis* (10^8 spores/ml) were sprayed on plants in the evening at two-time points with a four-day interval, starting 30 days after sowing (DAS). Agar plate technique was employed to confirm the absence of seed-borne pathogens before sowing. Disease incidence and severity were likely assessed throughout the experiment.

2.6 Field Evaluation of Biocontrol Based Strategy against *P. personata* and *P. arachidis*

A field experiment was carried out in an endemic field from June to September, 2022 in Sivapuri village, Cuddalore, Tamil Nadu. Groundnut cv. VRI-2 was sown in 33 x 13 feet plots. The biocontrol-based components tested in a previous greenhouse experiment were applied here except for pathogen control, following the same schedule. Standard agricultural practices were maintained throughout the season. PDI and

lesion frequency along with growth plant parameters and yield attributes were likely measured.

2.7 Effect of Eco-friendly Amendments on Activity of Defense Enzyme

In a poly house experiment on VRI-2 variety of groundnut (as mentioned in 2.3) subsequent to inoculation with the two target pathogens at 35 days post-sowing, plant samples were obtained at designated time points (1-, 3-, 5- and 7-days post inoculation) from each treatment. Groundnut plant tissues were immediately frozen in liquid N₂ and subsequently homogenized. A sample of 1g mixed with 2 mL of 0.1 M sodium phosphate buffer (pH 7.0) at 4°C. The homogenate was subsequently centrifuged at 10,000 rpm for 20 min. and used for enzyme assays.

Peroxidase (PO) activity was determined by following the method of Srivastava [14] and the enzyme activity was expressed as the change in absorbance/min/g of fresh weight tissue. The polyphenol oxidase (PPO) activity was assayed as per Mayer and Harel [15]. The change in absorbance at 495 nm/min/g of fresh-weight tissue is used to express PPO activity. Phenylalanine Ammonia-Lyase (PAL) was assayed following the procedure of Srivastava (14) and expressed as micromoles (μmol) of trans-cinnamic acid formed per minute per gram of fresh tissue weight (μmol min⁻¹ g⁻¹ fresh weight).

2.8 Statistical Data Analysis

A Completely Randomized Block Design (CRBD) and Randomized Block Design (RBD) with three replicates was used for the pot culture and field experiments respectively. Statistical analysis was performed using SAS software (version 9.4, SAS Institute, Cary, NC, USA) to assess treatment effects. Following a significant analysis of variance (ANOVA), Dunnett's multiple comparison test (DMRT) was employed to identify statistically significant differences (p < 0.05) between treatment means and the control. All the treatments were replicated thrice.

3. RESULTS AND DISCUSSION

3.1 Survey on the Disease Incidence

A substantial variation in disease incidence across the surveyed locations. LLS exhibited a

prevalence ranging from 8.24% to 32.84%, with the highest incidence observed in Mathur. Similarly, rust incidence varied between 7.21% and 30.78%, with Mathur again recording the most severe infection. These findings suggest that geographical factors, cultivar type, and pathogen genetics within the region may significantly influence the severity and distribution of the pathogens [16-18]. The co-infection of both *P. personata* and *P. arachidis* causes severe foliar disease in all major groundnut growing areas of Tamil Nadu.

3.2 Efficacy Biocontrol-Based Components under Pot Culture Conditions

A significant reduction in disease incidence of LLS and rust in plants treated with several biocontrol-based formulations was found. Among the treatments, the combination (seed and soil application) of *T. asperellum*, *P. fluorescens*, fortified lignite fly ash, and Annamalai mixture (denoted as T7) had most significantly lowered the lesion frequency of LLS and disease severity index (PDI). The same treatment also significantly reduced the disease incidence of rust and PDI of rust. Additionally, T7 resulted in a considerable increase in pod yield. The findings also suggested that strategically combined biocontrol-based formulations offer a promising approach for managing LLS and rust diseases, potentially surpassing the efficacy of individual eco-friendly components and even a standard chemical fungicide (Table 1). The microbial consortia could potentially proliferate in the soil and survive throughout the cropping season and protect the crop from harmful pathogenic infection [19,20]. Earlier workers have proven that the native isolate of bioagents could reduce the disease intensity of LLS and rust [21,22]. The biocontrol agents like *Trichoderma* and *Pseudomonas* produce antibiotics, iron-chelating agents, cell wall degrading enzymes and secondary metabolites that act on pathogens to suppress the pathogens' growth effectively [23,24]. Annamalai mixture has been proven as an effective disease suppressor and plant growth promoter [25]. The mixture contains animal excretes that improve the soil health and protect the plants against various diseases by promoting plant growth [13,12].

Table 1. Evaluation of biocontrol-based formulations against late leaf spot and rust under pot culture condition

Treatment	Disease incidence						PDI						Yield g/plant
	Late leaf spot (No. of lesion)			Rust (No. of pustule)			Late leaf spot			Rust			
	50 DAS	70 DAS	90 DAS	50 DAS	70 DAS	90 DAS	50 DAS	70 DAS	90 DAS	50 DAS	70 DAS	90 DAS	
T ₁	0.98 ^d	1.14 ^d	1.32 ^e	0.64 ^e	0.89 ^e	1.27 ^e	12.9 (21.04) ^e	13.6 (21.64) ^e	14.1 (22.05) ^e	10.8 (19.18) ^e	11.2 (19.55) ^e	13.8 (21.80) ^e	53 ^e
T ₂	1.34 ^g	1.52 ^g	2.02 ^h	1.10 ^g	1.48 ^h	1.63 ^h	15.9 (23.49) ^h	16.1 (73.65) ^h	16.9 (24.27) ^h	15.2 (22.94) ^h	16.4 (23.88) ^h	17.7 (24.87) ^h	50 ^h
T ₃	0.63 ^c	1.02 ^c	1.24 ^d	0.49 ^d	0.71 ^d	1.09 ^d	10.3 (18.71) ^d	11.9 (20.17) ^d	13.6 (21.64) ^d	7.0 (15.34) ^d	8.3 (16.74) ^d	11.7 (20.00) ^d	57 ^d
T ₄	1.19 ^e	1.26 ^e	1.57 ^f	0.87 ^f	1.16 ^f	1.32 ^f	14.1 (22.05) ^f	15.3 (23.02) ^f	15.9 (24.27) ^f	12.5 (20.70) ^f	13.0 (21.13) ^f	13.9 (21.89) ^f	53 ^f
T ₅	1.27 ^f	1.44 ^f	1.63 ^g	1.05 ^g	1.39 ^g	1.52 ^g	15.0 (22.78) ^g	15.9 (23.49) ^g	16.3 (23.81) ^g	14.5 (22.38) ^g	15.3 (23.02) ^g	16.1 (23.65) ^g	51 ^g
T ₆	0.42 ^b	0.97 ^c	1.13 ^c	0.37 ^c	0.67 ^c	1.03 ^c	9.7 (18.14) ^c	10.2 (18.62) ^c	12.9 (21.04) ^c	6.2 (14.41) ^c	7.9 (16.32) ^c	10.8 (19.18) ^c	59 ^c
T ₇	0.31 ^a	0.62 ^a	0.98 ^a	0.18 ^a	0.31 ^a	0.60 ^a	7.3 (15.67) ^a	8.9 (17.35) ^a	11.3 (19.64) ^a	5.2 (13.18) ^a	7.1 (15.45) ^a	9.0 (17.45) ^a	69 ^a
T ₈	0.37 ^b	0.84 ^b	1.02 ^b	0.29 ^b	0.53 ^b	0.89 ^b	8.9 (17.35) ^b	9.8 (18.24) ^b	12.6 (20.79) ^b	5.9 (14.05) ^b	7.6 (16.00) ^b	9.9 (18.33) ^b	63 ^b
T ₉	3.11 ⁱ	4.56 ⁱ	6.05 ^j	2.17 ⁱ	3.63 ^j	5.22 ^j	38.3 (38.23) ^j	41.8 (40.28) ^j	42.1 (40.45) ^j	35.5 (36.57) ^j	38.7 (38.46) ^j	41.2 (39.93) ^j	45 ^j
T ₁₀	2.45 ^h	3.26 ^h	3.40 ⁱ	2.03 ^h	3.25 ⁱ	3.47 ⁱ	23.8 (29.19) ⁱ	28.3 (32.13) ⁱ	34.9 (36.21) ⁱ	19.6 (26.27) ⁱ	23.7 (29.13) ⁱ	30.8 (33.70) ⁱ	51 ⁱ

* Means of three replications, Values in the column followed by same letters not differ significantly by DMRT (p=0.05)

T₁ – Talc formulation of *T. asperellum* as seed treatment at 5g/kg of seed, soil application at 10 kg/ha at 25 and 45 DAS, T₂ – Talc formulation of *P. fluorescens* as seed treatment at 10g/kg of seed, soil application at 10 kg/ha at 25 and 45 DAS, T₃ – T₁ +T₂, T₄ – Application of fortified lignite fly ash as seed treatment at 10g/kg of seed, soil application at 40 kg/ha at 25 DAS, T₅ – Application of Annamalai mixture as seed at 10 ml/kg of seed, foliar spray at 20 lit/ha at 25 and 45 DAS, T₆ – T₃ + T₄, T₇- T₆ + T₅, T₈ – Carbendazim seed treatment at 2g/kg, foliar application at 0.1 g/lit at 25 and 45 DAS, T₉ – Inoculated control and T₁₀ – Healthy control

Table 2. Evaluation of various eco-friendly components against *Phaeoisariopsis personata* and *Puccinia arachidis* under field condition

Treatment	Disease incidence						PDI						Yield kg/ha
	Late leaf spot (No. of lesion)			Rust (No. of pustule)			Late leaf spot			Rust			
	50 DAS	70 DAS	90 DAS	50 DAS	70 DAS	90 DAS	50 DAS	70 DAS	90 DAS	50 DAS	70 DAS	90 DAS	
T ₁	0.38 ^c	0.54 ^c	0.72 ^c	0.34 ^c	0.38 ^c	0.43 ^c	10.09 (18.52) ^e	10.79 (19.17) ^e	10.97 (19.34) ^e	6.80 (15.11) ^e	7.14 (15.49) ^e	8.83 (17.28) ^e	1893 ^e
T ₂	0.48 ^d	0.61 ^d	0.79 ^d	0.42 ^d	0.48 ^d	0.53 ^d	11.76 (20.05) ^h	12.13 (20.38) ^h	12.69 (20.86) ^h	8.87 (17.32) ^h	9.24 (17.67) ^h	10.19 (18.61) ^h	1828 ^h
T ₃	0.35 ^b	0.50 ^c	0.69 ^b	0.29 ^c	0.31 ^b	0.39 ^b	9.14 (17.59) ^d	9.39 (17.84) ^d	10.51 (18.91) ^d	5.93 (14.09) ^d	6.94 (15.27) ^d	8.02 (16.45) ^d	1897 ^d
T ₄	0.41 ^c	0.56 ^d	0.75 ^c	0.37 ^d	0.42 ^c	0.47 ^c	10.51 (18.91) ^f	10.93 (19.30) ^f	11.39 (19.72) ^f	7.46 (15.85) ^f	7.91 (16.33) ^f	9.09 (17.54) ^f	1868 ^f
T ₅	0.46 ^d	0.59 ^d	0.77 ^c	0.40 ^d	0.45 ^d	0.49 ^d	11.19 (19.54) ^g	11.78 (20.07) ^g	12.04 (20.30) ^g	8.51 (16.96) ^g	8.89 (17.34) ^g	9.71 (18.15) ^g	1843 ^g
T ₆	0.32 ^b	0.47 ^b	0.66 ^b	0.23 ^b	0.30 ^b	0.37 ^b	8.42 (16.86) ^c	8.96 (17.41) ^c	10.39 (18.80) ^c	5.02 (12.94) ^c	6.27 (14.50) ^c	7.98 (16.40) ^c	1938 ^c
T ₇	0.22 ^a	0.38 ^a	0.51 ^a	0.12 ^a	0.24 ^a	0.31 ^a	7.20 (15.56) ^a	8.14 (16.57) ^a	9.50 (17.95) ^a	3.92 (11.41) ^a	5.67 (13.77) ^a	6.81 (15.12) ^a	2060 ^a
T ₈	0.27 ^a	0.44 ^b	0.56 ^a	0.19 ^b	0.27 ^a	0.33 ^a	8.32 (16.76) ^b	8.89 (17.34) ^b	10.07 (18.50) ^b	4.19 (11.81) ^b	6.05 (14.23) ^b	7.37 (15.75) ^b	1984 ^b
T ₉	1.36 ^e	1.51 ^e	1.68 ^e	1.31 ^e	1.33 ^e	1.38 ^e	14.46 (22.34) ⁱ	16.89 (24.26) ⁱ	18.45 (25.43) ⁱ	10.67 (19.06) ⁱ	12.02 (20.28) ⁱ	18.48 (25.46) ⁱ	1739 ⁱ

* Means of three replications, Values in the column followed by same letters not differ significantly by DMRT ($p=0.05$)
(For treatment descriptions, please refer to the foot note in the Table 1. T₁₀ was not present in field experiment)




Fig. 1. Induction of defence enzymes by eco-friendly components in groundnut plants against *Phaeoisariopsis personata* and *Puccinia arachidis*. (a) Peroxidase, (b) Polyphenol oxidase, (c) Phenylalanine ammonia-lyase. (For treatment descriptions, please refer to the foot note in the Table 1)

3.3 Evaluation of Biocontrol-Based Formulations Components under Field Conditions

The results of the pot culture experiments were confirmed in the field. Again, the treatment (T7) demonstrated the most significant reductions in disease incidence for both LLS and rust. Treatment T7 displayed consistently lower lesion frequency and percent disease index (PDI) of LLS and rust diseases at 50, 70, and 90 DAS, respectively. Additionally, T7 resulted in the highest yield. The combined application of these biocontrol-based formulations was proven to be a more effective approach for managing LLS and rust in groundnut crops compared to individual applications or a chemical fungicide control (Table 2). Combined application of bioproducts have been known to control diseases effectively by employing more than one mechanism of direct or indirect pathogen suppression [26,27]. Such approaches include combination of organic products with biocontrol agents [13], a combination of plant products with biocontrol agents [28,29] and a combination of fungicides with biocontrol agents [30]. These bioproducts also promoted plant growth significantly and increased the yield of the crop [31]. The major reason for the increased plant growth and yield parameters have been attributed to the

significant induction of plant growth-promoting substances like auxin, gibberellic acid and abscisic acid induced by the bioproducts [32,33]. The lignite fly ash used as carrier material contains a trace amount of minor nutrients like Ca, K, and Si [34]. These minor nutrients trigger the plant defense response and protect the plants [35].

3.4 Defense Enzyme Activity in Groundnut Plants against *P. personata* and *P. arachidis*

Combined application of bioproducts (treatment T7) resulted in the most significant induction of activities of the defense enzymes, PO, PPO and PAL. This peak induction occurred 5 days post-challenge inoculation with the respective pathogens. Interestingly, plants solely inoculated with the pathogens also exhibited elevated levels of defense enzymes compared to the healthy control group up to the fifth day (Fig. 1). This stimulation of defense enzymes suggests that the combined application of these biocontrol components possibly are involved in systemic resistance in groundnut plants, potentially improving their tolerance against these fungal diseases [36]. Peroxidases are a diverse group of enzymes that detoxify reactive oxygen species (ROS) produced by pathogens [37]. These ROS

are highly reactive and can damage cellular components [38]. By detoxifying ROS, peroxidases prevent oxidative damage to proteins, lipids, and nucleic acids, maintaining cellular integrity and promoting plant survival [39]. PPO catalyzes the oxidation of phenolic compounds, leading to the formation of quinones. These quinones disrupt pathogen membranes and inactivates their enzymes, directly inhibiting infection [39].

4. CONCLUSION

Surveys of late leaf spot and rust in groundnut crops across the Cuddalore district, Tamil Nadu, India revealed significant geographical variation in disease incidence, with Mathur exhibiting the highest infection rates for both diseases. A combination of *Trichoderma asperellum*, *Pseudomonas fluorescens*, fortified lignite fly ash and Annamalai mixture most significantly reduced the disease incidence for both LLS and rust and increased groundnut yield under both pot culture and field conditions. Induction of defense enzyme activity in groundnut plants treated with these biocontrol-based components suggests a potential mechanism for disease resistance.

ACKNOWLEDGEMENTS

We thank the Head of the Department, Faculty of Agriculture, Annamalai University for encouragement and support for this investigation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Pal KK, Dey R, Tilak KV. Fungal diseases of groundnut: Control and future challenges. Future challenges in crop protection against fungal pathogens. 2014: 1–29. DOI: 10.1007/978-1-4939-1188-2
2. Joshi E, Sasode DS, Singh N, Chouhan N. Diseases of groundnut and their control measures. Biotica Res. Today. 2020;2(5): 232–237.
3. Kishore GK, Pande S. Integrated management of late leaf spot and rust diseases of groundnut (*Arachis hypogaea* L.) with *Prosopis juliflora* leaf extract and chlorothalonil. Int. J. Pest Manag. 2005; 51(4):325–332. DOI: 10.1080/09670870500404567
4. Subrahmanyam P, Rao VR, McDonald D, Moss JP, Gibbons RW. Origins of resistances to rust and late leaf spot in peanut (*Arachis hypogaea*, Fabaceae). Econ. Bot. 1989;43(4):444–455.
5. Cole DL. Rust disease of groundnut in southern Africa: Present situation and possible interactions with other groundnut foliar diseases. In Citation: ICRISAT (International Crops Research Institute for the Semi-Arid Tropics). 1987. Groundnut rust disease. Proceedings of a Discussion Group Meeting, 24-28 Sep 1984, ICRISAT Center, India. Patancheru, AP 502324, India: ICRISAT; 1987.
6. Goswami SK, Singh V, Chakdar H, Choudhary P. Harmful effects of fungicides-Current status. Int. J. Agric. Environ. Biotechnol. 2018;11:1011–1019. ISSN: 1974–1712.
7. Boudh S, Singh JS. Pesticide contamination: Environmental problems and remediation strategies. In: Bharagava R, Chowdhary P. (eds) Emerging and Eco-Friendly Approaches for Waste Management. Springer, Singapore. 2019: 245–269. Available: https://doi.org/10.1007/978-981-10-8669-4_12
8. Verma DK, Pandey AK, Mohapatra B, Srivastava S, Kumar V, Talukdar D, Yulianto R, Zuan AT, Jobanputra AH, Asthir B. Plant growth-promoting rhizobacteria: An eco-friendly approach for sustainable agriculture and improved crop production. In microbiology for sustainable agriculture, soil health, and environmental protection. Apple Academic Press. 2019: 3–80.
9. Raaijmakers JM, Mazzola M. Diversity and natural functions of antibiotics produced by beneficial and plant pathogenic bacteria. Annu. Rev. Phytopathol. 2012;50:403–424. Available: <https://doi.org/10.1146/annurev-phyto-081211-172908>.
10. Adam M, Heuer H, Hallmann J. Bacterial antagonists of fungal pathogens also control root-knot nematodes by induced systemic resistance of tomato plants. PloS One. 2014;9(2):e90402. Available: <https://doi.org/10.1371/journal.pone.0090402>
11. Subrahmanyam P, McDonald D, Waliyar F, Reddy LJ, Nigam SN, Gibbons RW, Rao VR, Singh AK, Pande S, Reddy PM, Rao PS. Screening methods and sources of

- resistance to rust and late leaf spot of groundnut. Information Bulletin no. 47. International Crops Research Institute for the Semi-Arid Tropics; 1995:1–20.
12. Thamamarai Selvi M, Kuruचेve V, Darwin Christdhas Henry I, Sutha Raja Kumar R, Jaiganesh V. Effect of animal excreta on mycelial growth of *Pyricularia oryzae*. J. Pharmacogn. Phytochem. 2019;8(3):3373–3375.
 13. Newton DK, Christopher DJ. Evaluation of Bio Inoculants Fortified Lignite Fly Ash (LFA) against Sheath Blight (*Rhizoctonia solani* Kuhn) and Sheath Rot (*Saracladium oryzae* (Sawada) Games and Haksworth) diseases of rice in cauvery delta region of Tamil Nadu. J. Biol. Cont. 2017;31(3):172–183.
DOI:<https://doi.org/10.18311/jbc/2017/15848>
 14. Srivastava SK. Peroxidase and poly-phenol oxidase in *Brassica juncea* plants infected with *Macrophomina phaseolina* (Tassai) Goid. and their implication in disease resistance. J. Phytopathol. 1987;120(3):249–254.
Available:<https://doi.org/10.1111/j.1439-0434.1987.tb04439.x>
 15. Mayer AM, Harel E. Polyphenol oxidases in plants. Phytochem. 1979;18(2):193–215.
Available:[https://doi.org/10.1016/0031-9422\(79\)80057-6](https://doi.org/10.1016/0031-9422(79)80057-6)
 16. Shaw MW, Osborne TM. Geographic distribution of plant pathogens in response to climate change. Plant Pathol. 2011; 60(1):31–43.
Available:<https://doi.org/10.1111/j.1365-3059.2010.02407.x>
 17. West JS, Townsend JA, Stevens M, Fitt BD. Comparative biology of different plant pathogens to estimate effects of climate change on crop diseases in Europe. Eur. J. Plant Pathol. 2012;133:315–331.
Available:<https://doi.org/10.1007/s10658-011-9932-x>
 18. Jain A, Sarsaiya S, Wu Q, Lu Y, Shi J. A review of plant leaf fungal diseases and its environment speciation. Bioengineered. 2019;10(1):409–424.
Available:<https://doi.org/10.1080/21655979.2019.1649520>.
 19. Vishwakarma K, Kumar N, Shandilya C, Mohapatra S, Bhayana S, Varma A. Revisiting plant–Microbe interactions and microbial consortia application for enhancing sustainable agriculture: A review. Front. Microbiol. 2020;11:560406.
Available:<https://doi.org/10.3389/fmicb.2020.560406>
 20. Mondal S, Halder SK, Yadav AN, Mondal KC. Microbial consortium with multifunctional plant growth-promoting attributes: Future perspective in agriculture. In: Yadav A, Rastegari A, Yadav N, Kour D. (eds) Advances in plant microbiome and sustainable agriculture. Microorganisms for Sustainability, Springer, Singapore. 2020;20:219–258.
Available:https://doi.org/10.1007/978-981-15-3204-7_10
 21. Dey U, Dhutraj DN, Suryawanshi AP, DAS A, Das SP. Comparative economics of spray scheduling for the management of late leaf spot of groundnut incited by *Phaeoisariopsis personata*. Indian Phytopathol. 2016;69(2):149–154.
 22. Kumhar DR, Meena AK, Meena PN. Efficacy of bio agents and fungicides against collar rot and early leaf spot of groundnut. Indian Phytopathol. 2018;71: 549–554.
Available:<https://doi.org/10.1007/s42360-018-0080-5>
 23. Prashar P, Kapoor N, Sachdeva S. Biocontrol of plant pathogens using plant growth promoting bacteria. Sustainable Agriculture Reviews. 2013;12:319–360.
Available:https://doi.org/10.1007/978-94-007-5961-9_10
 24. Lahlali R, Ezrari S, Radouane N, Kenfaoui J, Esmaeel Q, El Hamss H, Belabess Z, Barka EA. Biological control of plant pathogens: A global perspective. Microorganisms. 2022;10(3):596.
Available:<https://doi.org/10.3390/microorga-nisms10030596>
 25. Lananya J, Kave Newton DE, Jaiganesh V, John Christopher D. Evaluation of ecofriendly components viz., fortified lignite fly ash, Annamalai mixture and bio-inoculants against groundnut stem rot diseases caused by *Sclerotium rolfsii* Sacc. 2019;865-869.
 26. Lastochkina O, Seifikalhor M, Aliniaiefard S, Baymiev A, Pusenkova L, Garipova S, Kulabuhova D, Maksimov I. *Bacillus* spp.: Efficient biotic strategy to control postharvest diseases of fruits and vegetables. Plants. 2019;8(4):97.
Available:<https://doi.org/10.3390/plants8040097>
 27. Nguyen SD, Trinh TH, Tran TD, Nguyen TV, Chuyen HV, Ngo VA, Nguyen AD.

- Combined application of rhizosphere bacteria with endophytic bacteria suppresses root diseases and increases productivity of black pepper (*Piper nigrum* L.). Agriculture. 2020;11(1):15. Available: <https://doi.org/10.3390/agricultur e11010015>
28. Muthukumar A, Eswaran A, Nakkeeran S, Sangeetha G. Efficacy of plant extracts and biocontrol agents against *Pythium aphanidermatum* inciting chilli damping-off. Crop Protect. 2010;29(12):1483–1488. Available: <https://doi.org/10.1016/j.cropro.2 010.08.009>
29. Akila R, Rajendran L, Harish S, Saveetha K, Raguchander T, Samiyappan R. Combined application of botanical formulations and biocontrol agents for the management of *Fusarium oxysporum* f. sp. *ubense* (Foc) causing Fusarium wilt in banana. Biol. control. 2011;57(3):175–183. Available: <https://doi.org/10.1016/j.biocontr ol.2011.02.010>
30. Santos FM, Viera LS, Camargo DP, Muniz MF, Costa IF, Guedes JV, Santos JR, Silva JC. Integrating a Bacillus-based product with fungicides by foliar application to protect soybean: A sustainable approach to avoid exclusive use of chemicals. Pest Manag. Sci. 2022;78(11):4832–4840. Available: <https://doi.org/10.1002/ps.7104>
31. Radwan MA, Farrag SA, Abu-Elamayem MM, Ahmed NS. Biological control of the root-knot nematode, *Meloidogyne incognita* on tomato using bioproducts of microbial origin. Appl. Soil Ecol. 2012;56:58–62. Available: <https://doi.org/10.1016/j.apsoil.20 12.02.008>
32. Khalid A, Arshad M, Shaharoona B, Mahmood T. Plant growth promoting Rhizobacteria and sustainable agriculture. In: Khan M, Zaidi A, Musarrat J. (eds) Microbial strategies for crop improvement. Springer, Berlin, Heidelberg. 2009:133–160. Available: https://doi.org/10.1007/978-3- 642-01979-1_7.
33. Maheshwari DK, Dheeman S, Agarwal M. Phytohormone-Producing PGPR for sustainable agriculture. In: Maheshwari D. (eds) Bacterial metabolites in sustainable agroecosystem. Sustainable Development and Biodiversity, Springer, Cham. 2015;12:159–182. Available: https://doi.org/10.1007/978-3- 319-24654-3_7
34. Ram LC, Srivastava NK, Tripathi RC, Jha SK, Sinha AK, Singh G, Manoharan V. Management of mine spoil for crop productivity with lignite fly ash and biological amendments. J. Environ. Manag. 2006;79(2):173–187. Available: <https://doi.org/10.1016/j.jenvman. 2005.06.008>
35. Omidvari M, Abbaszadeh-Dahaji P, Hatami M, Kariman K. Biocontrol: A novel eco-friendly mitigation strategy to manage plant diseases. Plant Stress Mitigators. 2023:27–56. Available: <https://doi.org/10.1016/B978-0- 323-89871-3.00020-3>
36. Kawano T. Roles of the reactive oxygen species-generating peroxidase reactions in plant defense and growth induction. Plant Cell Rep. Plant Cell Rep. 2003;21:829–837. Available: <https://doi.org/10.1007/s00299- 003-0591-z>
37. Almagro L, Gómez Ros LV, Belchi-Navarro S, Bru R, Ros Barceló A, Pedreño MA. Class III peroxidases in plant defense reactions. J. Exp. Bot. 2009;60(2):377–390. Available: <https://doi.org/10.1093/jxb/ern27 7>
38. Madkour LH. Function of reactive oxygen species (ROS) inside the living organisms and sources of oxidants. Pharm. Sci. Anal. Res. J. 2019;2:180023. ISSN: 2640–6659.
39. Sharma P, Sharma P, Arora P, Verma V, Khanna K, Saini P, Bhardwaj R. Role and regulation of ROS and antioxidants as signaling molecules in response to abiotic stresses. In Plant Signaling Molecules, Woodhead Publishing; 2019.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/116170>