



Determining the Effect of Growing Media on Growth and Flowering of Rose

Lairingdi Bodo ^a, Sanjib Sharma ^b, Kankana Deka ^a,
Sangita Mahanta ^a, Bhaskarjyoti Sarma ^{c*},
Rupak Kumar Nath ^c, Nilay Bora ^d and Kaushik Das ^e

^a Department of Horticulture, Assam Agricultural University, Jorhat – 785013, Assam, India.

^b Department of Floriculture & Landscape Architecture, College of Horticulture & FSR, Lakhoupur, Nalbari – 781335, Assam, India.

^c SCS College of Agriculture, AAU, Rangamati, Chapar, Dhubri – 783376, Assam, India.

^d Department of Soil Science, Assam Agricultural University, Jorhat – 785013, Assam, India.

^e Department of Crop Physiology, Assam Agricultural University, Jorhat – 785013, Assam, India.

Authors' contributions

This work was carried out in collaboration among all Authors. Authors LB, SS and KD designed the study and conducted the experimentation. Authors SM, BS and RKN did the analysis and wrote the first draft of the manuscript. Author BS performed the literature searches. Authors NB and KD supervised the investigation.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i92895>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123884>

Original Research Article

Received: 14/07/2024

Accepted: 16/09/2024

Published: 18/09/2024

ABSTRACT

The present study aimed to determine the effect of Growing Media on Growth and Flowering of Rose. The study investigation was carried out in the Experimental Farm, Dept. of Horticulture, Assam Agricultural University, Jorhat for the year 2022-2023 with a view to assess the effect of

*Corresponding author: E-mail: bhaskarjyoti.sarma@aau.ac.in;

Cite as: Bodo, Lairingdi, Sanjib Sharma, Kankana Deka, Sangita Mahanta, Bhaskarjyoti Sarma, Rupak Kumar Nath, Nilay Bora, and Kaushik Das. 2024. "Determining the Effect of Growing Media on Growth and Flowering of Rose". *Journal of Experimental Agriculture International* 46 (9):985-996. <https://doi.org/10.9734/jeai/2024/v46i92895>.

growing media on growth and flowering of rose (Variety Revival). The experiment was carried out in Completely Randomized Design consisting of 9 treatment combination viz., T₁ {Sandy loam + Sand + FYM @ 1:1:1 (w/w)}, T₂ {Loam soil + Sand + FYM @ 1:1:1 (w/w)}, T₃ {Clay loam + Sand + FYM @ 1:1:1 (w/w)}, T₄ {Sandy loam + Sand + Vermicompost @ 1:1:1 (w/w)}, T₅ {Loam soil + Sand + Vermicompost @ 1:1:1 (w/w)}, T₆ {Clay loam + Sand + Vermicompost @ 1:1:1 (w/w)}, T₇ {Sandy loam + Sand + Vermicompost (10% substituted with Cocopeat) @ 1:1:1 (w/w)}, T₈ {Loam soil + Sand + Vermicompost (10% substituted with Cocopeat) @ 1:1:1 (w/w)}, T₉ {Clay loam + Sand + Vermicompost (10% substituted with Cocopeat) @ 1:1:1 (w/w)}, which were replicated three times. The test crop positively responded to the growing media and the highest performance was observed for the treatment T₈. Stem girth, number of nodes, intermodal length, leaf area and leaf area index, days taken to bud visibility, bud sprouting, half bloom and full bloom, length of stalk and diameter of bud, number of flowers, number of petals per flower and flowering duration (self-life) and diameter of flower were significantly enhanced by the treatment T₈.

Keywords: Rose; vermicompost; cocopeat; loam; Sand; FYM.

1. INTRODUCTION

The Rose, scientifically known as *Rosa indica*, is widely recognized as the "Queen of Flowers" due to its unique shape, varying sizes, alluring colours, delightful fragrance and has many practical applications [1]. Genus Rose comprises of more than 200 species and 18,000 cultivars that has since been used as vibrant plant in floriculture industry [2]. Nowadays, the potted production of rose plants is gaining popularity as potted plants are a good source for decreasing air pollution provide freshness in indoor premises. Important growing substrates that are ideally employed for the container production of both annual and perennial ornamental plants include garden soil, sand, vermicompost, cocopeat, FYM, and coir husk [3]. These materials help to boost the microbial activity of the soil which is necessary to improve and meet the nutritional needs and also decrease the nitrogen losses of the crop [4].

A good growing media is very important for the plant to be adequately anchored so that the supply of nutrients and water, as well as gas exchange between the roots and the atmosphere outside the growing medium is not hampered. A highly sought-after horticultural by-product utilized as a substrate for the growth of numerous floricultural crops is coco peat, which is made after the fibre from coconut husk is extracted because of its pH, electrical conductivity and other chemical properties [5].

It is necessary to amend cocopeat with sufficient organic composts in order to maintain the ideal

level of nutrients in growing media for the successful growth of plants. Vermicomposting is a technique for obtaining organic matter that is well known for producing highly stable, non-thermophilic products with a lot of potential as soil amendments [6]. Large levels of humic substances found in vermicomposts have some effects on plant growth that are comparable to those of soil-applied with plant growth regulators [6]. It is with this background that this study aimed to assess the effect of growing media on these aspects by analyzing some growth and flowering parameters in the rose plant.

2. MATERIALS AND METHODS

A field experiment on Effect of growing media on growth and flowering of rose was conducted during 2022-23 at the Experimental Farm, Department of Horticulture, Assam Agricultural University, Jorhat.

For this experiment, three different types of soil were utilized viz., sandy loam, loam and clay loam. The textural class of the soil samples were estimated by feel method for determining soil texture. Soil texture refers to the ratio of sand, silt, and clay present in the soil. In the experiment, Sand, Farmyard Manure, Vermicompost and Cocopeat were used.

2.1 Experimental Details

The details of the experiment with respect to treatment, design, plot size are given below:

List 1. Details of the experiment with respect to treatment, design, plot size

Period of Experiment	: 2022-2023
Crop Variety	: Revival
Design of the experiment	: CRD (Completely Randomized Design)
Number of replications	: 3
Number of treatments	: 9
Number of pots per treatment	: 5
Pot size	: 30 cm diameter and 28 cm height
Total number of pots	: 135
Total area	: 500 m ²

2.2 Treatment Details

The experiment consisted of 9 (nine) treatment combinations comprised of 3 (three) different types of soil incorporated with sand, FYM, vermicompost and cocopeat. The details of the treatment combinations are given in List 2:

2.2.1 Treatments

The soils used for the pot experiment were collected from three different locations of the Experimental Farm, Department of Horticulture, Assam Agricultural University, Jorhat. Each soil was mixed thoroughly and stones and debris were removed and mixed homogeneously with sand. Each soil sample was mixed with FYM, vermicompost and cocopeat in different ratios respectively.

2.3 Filling Up of Pot

For the pot experiment, 5 kg capacity pots were used. The pots were properly cleaned and sun-dried. Holes were made and a few broken pieces of bricks were placed at the bottom of the pots. Then the pots were filled with soil mixture and proper labelling of each pot was done.

2.4 Collection of Planting Material and Planting

35 days old grafted planting material were collected from a Government accredited nursery and were planted in individual pots on 19th August, 2022. Watering was done immediately after the completion of planting.

2.5 Crop Protection

Saaf @ 1.5gm/l and Blitox @ 2gm/l was sprayed alternately as a prophylactic measure against dieback at an interval of 15 days. 2-3 spray of Blitox @ 2gm/l and Imidacloprid @ 0.5 ml/l was sprayed at an interval of 10 days for insect attacks. 21

2.6 Observations Recorded

Growth characters:

- Plant height (cm)
- Number of leaves per plant
- Leaf area per plant (cm²) was measured with a Laser area Meter Model No: C1-203, C1-203A (Make-CID, Inc., USA). Leaf area index (LAI)
- The leaf area index was calculated according to Evans (1972) as follows:
LAI = Total leaf area of the plant / Total ground area covered by the plant
- Stem girth (cm)
- Intermodal length (cm)
- Plant spread (cm)

Flower characters:

- Days taken to bud visibility from planting
- Days taken to bud sprouting
- Days taken to half bloom
- Days taken to full bloom
- Bud length (cm)
- Bud diameter (cm)
- Stalk length (cm)
- Stalk diameter (cm)
- Flower diameter (cm)
- Number of flowers per plant
- Number of petals per flower
- Self-life (duration of the flowering) (days)

3. RESULTS AND DISCUSSION

3.1 Effect of Growing Media on Soil Physico-Chemical Properties (After Mixing)

Results on soil analysis after mixing showed a significant variation between the treatments in respect of soil pH, organic carbon and available N, P and K in the growing media.

List 2. Details of the treatment combinations

Notation	Treatment
Treatment 1 (T ₁)	Sandy loam + Sand + FYM @ 1:1:1 (w/w)
Treatment 2 (T ₂)	Loam soil + Sand + FYM @ 1:1:1 (w/w)
Treatment 3 (T ₃)	Clay loam + Sand + FYM @ 1:1:1 (w/w)
Treatment 4 (T ₄)	Sandy loam + Sand + Vermicompost @ 1:1:1 (w/w)
Treatment 5 (T ₅)	Loam soil + Sand + Vermicompost @ 1:1:1 (w/w)
Treatment 6 (T ₆)	Clay loam + Sand + Vermicompost @ 1:1:1 (w/w)
Treatment 7 (T ₇)	Sandy loam + Sand + Vermicompost (10% substituted with Cocopeat) @ 1:1:1 (w/w)
Treatment 8 (T ₈)	Loam soil + Sand + Vermicompost (10% substituted with Cocopeat) @ 1:1:1 (w/w)
Treatment 9 (T ₉)	Clay loam + Sand + Vermicompost (10% substituted with Cocopeat) @ 1:1:1 (w/w)

It was found that the treatment T₁ {Sandy loam + Sand + FYM @ 1:1:1 (w/w)} exhibited the highest soil pH of 5.91 (Table 1). The probable reason for this may be because of the application of FYM in sandy loam soil which is responsible for the moderately acidic pH as pH of FYM is higher than that of any other organic inputs amended in the growing media treatments. Similar finding was reported by [7]. However, highest organic carbon of 23.20 g kg⁻¹ (Table 1), available Phosphorus of 23.15 kg ha⁻¹ and Potassium of 152.54 kg ha⁻¹ (Table 2) was recorded in treatment T₉. This might be because of the presence of cocopeat in the media containing clay loam soil as cocopeat has organic matter of 85-90 % and as soil carbon is a component of soil organic matter leading to higher organic carbon in the growing media. Similar findings were reported by [8]. Organic amendments can significantly enhance soil organic C and provide considerable effects on soil microbes, nutrient availability, and uptake for plant [4]. The probable reason for higher P and K content might be because of the presence of vermicompost and cocopeat in the media both containing adequate quantities of N, P, K and several micronutrients essential for plant growth [9]. The probable reason for the increase in the available

potassium content might be due to the release of potassium from the organic inputs and also release of potassium [10]. The increase in available phosphorus in soil might be due to the incorporation of compost which is enriched with PSB and other microorganism helping in phosphate solubilizing and mineralization process thus increases the soluble form of Phosphorus [11].

The available nitrogen of 309.42 kg ha⁻¹ (Table 2) was recorded highest in treatment T₆ {Clay loam + Sand + Vermicompost @ 1:1:1 (w/w)}. This might be because of the presence of vermicompost in clay loam soil which is rich in nitrogen leading to higher nitrogen availability of the growing media. Vermicompost is an organic amendment that is rich in nutrients and microorganisms and is produced as a result of the interactions between earthworms and microorganisms during the decomposition of organic materials. It is a stabilized, finely divided peat-like substance that has a low C:N ratio, high porosity and water-holding capacity. The majority of the nutrients are present in forms that plants may easily absorb as reported by [12]. These results are in close conformity with the findings of [13].

Table 1. pH and organic carbon content of different media (24 hours after mixing)

Treatment	pH	Organic carbon (g kg⁻¹)
Treatment 1	5.91	15.63
Treatment 2	5.72	16.33
Treatment 3	5.68	16.53
Treatment 4	5.23	11.27
Treatment 5	5.20	12.03
Treatment 6	5.12	15.27
Treatment 7	5.12	20.20
Treatment 8	5.11	20.40

Treatment	pH	Organic carbon (g kg ⁻¹)
Treatment 9	5.05	23.20
S. Ed. (±)	0.12	2.09
C.D. _{P=0.05}	0.25	4.39

Table 2. Available nutrient content of different media (24 hours after mixing)

Treatment	Available nutrients (kg ha ⁻¹)		
	N	P	K
Treatment 1	269.83	15.68	112.34
Treatment 2	271.79	16.05	132.53
Treatment 3	275.97	16.43	142.50
Treatment 4	292.69	17.92	118.57
Treatment 5	301.05	18.67	133.69
Treatment 6	309.42	19.79	148.51
Treatment 7	280.15	21.65	135.15
Treatment 8	284.33	22.03	141.23
Treatment 9	288.51	23.15	152.54
S. Ed. (±)	12.52	1.19	3.92
C.D. _{P=0.05}	NS	2.50	8.23

Table 3. Plant height (cm) and number of leaves in different growth stages

Treatment	Plant height (cm)			No. of leaves		
	30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP
Treatment 1	15.00	15.16	24.33	7.67	12.33	26.47
Treatment 2	23.33	21.50	32.70	9.67	20.67	35.03
Treatment 3	24.16	26.50	35.53	10.37	23.03	36.07
Treatment 4	19.66	20.83	32.07	8.00	16.07	29.67
Treatment 5	21.33	23.16	32.17	9.00	19.77	32.73
Treatment 6	28.16	31.00	40.67	13.07	25.04	38.37
Treatment 7	27.83	29.00	41.73	12.00	22.07	38.00
Treatment 8	29.50	34.33	42.10	14.40	26.83	39.10
Treatment 9	24.66	28.50	39.43	10.97	23.67	36.40
S. Ed. (±)	2.05	2.75	3.18	1.19	2.05	3.04
C.D. _{P=0.05}	4.31	5.79	6.68	2.50	4.30	6.38

3.2 Effect of Growing Media on Growth

Data showing the plant height (cm) and no. of leaves in different growth stages of rose i.e., at 30 days, 60 days, and 90 days of planting are presented in Table 3. From the investigation it was found that there were significant differences among the treatments and also noticed that there was gradual increase in height of the plants of different treatments with the growth advancement. The treatment containing loam soil, sand, vermicompost and cocopeat have been found to increase the plant height, leaf number, and leaf area at vegetative stage and also decrease in the number of days required for flowering. After 30, 60 and 90 days of planting, the tallest plant height were recorded in T₈ with 29.50, 34.33 and 42.10 cm, respectively (Table 3).

Similarly, after 30, 60 and 90 days of planting, the maximum number of leaves (14.40, 26.83 and 39.10) were recorded in T₈ and minimum number of leaves were recorded in T₁. This indicates that the response of rose was positive to the growing media inputs used for this pot mixture. Similar results have been reported by [14] and [15] in rose. The possible reason for better growth characters might be due to the presence of vermicompost which significantly augmented soil enzymatic activities and beneficial microbes which ultimately led to an increase in a number of growth and flowering indices, including plant height, number of leaves, leaf area, plant spread, stem diameter and early flowering. These findings are in conformity with [16] in rose.

Table 4. Stem girth (cm) and number of nodes in different growth stages

Treatment	Stem girth (cm)			No. of nodes		
	30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP
Treatment 1	0.43	0.43	0.50	5.43	17.00	22.47
Treatment 2	0.50	0.56	0.60	6.40	23.37	27.33
Treatment 3	0.53	0.60	0.66	8.00	24.13	28.37
Treatment 4	0.46	0.50	0.56	10.67	18.70	24.17
Treatment 5	0.50	0.53	0.56	7.70	21.40	26.10
Treatment 6	0.66	0.70	0.73	13.33	24.67	29.73
Treatment 7	0.73	0.83	0.86	9.00	30.10	33.43
Treatment 8	0.86	0.90	0.93	11.67	31.70	36.37
Treatment 9	0.63	0.66	0.73	10.03	24.07	29.00
S. Ed. (\pm)	0.05	0.06	0.05	2.19	2.73	2.18
C.D. _{P=0.05}	0.12	0.13	0.12	4.60	5.78	4.58

The reason behind the increase in the plant height may be due to the presence of cocopeat and vermicompost in the media, which considerably improved the aeration, water holding capacity and nutrients uptake by the root system. Similar results were recorded by [17] in rose.

Likewise, the reason behind the increased leaves number might be attributed to the increased availability of N in the media. High nitrogen concentration in the media will cause the plant to produce a lot of leaves because nourishment is a key factor in growing more leaves. The increase in number of leaves as a result of vermicompost application are in accordance with the findings of [18] and [19] in rose. Likewise, the maximum stem girth of 0.93 cm was recorded in the treatment T₈ (Table 4). The reason behind the increase in stem girth might be due to the cocopeat application which increased the readily available nitrogen in the media, which enhanced plant metabolic activities and thus increased various stem girth. The rise in organic carbon content can enhance the activity of helpful soil microorganisms and improve soil fertility by increasing the accessibility of nutrients for plant growth [20]. It also might be due to the increase

in humic materials and plant growth promoting substances by application of vermicompost that ultimately promoted the plant growth. These results are in harmony with the findings of [21] and [15] in rose. The maximum number of nodes was recorded in T₈ (36.37). This might be because of the increased plant spread and the overall growth of different parameters. Vermicompost gives plants the right amount of nutrients, which results in more carbohydrates absorption and results in better utilisation of nutrients. These results are in close conformity with the findings of [16] in rose.

The maximum inter-nodal length of 4.83 cm was recorded in the treatment T₈ (Table 5). This might be because of the increased plant height. The use of vermicompost and cocopeat together had a significant impact on plant growth parameters. This is likely because the combination of both factors improved the physical conditions of the media and provided better nutrition. Scientists [22] and [23] suggested that when mixed with vermicompost, coir dust may offer a superior growth medium for establishing plants despite being low in nutrients. Similar results were recorded by [17] in rose.

Table 5. Inter-nodal length (cm) and plant spread (cm) in different growth stages

Treatment	Inter-nodal length (cm)			Plant spread (cm)		
	30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP
Treatment 1	1.63	1.60	2.86	11.37	15.23	22.70
Treatment 2	2.50	3.20	4.10	15.40	21.33	35.17
Treatment 3	2.60	4.03	4.20	17.83	22.43	37.47
Treatment 4	2.06	2.76	3.50	19.50	20.44	29.73
Treatment 5	2.16	2.83	4.03	13.17	20.37	34.80
Treatment 6	3.80	3.80	4.40	24.07	30.33	41.40
Treatment 7	3.03	4.16	4.70	22.01	27.43	38.44
Treatment 8	4.40	4.20	4.83	27.37	32.47	41.73

Treatment	Inter-nodal length (cm)			Plant spread (cm)		
	30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP
Treatment 9	2.83	3.86	4.36	20.00	25.40	38.70
S. Ed. (\pm)	0.41	0.32	0.29	4.50	2.92	1.87
C.D. _{P=0.05}	0.86	0.68	0.63	9.45	6.13	3.92

Table 6. Leaf area and Leaf Area Index

Treatment	Leaf area (cm ²)	Leaf Area Index
Treatment 1	18.76	2.63
Treatment 2	24.20	3.41
Treatment 3	25.73	3.47
Treatment 4	23.36	2.77
Treatment 5	23.43	2.80
Treatment 6	29.20	3.86
Treatment 7	31.60	4.37
Treatment 8	34.10	4.77
Treatment 9	29.16	3.79
S. Ed. (\pm)	2.33	0.22
C.D. _{P=0.05}	4.89	0.48

The plant spread records furnished was found to increase at all the growth stages. The maximum plant spread of 41.73 cm was recorded in the treatment T₈. The plant spread may have been boosted by improved physiochemical properties of the growing media, such as lower bulk density, higher total porosity, increased water retention, and greater nitrogen availability. Another contributing factor may be the enhanced air circulation in the soil resulting from the incorporation of cocopeat into the media. The above results are in line with the findings of [24] in petunia.

Table 6 depicted that the maximum leaf area of 34.10 cm² was recorded in the treatment T₈ {Loam soil + Sand + Vermicompost (10% substituted with Cocopeat) @ 1:1:1 (w/w)}. One reason for the increase in leaf area may be due to an increase in the number of leaves. This increase helps maintain a high source-sink relationship, as there is an enhanced photosynthetic area. These findings agree with the findings of [25] in rose and [18] in rose. The maximum LAI of 4.77 was recorded in the treatment T₈. The increase in leaf area index may be attributed to a sufficient supply of nitrogen and other nutrients that play a vital role in plant metabolism, the synthesis of carbohydrates, and the activity of phytohormones. The above results are in line with the findings of [26] in pistachio seedlings.

3.3 Effect of Growing Media on Flowering

The minimum days taken to bud visibility and to bud sprouting of 39.10 and 9.40 respectively

(Table 7) was recorded in the treatment T₈. It is possible that the plant's rapid growth in the media, coupled with its quick absorption of nutrients and water, contributes to its early production. This may be attributed to an increase in protein formation from carbohydrate deposits in its vegetative part, as well as in protoplast formation. Similar result was also found by [27]. The possible reason for early bud sprouting maybe attributed to the accumulation of more photosynthates in the plants resulting in early bud sprouting. Similar result was also found by [21].

Likewise, the minimum days taken to half bloom of 12.33 and days taken to full bloom from bud sprouting of 19.60 was recorded in the treatment T₈ (Table 8). This observation has similarity with the findings of [28] who reported early flowering in lilies and pelargonium, when these were grown in media containing cocopeat.

It was observed that the treatment T₇ {Sandy loam + Sand + Vermicompost (10% substituted with Cocopeat) @ 1:1:1 (w/w)} showed the highest length of bud (3.53 cm) (Table 9). The reason for this is may be the use of vermicompost, which can enhance the presence of growth-promoting chemicals and nutrients in the soil-plant system. This can lead to better absorption of nutrients and an increase in the accumulation of photosynthates within the plant. Similar findings were reported by [14] in rose cv. Gladiator.

Table 7. Days taken to bud visibility and days taken to bud sprouting

Treatment	Days taken to bud visibility	Days taken to bud sprouting
Treatment 1	74.43	16.01
Treatment 2	59.03	14.70
Treatment 3	57.77	12.37
Treatment 4	69.04	16.47
Treatment 5	62.73	16.40
Treatment 6	65.00	16.70
Treatment 7	47.67	10.40
Treatment 8	39.10	9.40
Treatment 9	50.40	12.10
S. Ed. (\pm)	3.83	1.06
C.D. _{P=0.05}	8.04	2.24

Table 8. Days taken to half bloom and days taken to full bloom

Treatment	Days taken to half bloom	Days taken to full bloom
Treatment 1	21.47	30.81
Treatment 2	17.17	25.57
Treatment 3	15.73	24.50
Treatment 4	20.67	29.93
Treatment 5	18.43	27.00
Treatment 6	19.80	28.14
Treatment 7	13.73	21.03
Treatment 8	12.33	19.60
Treatment 9	14.00	22.37
S. Ed. (\pm)	1.07	1.87
C.D. _{P=0.05}	2.24	3.97

Table 9. Bud length (cm) and bud diameter (cm)

Treatment	Bud length (cm)	Bud diameter (cm)
Treatment 1	1.60	0.90
Treatment 2	2.57	1.30
Treatment 3	2.80	1.37
Treatment 4	1.63	1.17
Treatment 5	2.50	1.27
Treatment 6	2.10	1.20
Treatment 7	3.53	1.63
Treatment 8	3.47	1.77
Treatment 9	3.30	1.57
S. Ed. (\pm)	0.19	0.07
C.D. _{P=0.05}	0.40	0.16

Likewise, the bud diameter and highest length of stalk value (1.77 and 27.27) was exhibited by treatment T₈ (Table 10). Superior bud diameter might be due to better growing conditions and good physico-chemical properties of media containing cocopeat with loam soil. These finding are in agreement with the results obtained by [8] in *Celosia cristata* and [29] in gerbera.

The increase in the stalk length of roses grown in cocopeat may be due to a decrease in phenolic

compounds or organic acids. Similar finding was reported by Yonis et al. (2015) in rose. The highest value of stalk diameter of 0.80 cm (Table 10) was exhibited by treatment T₇. This increase in flower stalk diameter may be attributed to overall good growth of plants in these cocopeat and vermicompost amended medium due to its favourable physical and chemical properties. The results are in line with the findings of [30] and [31] in gerbera.

Table 10. Stalk length (cm) and stalk diameter (cm)

Treatment	Stalk length (cm)	Stalk diameter (cm)
Treatment 1	9.73	0.37
Treatment 2	19.90	0.67
Treatment 3	19.17	0.70
Treatment 4	15.93	0.50
Treatment 5	18.53	0.63
Treatment 6	18.17	0.60
Treatment 7	23.50	0.80
Treatment 8	27.27	0.77
Treatment 9	22.67	0.73
S. Ed. (\pm)	1.38	0.05
C.D. _{P=0.05}	2.91	0.10

Table 11. Flower diameter (cm) and number of flowers

Treatment	Flower diameter (cm)	No. of flowers
Treatment 1	5.57	4.74
Treatment 2	7.57	13.07
Treatment 3	7.57	14.02
Treatment 4	6.40	10.41
Treatment 5	7.43	12.47
Treatment 6	7.27	11.33
Treatment 7	7.70	17.74
Treatment 8	8.10	20.04
Treatment 9	7.63	16.48
S. Ed. (\pm)	0.20	1.49
C.D. _{P=0.05}	0.42	3.13

The largest size of flower (8.10 cm) and highest number of flower (20.04) was exhibited by treatment T₈. (Table 11). It is possible that the improved size of flowers grown in cocopeat-amended soil is a result of increased potassium availability in the soil mixture. The larger flower size may be a result of all growth factors working together, including an increase in leaf area. This could be caused by heightened production and accumulation of photosynthates in the leaves, which are then transported to the flowers. The above results are in line with the findings of [18] and [32] in rose.

The higher number of flowers per plant and nitrogen content were recorded in potting media containing cocopeat as cocopeat adds organic matter to the medium and contains more nitrogen as a source of nutrition for the development of more numbers of flower [33].

The data revealed that the highest number (41.00) of petals per flower and the maximum self-life (19.00 days) was exhibited by treatment T₈ (Table 12). The use of cocopeat in the potting media has been found to contribute to an increase in potassium content and positive physico-chemical properties such as high

Table 12. Number of petals per flower and self-life

Treatment	No. of petals per flower	Self-life (days)
Treatment 1	20.74	8.07
Treatment 2	36.37	12.72
Treatment 3	36.38	13.71
Treatment 4	30.43	9.67
Treatment 5	35.07	12.37
Treatment 6	32.67	11.67
Treatment 7	40.75	18.33
Treatment 8	41.00	19.00
Treatment 9	36.10	16.77
S. Ed. (\pm)	1.85	1.06
C.D. _{P=0.05}	3.88	2.22

porosity, good water retention, and improved moisture retention. These factors have been shown to improve the length and width of rose petals, increase the number of petals per flower, and enhance the diameter of the flower. These results are in conformity with [32] in rose and [34] in Carnation. Better flowering duration in cocopeat amended media might be due to higher availability of potassium in media mixture. These results are supported by [19] in rose.

4. CONCLUSION

Rose positively responded to the growing media and the highest performance was observed for the treatment T₈ {Loam soil + Sand + Vermicompost (10% substituted with Cocopeat) @ 1:1:1 (w/w)} which was at par with T₆ {Clay loam + Sand + Vermicompost @ 1:1:1 (w/w)}, as regards to plant height, number of leaves and plant spread. Stem girth, number of nodes, internodal length, leaf area and leaf area index, days taken to bud visibility, bud sprouting, half bloom and full bloom, length of stalk and diameter of bud, number of flowers, number of petals per flower and flowering duration (self-life) and diameter of flower were significantly enhanced by the treatment T₈.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Gauchan DP, Pokhrel AR, Pratap M, Lama P. Current status of cut flower business in Nepal. Kathmandu Univ. J. Sci. Engg. Tech.. 2009;5(1):87-98.
2. Kiran M, Baloch J, Waseem K, Jilani MS. Effect of different growing media on the growth and development of Dahlia (*Dahlia pinnata*) under the agro-climatic condition of Dera Ismail Khan. Pakistan J. Biol. Sci.. 2007;10(22):4140-4143.

3. Tariq U, Rehman SU, Khan MA, Younis A, Yaseen M, Ahsan M. Agricultural and municipal waste as potting media components for the growth and flowering of *Dahlia hortensis* 'Figaro'. Turkish J. Bot.. 2012;36:378-385.
4. Mahmood F, Khan I, Ashraf U, Shahzad T, Hussain S, Shahid M, et al. Effects of organic and inorganic manures on maize and their residual impact on soil physico-chemical properties. J. Soil Sci. Plant Nutrition. 2017;17(1):22-32.
5. Abad M, Noguera P, Puchades R, Maquieira A, Noguera V. Physico-chemical and chemical properties of some coconut dusts for use as a peat substitute for containerized ornamental plants. Biores. Technol. 2002;82:241-245.
6. Manono B. Agro-ecological role of earthworms (*Oligochaetes*) in sustainable agriculture and nutrient use efficiency: A review. Journal of Agriculture and Ecology Research International. 2016;8(1): 1-18.
7. Walie M, Tegegne F, Mekuriaw Y, Tsunekawa A, Kobayashi N, Ichinohe T et al. Biomass yield, quality, and soil nutrients of pasture influenced by farmyard manure and enrichment planting. Rangeland Ecol. Mangt. 2023;88:174-181.
8. Awang Y, Shaharom AS, Mohamad RB, Selamat A. Chemical and physical characteristics of cocopeat-based media mixtures and their effects on the growth and development of *Celosia cristata*. American J. Agril. Biol. Sci.. 2009;4(1):63-71.
9. Kaushal S, Kumari P. Growing media in floriculture crops. J. Pharmacognosy Phytochem. 2020;9(2):1056-1061.
10. Srikanth K, Srinivasamurthy CA, Siddaramappa R, Parama VR. Direct and residual effect of enriched composts, FYM, vermicompost and fertilizers on properties of an Alfisol. J. Indian Soc. Soil Sci.. 2000;48(3):496-499.
11. Tao G, Jian S, Cai M, Xie G. Phosphate-solubilizing and-mineralizing abilities of bacteria isolated from soils. Pedosphere. 2008;18(4):515-523.
12. Domínguez J. State of the art and new perspectives on vermicomposting research. CRC Press LLC. 2004; 401-424.

13. Lazcano C, Domínguez J. The use of vermicompost in sustainable agriculture: Impact on plant growth and soil fertility. *Soil Nutrients* 2011;10(1-23):187.
14. Patel VS, Malam VR, Nurbhanej KH, Vihol AN, Chavada JR. Effect of organic manures and biofertilizers on growth, flowering and flower yield of Rose (*Rosa hybrida* L.) cv. Gladiator. *Int. J. Chemical Stud.* 2017;5(5):1924-1927.
15. Sendhilnathan R, Madhubala V, Rajkumar M, Sureshkumar R. Effect of organic manures and micronutrients on growth and flowering attributes of rose cv. Andhra Red (*Rosa centifolia*). *Plant Archives*. 2019;19(2):3633-3637.
16. Kumar P, Sheoran S, Beniwal BS. Flowering and yield parameters of rose as influenced by different organic manures and their levels. *J. Agri. Res. Technol.* 2023;48(1):39.
17. Hazarika A, Dhaduk, BK, Yadav MK. Vegetative growth and flowering of greenhouse Dutch rose cv. Naranga influenced by different potting media. *Hort. J.* 2010; 23(2):85-87.
18. Yonis A, Raiz A, Javaid F, Ahasan M, Aslam S. Influence of various growing substrate on growth and flowering of potted miniature Rose cv. Baby Boomer. *Curr. Sci. Perspect.* 2015;1(1):16-21.
19. Ahmed I, Aslam K, Qasim M, Zafar S, Ahmad Z. Substrates effects on growth, yield and quality of Rose (*Rosa hybrid* L.). *Pakistan J. Bot.* 2012; 44(1):177-185.
20. Anozie EL, Egwunatum AE, Ezenwenyi JU, Okonkwo CI. Effects of different potting media on the germination and early growth of *Newbouldia laevis* (P. Beauv.) Seem. *Asian J. Res. Agri. Forestry.* 2022;8(4):220-234.
21. Olubode O, Adekola S, Idowu S. Evaluation of flowering pattern, yield and yield determinants of hybrid tea rose in response to seasonal variations and applied organic manure rates. *American J. Plant Sci.* 2015;6:464-482.
22. Campos ML, Van MU, Blok C. Comparison of physical properties of vermicompost from paper mill sludge and green compost as substitutes for peat based potting media. *Int. Symp. Growing Media.* 2009;819:227-234.
23. Abirami K, Rema J, Mathew PA, Srinivasan V, Hamza S. Effect of different propagation media on seed germination, seedling growth and vigour of nutmeg (*Myristica fragrans* Houtt.). *J. Med. Plant Res.* 2010;4:2054-2058.
24. Dubey RK, Simrat S, Kukal SS, Kalsi HS. Evaluation of different organic growing media for growth and flowering of petunia. *Comm. Soil Sci. Plant Anal.* 2013;44(12):1777-1785.
25. Fascella G, Zizzo GV, Agnello S. Evaluating the productivity of red rose cultivars in soilless culture. In: IV Int. Symp. Rose Res. Cult. 2005;751: 99-105.
26. Golchin A, Nadi M, Mozaffari V. The effects of vermicomposts produced from various organic solid wastes on growth of pistachio seedlings. In: IV Int. Symp. Pistachios and Almonds. 2005;726:301-306.
27. Gupta YC, Quec Dein LE, Dhiman SR, Jain R. Standardization of growing media under protected environment for gerbera in mid hill of Himachal Pradesh J. Orn.. *Hort.* 2004;7(1):99-102.
28. Treder J. The effects of cocopeat and fertilization on the growth and flowering of oriental lily 'star gazer'. *J. Fruit Orn. Plant Res.* 2008;16:361-370.
29. Kale RD, Jagtap KB, Badgujar CD. Effect of different containers and growing media on yield and quality parameters of gerbera (*Gerbera jamesonii* Bolus ex. Hooker F.) under protected cultivation. *J. Orn. Hort.* 2009;12(4):261-264.
30. Sindhu SS, DB G, Singh MC, Dhiman MR. Effect of medium amendments on growth and flowering in gerbera. *Indian J. Hort.* 2010;67(4):391-394.
31. Khalaj MA, Amiri M, Sindhu SS. Study on the effect of different growing media on the growth and yield of gerbera (*Gerbera jamesonii* L.). *J. Orn. Plant.* 2015;1(3):185-189.
32. Barman D, Rajni K, Upadhyaya RC, Singh DK. Effect of horticultural practices for sustainable production of rose in partially modified greenhouse. *Indian J. Hort.* 2006;63(4):415-418.
33. Awang Y, Shaharom AS, Mohamad RB, Selamat A. Growth dynamics of *Celosia cristata* grown in cocopeat, burnt rice hull and kenaf core fiber mixtures. *American J. Agri. Biol. Sci.* 2010;5(1): 70-76.

34. Bhatia S, Gupta YC, Dhiman SR. Effect of growing media and fertilizers on growth and flowering carnation under protected condition. J. Orn. Hort.. 2004;7(2):174-178.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/123884>