



Xanthine Oxidase Inhibitory Activity and Antioxidant Property of Various Extracts from the Medicinal Plant *Paronychia argentea* L.

Moufida Adjadj^{1*}, Meriem Djarmouni¹ and Abderrahmane Baghiani¹

¹Laboratory of Applied Biochemistry, Department of Biochemistry, Faculty of Natural and Life Sciences, University Ferhat Abbas, Setif 1, Algeria.

Authors' contributions

This work was carried out in collaboration between all authors. Author MA designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors MD and MA managed the analyses of the study. Author AB managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJB2T/2017/39081

Editor(s):

(1) Armando Zepeda Bastida, Professor, Veterinary Biotechnology, Agricultural Science Institute of Autonomous University, Hidalgo State, Mexico.

Reviewers:

- (1) Muhammad Ali, Kano University of Science and Technology, Nigeria.
(2) Rojita Mishra, Polasara Science College, India.
(3) Daohong Chen, New Drug Discovery Center, Yiling Pharmaceutical, China.
Complete Peer review History: <http://prh.sdiarticle3.com/review-history/22897>

Original Research Article

Received 28th October 2017
Accepted 17th January 2018
Published 27th January 2018

ABSTRACT

Aim: *Paronychia argentea* L. is used in traditional medicine to treat various diseases in Algeria especially kidney stones. This study focused on the antioxidant and xanthine oxidase inhibitory activities of these plant extracts.

Methods: The different extracts were prepared using solvents of increasing polarity, and their total polyphenols and flavonoids contents were determined using spectrophotometric methods. The antioxidant activities of *Paronychia argentea* extracts (PAE) were assessed by their inhibitory effect on xanthine oxidase (XO) and their scavenger ability on superoxide radical ($O_2^{\cdot-}$) generated by this enzyme.

Results: The phytochemical investigation of the plant extracts showed that the ethyl acetate extract (EaE) has the highest concentration of total polyphenols and flavonoids, followed by

*Corresponding author: E-mail: moufidatoxicology@gmail.com;

chloroform (ChE) and crud (CE) extracts. Results of the antioxidant activity revealed that all PAE were effective in both XO inhibiting and superoxide radical scavenging, with the following order for the two assays: ChE > EaE > CE.

Conclusion: This study provided evidence that PAE had interesting antioxidant potential which was confirmed using two enzymatic methods. Therefore, this plant could be used to treat gout and lot of diseases, where inhibition of XO and scavenging of superoxide radical are necessary.

Keywords: Flavonoids; oxidative stress; *Paronychia argentea* L.; superoxide anion radical; traditional medicine; xanthine oxidase.

1. INTRODUCTION

Oxidative stress occurs when the balance between the production of reactive oxygen species (ROS) and the quantity of antioxidants is interrupted. This can lead to damage of biomolecules and ultimately leads to cell death causing physiological disorders such as cancer, diabetes, asthma, premature aging, cardiovascular, neurodegenerative and inflammatory diseases [1]. ROS can be synthesized by a variety of enzymes, including xanthine oxidase (XO) [2], which catalyzes the metabolism of purines, converting hypoxanthine to xanthine and xanthine to uric acid with reduction of molecular oxygen to hydrogen peroxide and superoxide anion radical ($O_2^{\cdot-}$) [3-4]. XO causes gout; an inflammatory disease due to elevated levels of uric acid crystals in the serum [5-6]. In addition, oxidizing products of XO involved in the development of cardiovascular and metabolic diseases, leading to atherosclerosis [7]. Thus, the inhibitors of XO may be used for the treatment of gout or other XO induced diseases [8]. Several works have shown that phenolic compounds and flavonoids function as antioxidants including: inhibition of oxidative enzymes such as XO [9], and scavenging of free radicals [10].

Plants are a rich source of antioxidants such as phenolic compounds, anthocyanins and flavonoids. Herbal antioxidants can reduce the development of several human diseases related to oxidative stress [11]. *Paronychia argentea* L. Caryophyllaceae is one of the most used plants in folk medicine in Algeria. It is popularly known: Arabic tea (Fettatet lahdjer or Bissat elmoulouk). The aerial parts are used as diuretic. They are also used to treat renal diseases as antiurolithiasis [12]. This plant was reported to contain the flavonoids isorhamnetin, quercetin, and luteolin [13]. It was reported to have digestive [14], hypoglycemic [15], and antimicrobial activities [16]. Furthermore, Dafni et al. [17] reported the use of leaf decoction of this

plant as diuretic and to treat kidney stones, diabetes and heart ailments. It was also used as a blood purifier [18]. In Portugal, *Paronychia argentea* is used as analgesic, for stomach ulcer, anorexia, and flatulence [19]. This paper aimed at studying the XO inhibitory activity and radical scavenging property of *Paronychia argentea* extracts by applying *in vitro* enzymatic assays.

2. MATERIALS AND METHODS

2.1 Plant Material and Chemicals

Aerial parts of *Paronychia argentea* were collected, at the flowering stage (April - May 2011), from Ouled Rahmoune, Constantine, Algeria, and air-dried at room temperature. The plant was identified and authenticated by Prof. Oudjehih Bachir, a botanist at University El Hadj Lakhdar, Batna, Algeria. All chemicals were purchased from Sigma (Germany), Pfizer Health AB (Sweden), Prolabo, Aldrich and Fluka.

2.2 Methods

2.2.1 Purification of bovine milk xanthine oxidoreductase (XOR)

XOR was purified from bovine milk in the presence of 10 mM of dithiothreitol, by ammonium sulfate fractionation, followed by affinity chromatography on heparin agarose, according to Baghiani et al. [20]. The concentration of XOR was determined *via* UV-visible spectroscopy using an absorption coefficient of $36000 \text{ M}^{-1}\text{cm}^{-1}$ at 450 nm. Estimation of the purity of the enzyme was based on protein/flavin ratio ($\text{PFR} = A_{280}/A_{450}$) and 10% sodium dodecyl sulfate polyacrylamide gel electrophoresis (10% SDS-PAGE), whereas XOR activity was assessed spectrophotometrically by measuring the production of uric acid obtained from xanthine (100 μM , final concentration) at 295 nm using an absorption coefficient of $9600 \text{ M}^{-1}\text{cm}^{-1}$. Assays were performed at room temperature in air-

saturated 50 mM phosphate buffer, pH 7.4, supplemented with 0.1 mM EDTA.

2.2.2 Extraction of phenolic compounds

The extraction was carried out using polar and non-polar solvents (methanol, *n*-hexane, chloroform, ethyl acetate) according to a procedure outlined by Baghiani et al. [21]. One hundred grams of dried and powdered plant material was soaked in 1 L of 85 % methanol at 4 °C for 16 h. Then, the residue obtained after filtration was re-extracted with 1 L of 50 % methanol for 4 h. The resulting solutions from the first and the second extractions were concentrated by a rotary evaporator to obtain the crude extract (CE). The CE was then fractionated with *n*-hexane, chloroform, and ethyl acetate to obtain these solvent extracts (HxE, ChE and EaE, respectively), and the remaining aqueous extract was labeled AqE. Solvents were removed by evaporation under reduced pressure, and the obtained dried extracts were stored until use.

2.2.3 Total polyphenols contents determination

Total polyphenols contents of *Paronychia argentea* extracts (PAE) were determined with the Folin-Ciocalteu reagent using gallic acid as a standard according to Li et al. [22]. In brief, 0.1 mL of PAE was mixed with 2.5 mL of distilled water and 0.5 mL of Folin-Ciocalteu stock reagent. After 5 min, 1.0 mL of 20% aqueous Na₂CO₃ solution was added to the mixture. The mixture was then incubated at room temperature for 1 h and its absorbance was measured at 760 nm. The amount of total polyphenols in different extracts was determined from standard curve of gallic acid, and results are expressed in milligrams of gallic acid equivalents per gram of dried PAE (mgGA-Eq/gE).

2.2.4 Total flavonoid contents determination

Total flavonoid contents in PAE were determined according to the method of Bahorun et al. [23]. Briefly, 1 mL of each sample was mixed with 1 mL of aluminium chloride (AlCl₃) solution (2%, in methanol). After incubation for 10 min, absorbance of mixture was measured at 430 nm versus a prepared methanol blank. Quercetin and rutin were used as standards. Results are expressed as milligrams of quercetin and rutin equivalents per gram of dried PAE (mgQ-Eq/gE and mgR-Eq/gE, respectively).

2.2.5 Effect of *Paronychia argentea* extracts on superoxide anion radicals generated by xanthine oxidase

Superoxide radicals generated by XO are able to reduce cytochrome c. Free radical scavenging activity of PAE was evaluated according to the method outlined by Boumerfeg et al. [24]. In this method, a mixture containing xanthine (100 µM), horse heart cytochrome c (25 µM), in air-saturated sodium phosphate buffer (50 mM, pH 7.4), supplemented with 0.1 mM EDTA and various concentrations of PAE was obtained. The reaction started upon addition of XO, and the extent of cytochrome c reduction was determined at 550 nm against enzyme free mixture using an absorption coefficient of 21.100 M⁻¹ cm⁻¹. Bovine erythrocytes superoxide dismutase (SOD), (330 U/mL final concentration) was employed to assess the sensitivity of the reaction. The inhibitory activity of cytochrome c reduction by PAE was expressed as percent inhibition (I%) calculated as follows:

$$I (\%) = [(A_{\text{control}} - A_{\text{sample}})/A_{\text{control}}] \times 100$$

Where A_{control} is the absorbance of the control reaction (containing all reagents except the test solutions), and A_{sample} is the absorbance of the test compound. Extract concentration providing 50% inhibition (IC₅₀) was calculated from the plot of inhibition percentage against extract concentration.

2.2.6 Effect of *Paronychia argentea* extracts on xanthine oxidase activity

The effect of PAE on XO activity was evaluated spectrophotometrically at 295 nm by measuring the formation of uric acid from xanthine at room temperature following the procedure published by Boumerfeg et al. [25]. Mixtures containing a final concentration of 100 µM of xanthine, and various amounts of PAE were made. Addition of 1176 nmol of urate /min/mg XOR protein to each mixture initiated the reaction; enzyme activity of the control was set as 100 % activity. Allopurinol, a clinical drug for XO inhibition, was used as standard inhibitor. The percent inhibition was calculated by using the following formula:

$$I (\%) = [(A_{\text{control}} - A_{\text{sample}})/A_{\text{control}}] \times 100$$

Where A_{control} is the absorbance of the control reaction (containing all reagents except the test solutions), and A_{sample} is the absorbance of the test compound. Extract concentration providing 50% inhibition (IC₅₀) was calculated from the plot

of inhibition percentage against extract concentration.

2.2.7 Statistical analysis

All determinations were conducted in triplicate. Results in both XO inhibition and superoxide radical scavenging assays were calculated as mean \pm standard deviation (SD). While yield percentages and amounts of total polyphenols and flavonoids are expressed as mean \pm standard error of the mean (SEM). Statistical analysis was performed using Student's t-test for significance and analysis of variance (ANOVA) followed by Dunnet's test for the multiple effects comparison of the different extracts. The p values less than 0.05 were considered statistically significant.

3. RESULTS AND DISCUSSION

3.1 Purification of Bovine Milk Xanthine Oxidoreductase

Xanthine oxidoreductase purified from fresh bovine milk yielded 23.21 mg of XOR protein per liter, which is comparable to the amounts reported by Baghiani et al. [20]. The obtained enzyme was largely (more than 90%) under the oxidase form. The freshly purified bovine milk XOR exhibited an ultraviolet/visible spectrum with three major peaks at 280, 330, and 450 nm (Fig. 1A), with A280/A450 (protein to flavin ratio, PFR) of 5.15 which indicates a high degree of purity. These results are in agreement with those reported in the literature [21,26]. The purified enzyme showed one major band of an approximate molecular weight of 150 KDa when run on SDS/PAGE (Fig. 1B). The oxidase form of the purified enzyme showed an activity of **1988.55** nmol of urate/min/mg protein. These results are similar to those reported by Baghiani et al. [9,20,21] and Atmani et al. [27].

3.2 Extraction of Phenolic Compounds and Determination of Total Polyphenol and Flavonoid Contents

Table 1 lists the yields, percentages, and amounts of total polyphenols and flavonoids. Results show that aqueous extract (AqE) has the highest yield, followed by the crude extract (CE), whereas the other extracts displayed lower yields. On the other hand, results also show that ethyl acetate (EaE) extract contained the highest total polyphenols and flavonoids contents.

3.3 Effect of *Paronychia argentea* Extracts on the Generation of Superoxide Anion Radicals by XO System

The ability of PAE to scavenge superoxide anion ($O_2^{\cdot-}$) radicals was determined by following reduction of cytochrome c (cyt c) at 550 nm. Results revealed that SOD (330 U/mL) has totally inhibited the reduction of cyt c (1% = 100%), and show that all PAE significantly inhibit the cyt c reduction in a concentration-dependent manner ($p < 0.0001$) (Fig. 2A). Comparison between these extracts reveal that ChE has the highest ability to scavenge superoxide anion radicals followed by EaE and CE, with IC_{50} (mg/mL) values of 0.092 ± 0.00014 , 0.098 ± 0.0002 , and 0.277 ± 0.0015 , respectively (Fig. 2B).

Superoxide anion formed during XO reaction can be quantified using cytochrome c [28]. It causes damage to biomolecules by forming H_2O_2 , OH^{\cdot} that can initiate lipid peroxidation [29]. Since the presence of proteins, sugars, and polyphenols in the extracts may affect their antioxidant activity [30], it is possible that the antioxidative properties of PAE are caused by the presence of polyphenols and flavonoids. Moreover, the antioxidant properties of polyphenols are directly linked to their structure [31]. Phenols are made up of one or more aromatic rings with one or more hydroxyl groups. Therefore, these phenols are potentially able to quench free radicals by forming resonance-stabilized phenoxyl radicals [32]. On the other hand, structure-activity relationships of flavonoids in scavenging superoxide anion were studied by Cos et al. [33] and Dugas et al. [34]. They discovered that the presence of hydroxyl groups and carbon-carbon double bonds in certain positions enhance flavonoids scavenging activity.

3.4 Effect of *Paronychia argentea* Extracts on XO Activity

Since an inhibitory effect on the enzyme itself would lead to a decrease in cyt c reduction [35], the effect of PAE on the inhibition of XO was studied. Results demonstrated that all extracts exerted a very significant concentration-dependent inhibition of XO activity ($p < 0.0001$) (Fig. 3A). The IC_{50} values revealed that the highest XO inhibitory effect was shown with ChE followed by EaE and CE (Fig. 3B).




Fig. 1. UV/Visible absorbance spectrum of Bovine XOR (A), 10% SDS-PAGE of XOR preparation (B)

Lan1: purified bovine milk XOR; Lan2: crud bovine milk XOR and Lan3: molecular weight markers: Myosin 205 000; β -galactosidase 116 000; Phosphorylase 97 400; Serum albumin 66 000; Ovalbumin 45 000; Carbonic anhydrase 29 000

Table 1. The yields and the amounts of total polyphenols and flavonoids compounds in different extracts of *Paronychia argentea*

Extracts	% yield (w/w)	Total polyphenols (mg GA-Eq / g E)	Total flavonoids	
			(mg Q-Eq / g E)	(mg R-Eq / g E)
CE	10.971±0.637	217.463±0.87	13.349±0.562	26.765±0.837
HxE	0.314±0.097	nm	nm	nm
ChE	0.212±0.015	211.444±0.778	38.621±1.303	87.717±1.811
EaE	0.786±0.175	525.796±0.796	194.193±8.622	382.176±4.74
AqE	11.271±1.831	nm	nm	nm

nm: not mentioned. Values are expressed as mean \pm SEM (n =3)




Fig. 2. The inhibitory effect of *P. argentea* extracts on the reduction of cyt c by $O_2^{\cdot -}$ anion radical produced by XO (A), concentrations of *P. argentea* extracts which inhibit 50% reduction of cyt c by superoxide produced by XO (scavenger effect, IC_{50}) (B)

CE: methanol extract; ChE: chloroform extract; EaE: ethyl acetate extract.

Values are means \pm SD (n = 3). *** p < 0.0001, comparing with the control in absence of extracts

The presence of polyphenols and flavonoids in the extract can enhance XO inhibition [36-37]. Thus, the higher XO inhibitory effect of ChE and EaE can be due to their richness in flavonoids. XO causes hyperuricemia, gout and

cardiovascular disease, and their inhibitors are considered effective drugs to control these uric acid-related problems [38-39]. Allopurinol is the most used XO inhibitor drug, but it has several adverse effects including: gastrointestinal

irritation, hypersensitivity syndromes, fever, hepatitis, eosinophilia and worsening renal function [40]. Thus, due to the limitations of currently available XO inhibitory drugs, the development of new ones with increased therapeutic activity and less side effects is an active field of research [38]. XO contains a molybdopterin (Mo) domain which is the active site of the enzyme [41-42]. The inhibitors of the enzyme; allopurinol and 3,4-dihydroxy-5-nitrobenzaldehyde (DHNB) inhibit the XO activity via interaction with its Mo center [38,43]. Therefore, it is possible that XO inhibitory activity of PAE is due to the presence of various compounds which are fixed to this active site.

On the other hand, the inhibition of cyt c reduction is due to the XO inhibitory effect and/or the scavenger effect on $O_2^{\cdot -}$ produced by this enzyme [44]. Fig. 4 shows a comparison between IC_{50} values of cyt c reduction and XO inhibition effects of PAE. The ChE and EaE IC_{50} values of cyt c reduction are higher than those of XO inhibition. Thus, according to Cos et al. [33] the inhibition of cyt c reduction by these extracts is due to their XO inhibitory effect and to a weak $O_2^{\cdot -}$ radical scavenger effect. CE has an IC_{50} of cyt c reduction almost identical to that of XO inhibition, therefore according to Cos et al. [33], it is considered as an antioxidant because of its ability to inhibit XO.




Fig. 3. The inhibitory effect of *P. argentea* extracts on the activity of XO (A), concentrations of *P. argentea* extracts and allopurinol that inhibit 50% of XO activity (IC_{50}) (B)

CE: methanol extract; ChE: chloroform extract; EaE: ethyl acetate extract.

Values are means \pm SD ($n = 3$). *** $p < 0.0001$, comparing with the control in absence of extracts




Fig. 4. Comparison of IC_{50} values of cyt c reduction and XO inhibition effects of *Paronychia argentea* extracts

CE: methanol extract; ChE: chloroform extract; EaE: ethyl acetate extract.

Values are expressed as mean \pm SD ($n = 3$).

*** $p < 0.0001$, comparing with the control in absence of extracts

4. CONCLUSION

Paronychia argentea extracts contain significant amounts of flavonoids and polyphenols and have a strong antioxidant activity. Consequently, these plant extracts can be used as a source of bioactive compounds which may function as useful natural antioxidants and therapeutic agents for hyperuricemia, gout and other related diseases, where inhibition of XO and scavenging of superoxide radicals are necessary. However, further investigations to isolate and identify the antioxidant compounds present in the plant extracts, and further studies on definitive mechanisms of their therapeutic activities *in-vivo* are needed.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Subedi L, Timalansa S, Duwadi P, Thapa R, Paudel A, Parajuli K. Antioxidant activity and phenol and flavonoid contents of eight medicinal plants from Western Nepal. *Journal of Traditional Chinese Medicine*. 2014;34(5):584-90. DOI: 10.1016/S0254-6272(15)30067-4.
- Kundu TK, Velayutham M, Zweier JL. Aldehyde oxidase functions as a superoxide generating NADH oxidase: an important redox regulated pathway of cellular oxygen radical formation. *Biochemistry*. 2012;51(13):2930-9. DOI: 10.1021/bi3000879.
- Kardeh S, Ashkani-Esfahani S, Alizadeh AM. Paradoxical action of reactive oxygen species in creation and therapy of cancer. *European Journal of Pharmacology*. 2014;735(1):150-68. DOI: 10.1016/j.ejphar.2014.04.023.
- Kelley EE. A new paradigm for XOR-catalyzed reactive species generation in the endothelium. *Pharmacological Reports*. 2015;67(4):669-74. DOI: 10.1016/j.pharep.2015.05.004.
- Masuda T, Shingai Y, Takahashi C, Inai M, Miura Y, Honda S, Masuda A. Identification of a potent xanthine oxidase inhibitor from oxidation of caffeic acid. *Free Radical Biology and Medicine*. 2014;69:300-7. DOI:10.1016/j.freeradbiomed.2014.01.016.
- Masuda T, Nojima S, Miura Y, Honda S, Masuda A. An oxidative coupling product of luteolin with cysteine ester and its enhanced inhibitory activity for xanthine oxidase. *Bioorganic & Medicinal Chemistry Letters*. 2015;25(16):3117-9. DOI: 10.1016/j.bmcl.2015.06.016.
- Battelli MG, Bolognesi A, Polito L. Pathophysiology of circulating xanthine oxidoreductase: New emerging roles for a multi-tasking enzyme. *Biochimica et Biophysica Acta*. 2014;1842(9):1502-1517. DOI: 10.1016/j.bbadis.2014.05.022.
- Sweeney AP, Wyllie SG, Shalliker RA, Markhan JL. Xanthine oxidase inhibitory activity of selected Australian native plants. *Journal of Ethnopharmacology*. 2001;75(2-3):273-7. DOI: 10.1016/S0378-8741(01)00176-3.
- Baghiani A, Ameni D, Boumerfeg S, Adjadj M, Djarmouni M, Charef N, Khennouf S, Arrar L. Studies of Antioxidants and Xanthine Oxidase Inhibitory Potentials of Root and Aerial Parts of Medicinal Plant *Capparis Spinosa* L. *American Journal of Medicine and Medical Sciences*. 2012;2(1):25-32. DOI: 10.5923/j.ajmms.20120201.06.
- Custódia L, Patarra J, Alberício F, Neng N, Nogueira JMF, Romano A. Phenolic composition, antioxidant potential and *in vitro* inhibitory activity of leaves and acorns of *Quercus suber* on key enzymes relevant for hyperglycemia and Alzheimer's disease. *Industrial Crops and Products*. 2015;64:45-51. DOI: 10.1016/j.indcrop.2014.11.001.
- Tauchen J, Doskocil I, Caffi C, Lulekal E, Marsik P, Havlik J, Van Damme P, Kokoska L. *In vitro* antioxidant and anti-proliferative activity of Ethiopian medicinal plant extracts. *Industrial Crops and Products*. 2015;74:671-9. DOI: 10.1016/j.indcrop.2015.05.068
- Adjadj M, Baghiani A, Boumerfeg S, Nouredine C, Khennouf S, Arrar L, Mubarak MS. Protective Effect of *Paronychia argentea* L. on Acetic Acid Induced Ulcerative Colitis in Mice by Regulating Antioxidant Parameters and Inflammatory Markers. *Der. Pharma. Chemica*. 2016;8(4):207-218.
- Rizk AM. *The Phytochemistry of the Flora of Qatar*. Doha, State of Qatar: The Scientific and Applied Research Center, University of Qatar. 1986;22.
- Novais HM, Santos I, Mendes S, Pinto-Gomes C. Studies on pharmaceutical ethnobotany in Arrabida Natural Park.

- Journal of Ethnopharmacology. 2004;93(2-3):183-95.
DOI: 10.1016/j.jep.2004.02.015
15. Carmona MD, Llorach R, Obon C, Rivera D. 'Zahraa', a Unani multicomponent herbal tea widely consumed in Syria: components of drug mixtures and alleged medicinal properties. Journal of Ethnopharmacology. 2005;102(3):344-50. DOI: 10.1016/j.jep.2005.06.030.
 16. Al-Bakri AG, Affifi FU. Evaluation of antimicrobial activity of selected plant extracts by rapid XTT colorimetry and bacterial enumeration. Journal of Microbiological Methods. 2007;68(1):19-25. DOI: 10.1016/j.mimet.2006.05.013.
 17. Dafni A, Yanif Z, Palevitch D. Ethnobotanical survey of medicinal plants in northern Israel. Journal of Ethnopharmacology. 1984;10(3):295-310. DOI: 10.1016/0378-8741(84)90017-5.
 18. González-Tejero MR, Molero Mesa J, Casares M, Martínez MJ. New contributions to the ethnopharmacology of Spain. Journal of Ethnopharmacology. 1995;45(3):157-65. DOI: 10.1016/0378-8741(94)01208-H.
 19. Ferreira A, Proenca C, Serralheiro MLM, Araújo MEM. The in vitro screening for acetylcholinesterase inhibition and antioxidant activity of medicinal plants from Portugal. Journal of Ethnopharmacology. 2006;108(1-3):31-7. DOI: 10.1016/j.jep.2006.04.010.
 20. Baghiani A, Harrison R, Benboubetra M. Purification and partial characterization of camel milk xanthine oxidoreductase. Archives of Physiology and Biochemistry. 2003;111(5):407-14. DOI: 10.3109/13813450312331342265.
 21. Baghiani A, Boumerfeg S, Adjadj M, Ameni D, Djarmouni M, Khelifi Touhami F, Charef N, Khennouf S, Arrar L. Antioxidants, Free Radicals Scavenging and Xanthine Oxidase Inhibitory Potentials of *Ajuga iva* L. Extracts. Free Radicals and Antioxidants. 2011;1(4):21-30. DOI: 10.5530/ax.2011.4.5.
 22. Li HB, Cheng KW, Wong CC, Fan KW, Chen F, Jiang Y. Evaluation of antioxidant capacity and total phenolic content of different fraction of selected microalgae. Food Chemistry. 2007;102(3):771-6. DOI: 10.1016/j.foodchem.2006.06.022.
 23. Bahorun T, Gressier B, Troitin F, Brunete C, Dine T, Vasseur J, Gazin JC, Pinkas M, Luycky M, Gazin M. Oxygen species scavenging activity of phenolic extract from hawthorn fresh plant organs and pharmaceutical preparation. Arzneimittel Forsh / Drug Res. 1996;46(11):1086-9.
 24. Boumerfeg S, Baghiani A, Djarmouni M, Ameni , Adjadj M, Belkhiri F, Charef N, Khennouf S, Arrar L. Inhibitory Activity on Xanthine Oxidase and Antioxidant Properties of *Teucrium polium* L. Extracts. Chinese Medicine. 2012;3(1):30-41. DOI: 10.4236/cm.2012.31006.
 25. Boumerfeg S, Baghiani A, Messaoudi D, Khennouf S, Arrar L. Antioxidant properties and xanthine oxidase inhibitory effects of *Tamus communis* L root extracts. Phytother. Res. 2009;23(2):283-8. DOI: 10.1002/ptr.2621.
 26. Benboubetra M, Baghiani A, Atmani D, Harrison H. Physicochemical and kinetic properties of purified sheep's milk xanthine oxidoreductase. J. Dairy Sci. 2004;87(6):1580-4. DOI: 10.3168/jds.S0022-0302(04)73311-1.
 27. Atmani D, Benboubetra M, Harrison H. Goat's milk xanthine oxidoreductase is grossly deficient in molybdenum. J. Dairy Res. 2004;71(1):7-13.
 28. Flemmig J, Kuchta K, Arnholda J, Rauwald HW. *Olea europaea* leaf (Ph.Eur.) extract as well as several of its isolated phenolics inhibit the gout-related enzyme xanthine oxidase. Phytomedicine. 2011;18(7):561-6. DOI: 10.1016/j.phymed.2010.10.021.
 29. Giese EC, Gascon J, Anzelmo G, Barbosa AM, Cunha MAA, Dekker RFH. Free radical scavenging properties and antioxidant activities of botryosphaeran and some other β -D-glucans. International Journal of Biological Macromolecules. 2015;72:125-30. DOI: 10.1016/j.ijbiomac.2014.07.046.
 30. Renuka Devi R, Arumughan C. Antiradical efficacy of phytochemical extracts from defatted rice bran. Food and Chemical Toxicology. 2007;45(10):2014-21. DOI: 10.1016/j.fct.2007.04.020.
 31. Balan T, Sani MHM, Ahmad SHM, Suppaiah V, Mohtarrudin N, Jamaludin F, Zakaria ZA. Antioxidant and anti-inflammatory activities contribute to the prophylactic effect of semi-purified fractions obtained from the crude methanol extract of *Muntingia calabura* leaves against gastric ulceration in rats. Journal of Ethnopharmacology. 2014;164:1-15. DOI: 10.1016/j.jep.2014.12.017.

32. Bors W, Michel C. Chemistry of the antioxidant effect of polyphenols. *Annals of the New York Academy of Sciences*. 2002;957:57-69.
DOI: 10.1111/j.1749-6632.2002.tb02905.x.
33. Cos P, Ying L, Calomme M, Hu JP, Cimanga K, Van-Poel B, Pieters L, Vlietinck AJ, Vanden Berghe D. Structure-activity relationship and classification of flavonoids as inhibitors of xanthine oxidase and superoxide scavengers. *J. Nat. Prod.* 1998;61(1):71-6.
DOI: 10.1021/np970237h.
34. Dugas AJ, Castaneda-Acosta J, Bonin GC, Price KL, Fischer NH, Winston GW. Evaluation of the total peroxyl radical-scavenging capacity of flavonoids: structure-activity relationships. *J. Nat. Prod.* 2000;63(3):327-31.
DOI: 10.1021/np990352n.
35. Valentao P, Fernandes E, Carvalho F, Andrade PB, Seabra RM, Bastos ML. Antioxidant activity of *Centaureum erythraea* infusion evidenced by its superoxide radical scavenging and xanthine oxidase inhibitory activity. *J. Agric. Food Chem.* 2001;49(7):3476-9.
DOI: 10.1021/jf001145s.
36. Abdullahi A, Hamzah RU, Jigam AA, Yahya A, Kabiru AY, Muhammad H, Sakpe S, Adefolalu FS, Isah MC, Kolo MZ. Inhibitory activity of xanthine oxidase by fractions *Crateva adansonii*. *Journal of Acute Disease*. 2012;1(2):126-9.
DOI: 10.1016/S2221-6189(13)60029-3.
37. Kilani-Jaziri S, Ghedira Z, Nasr N, Krifa M, Ghedira K, Franca Dijoux MG, Ghedira-Chekir L. Evaluation of in vitro antioxidant and apoptotic activities of *Cyperus rotundus*. *Asian Pacific Journal of Tropical Medicine*. 2014;7(2):105-12.
DOI: 10.1016/S1995-7645(14)60004-3.
38. Lu JM, Yao Q, Chen C. 3,4-Dihydroxy-5-nitrobenzaldehyde (DHNB) is a potent inhibitor of xanthine oxidase: A potential therapeutic agent for treatment of hyperuricemia and gout. *Biochemical Pharmacology*. 2013;86(9):1328-37.
DOI: 10.1016/j.bcp.2013.08.011.
39. Li DQ, Zhao J, Li SP. High-performance liquid chromatography coupled with post-column dual-bioactivity assay for simultaneous screening of xanthine oxidase inhibitors and free radical scavengers from complex mixture. *Journal of Chromatography A*. 2014;1345:50-6.
DOI: 10.1016/j.chroma.2014.03.065.
40. Lin WQ, Xie JX, Wu XM, Yang L, Wang HD. Inhibition of Xanthine Oxidase Activity by *Gnaphalium affine* Extract. *Chin. Med. Sci. J.* 2014;29(4):225-30.
DOI: 10.1016/S1001-9294(14)60075-4.
41. Enroth C, Eger BT, Okamoto K, Nishino T, Pai EF. Crystal structures of bovine milk xanthine dehydrogenase and xanthine oxidase: structure-based mechanism of conversion. *Proc. Natl. Acad. Sci. USA*. 2000;97(20):10723-8.
DOI: 10.1073/pnas.97.20.10723.
42. Metz S, Thiel W. A combined QM/MM study on the reductive half-reaction of xanthine oxidase: substrate orientation and mechanism. *J. Am. Chem. Soc.* 2009;131(41):14885-902.
DOI: 10.1021/ja9045394.
43. Pauff JM, Hille R. Inhibition studies of bovine xanthine oxidase by luteolin, silibinin, quercetin and curcumin. *J. Nat. Prod.* 2009;72(4):725-31.
DOI: 10.1021/np8007123.
44. Valentao P, Fernandes E, Carvalho E, Andrade PB, Seabra RM, Bastos ML. Antioxidant activity of *Hypericum androsaenum* infusion scavenging effect on superoxide radical, hydroxyl radical and hypochlorous acid. *Biological and Pharmaceutical Bulletin*. 2002;25(10):1324-7.

© 2017 Adjadj et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://prh.sdiarticle3.com/review-history/22897>