



Impact of Zinc and Thiourea Application on Growth and Yield of Summer Green Gram (*Vigna radiata* L.)

D. J. Patel ^{a++}, N. I. Patel ^{b#}, R. H. Kotadiya ^{c++},
K. H. Chaudhary ^{a++}, Devilal Birla ^{c++*} and Manoj Dohat ^{c†}

^a Department of Soil Science & Agricultural Chemistry, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar, Gujarat-385506, India.

^b Centre for Natural Resources Management, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar, Gujarat-385506, India.

^c Department of Agronomy, Anand Agricultural University, Anand, Gujarat-388110, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i183366

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103036>

Original Research Article

Received: 10/05/2023

Accepted: 13/07/2023

Published: 26/07/2023

ABSTRACT

Aims: To find out the Impact of zinc and thiourea application on growth and yield of summer green gram (*Vigna radiata* L.).

Study Design: The experiment was laid out in Randomized Block Design (factorial concepts) with three replications.

Place and Duration of Study: A field experiment was conducted during summer season of 2020 at Agronomy Instructional Farm, SDAU, Dantiwada (Gujarat). The soil of experimental plot was loamy sand in texture.

⁺⁺ Research Scholar;

[#] Associate Research Scientist;

[†] Assistant Professor;

*Corresponding author: E-mail: devilalbirla130596@gmail.com;

Methodology: The experiment comprising three levels of zinc (1.5, 3.0 and 4.5 kg ha⁻¹) and five treatments of thiourea application (control (no spray), seed soaking with thiourea @ 500 ppm, foliar spray of thiourea @ 500 ppm at 30 DAS, foliar spray of thiourea @ 500 ppm at 45 DAS and foliar spray of thiourea @ 500 ppm at 30 and 45 DAS).

Results: Almost all the growth and yield attributes as well as yield were significantly influenced due to various treatments. The plant height (47.02 cm and 46.60 cm, respectively), total number of pods plant⁻¹ (15.92 and 19.82, respectively) and number of seeds pod⁻¹ (7.91 and 8.26, respectively) and seed yield and stover yield were also found significantly higher in the application of either 4.5 kg Zn ha⁻¹ or foliar sprays of thiourea @ 500 ppm at 30 and 45 DAS.

Conclusion: On the basis of experimental results, it can be concluded that to obtain higher yield of green gram crop responds positively to the application of zinc and thiourea with enhancement in the agronomical performance and productivity.

Keywords: Green gram; soil application; foliar application; thiourea and zinc.

1. INTRODUCTION

“Pulses are agriculture’s lifeblood since they play a distinct role in every system as the primary crop, catch crop, cover crop, green manure, intercrop and mix crop. One of India’s principal pulse crops is greengram, often known as moong. Along with fibre and iron, it provides a rich source of protein. Green gram has extremely little cholesterol and is low in sodium and saturated fat. Along with being a very good source of dietary fibre, vitamin C, vitamin K, riboflavin, folate, copper, manganese, and protein, it is also a good source of 1008adiate1008, niacin, vitamin B₆, pantothenic acid, iron, magnesium, phosphorus and potassium. Additionally, 100 g of dry green gram seeds contain 104.5% copper, 84% iron, 45% manganese, 45% phosphorus, 52% selenium, 15% calcium, and 24% zinc” [1].

“Such limiting conditions are brought about by the alkaline reaction, low organic matter, high calcium carbonate, low soil moisture and coarse texture of soils. Crops develop slowly and have very low Zn concentrations under these Zn-limiting conditions, especially in edible sections” [1]. “One of the main causes of the prevalent Zn deficits in human populations is low dietary intake of Zn. The yield of green gram (*Vigna radiata* L.) decreases when cultivated in Zn-deficient soils. Zn application to the soil is effective in addressing their shortages, but it may be less effective at boosting its concentration in edible components, which is the main goal in addressing micronutrient deficiency” [2].

Thiourea is an essential component of plant physiology, both as a sulphhydryl molecule and as an amino compound similar to urea. Thiourea, a fertilizer that belongs to the SH group and has a

variety of biological actions in plants, is one of the fertilizers with 36.8 and 42.1% of N and S, respectively [3]. It has been observed that thiourea foliar spray and seed soaking boost plant growth and development as well as dry matter partitioning for higher grain production [4]. Sulphydryl compound thiourea (NH₂-CS-NH₂) is well known for promoting germination and breaking dormancy [5]. Use of thiourea, recognized as plant growth regulator may be helpful in this regard. The exogenous supply of growth regulators also affects essential physiological processes such carbon and nutrient assimilation, partitioning of photosynthates and utilisation efficiency in addition to hormonal regulation, differentiation and morphogenesis [6]. Keeping the above things in mind a study was carried out to Impact of zinc and thiourea application on growth and yield of summer green gram in loamy sand soils of Dantiwada (Gujarat).

2. MATERIALS AND METHODS

Geographically, Sardarkrushinagar is situated at 24° 19' North latitude and 72° 19' East longitude with an elevation of 154.52 meter above the mean sea level and situated in the North Gujarat Agro-climatic region. The current study was carried out at the Agronomy Instructional Farm, C. P. College of Agriculture, Sardarkrushinagar, during summer of 2020. The standard week wise meteorological data for the period of this investigation recorded at the Meteorological Observatory at Agronomy Instructional Farm, Chimanbhai Patel College of Agriculture, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar (Banaskantha). Climate of this region is sub-tropical monsoon type and falls under semi-arid region. In general, the monsoon is warm and moderately humid, winter is fairly cold and dry, while summer is

largely hot and dry. The experimental soil was loamy sand with 7.5% clay, 8.3% silt and 84.2% sand. The soil exhibited a low available N content (157.8 kg ha⁻¹), a medium available P₂O₅ content (33.8 kg ha⁻¹) and a medium available K₂O content (259.7 kg ha⁻¹) and available Zn (0.38 mg kg⁻¹).

Green gram was sown in field plots on first fortnight of March. With three replications, fifteen treatments were laid up in randomised block design (with factorial concept). The experiment included three levels of zinc and five treatments of thiourea application (Table 1). The variety of the crop was *Gujarat Mung 4*. The plant to plant and row to row spacing was maintained at 10 x 45 cm. The RDF *i.e* nitrogen (20 kg ha⁻¹) was applied as Urea, while phosphorous (40 kg ha⁻¹) was applied as DAP. Plant height and number of branches were measured on five randomly selected plants in a plot of each treatment during harvest in order to analyse the crop's growth pattern. According to established protocols, observations on yield parameters such as number of pods plant⁻¹, number of seeds pod⁻¹, seed index, seed yield, and stover yield hectare⁻¹ were recorded [7].

The data on different aspects of green gram crop were subjected to statistical analysis by adopting analysis of variance [7] as per the procedure of randomized block design with factorial concept by computer system at the computer centre, Department of Agricultural Statistics, Chimanbhai Patel College of Agriculture, Sardarkrushinagar. The value of calculated 'F' was worked out and compared with the value of table "F" at 5% level of significance. The value of S.Em.±, C.D. at 5% and coefficient of variation (C.V. %) were also calculated.

3. RESULTS AND DISCUSSION:

3.1 Effect of Zinc

3.1.1 Growth attributes

The number of branches plant⁻¹ is a key morphological feature that is directly associated to yield in green gram. With the application of 4.5 kg Zn ha⁻¹ along with the recommended dose of NPK (20-40-00 kg ha⁻¹), the maximum plant height (47.02 cm) and number of branches plant⁻¹ (7.25) were measured (Table 2). The positive effects of zinc application on these traits can be attributed to its catalytic or stimulatory effects on the majority of plant physiological and metabolic processes [6]. Similarly, Kuniya et al. reported that "application of 5.0 kg Zn/ha gave significantly higher plant height and number of branches plant⁻¹ of clusterbean" [8].

3.1.2 Yield attributes

Plant yield is directly correlated with yield attributing variables such pods plant⁻¹, seeds pod⁻¹ and seed index. The results showed that the application of 4.5 kg Zn ha⁻¹ along with recommended dose of NPK (20-40-00 kg ha⁻¹) resulted in the highest pods plant⁻¹ (15.92), seeds pod⁻¹ (7.91) and seed index (4.40) (Table 2). Application of Zn encouraged greater root growth and sink size development (number of pods plant⁻¹), which ultimately led to increased seed yield [9]. Ram and Katiyar also concluded that application of 10 kg Zn/ha significantly increased number of nodules and pods per plant, seeds per pod, test weight over preceding levels [10].

Table 1. Treatment details of experiment

Treatments	
A] Zinc (kg ha⁻¹)	
Zn ₁	1.5
Zn ₂	3.0
Zn ₃	4.5
B] Thiourea	
TU ₀	Control
TU ₁	Seed soaking with thiourea @ 500 ppm
TU ₂	Foliar spray of thiourea @ 500 ppm at 30 DAS
TU ₃	Foliar spray of thiourea @ 500 ppm at 45 DAS
TU ₄	Foliar spray of thiourea @ 500 ppm at 30 and 45 DAS

Table 2. Effect of zinc and thiourea on plant height, branches plant⁻¹, pods plant⁻¹, seeds pod⁻¹, seed index, seed yield and stover yield of green gram

Treatments	Plant height (cm)	Branches plant ⁻¹	Pods plant ⁻¹	Seeds pod ⁻¹	Seed index (g)	Yield (kg ha ⁻¹)	
						Seed	Stover
[A] Zinc (kg ha⁻¹)							
Zn ₁ 1.5	41.17	6.60	14.84	7.29	4.27	722	1234
Zn ₂ 3.0	42.63	6.67	15.26	7.59	4.36	816	1306
Zn ₃ 4.5	47.02	7.25	15.92	7.91	4.40	873	1378
S.Em.±	0.92	0.15	0.30	0.13	0.05	18	34
C.D. ($p \leq 0.05$)	2.65	0.44	0.86	0.38	NS	54	100
[B] Thiourea							
TU ₀ Control	40.84	5.90	12.52	6.89	4.23	739	1165
TU ₁ Seed soaking with thiourea @ 500 ppm	41.70	6.30	13.16	7.22	4.29	786	1263
TU ₂ Foliar spray of thiourea @ 500 ppm at 30 DAS	43.48	6.75	14.93	7.66	4.32	798	1305
TU ₃ Foliar spray of thiourea @ 500 ppm at 45 DAS	45.40	7.49	16.27	7.97	4.39	827	1359
TU ₄ Foliar spray of thiourea @ 500 ppm at 30 and 45 DAS	46.60	7.75	19.82	8.26	4.48	868	1438
S.Em.±	1.18	0.20	0.39	0.17	0.06	24	44
C.D. ($p \leq 0.05$)	3.42	0.57	1.11	0.49	NS	69	129
Interaction							
(Zn × TU)	NS	NS	NS	NS	NS	NS	NS
C. V. %	8.11	8.51	7.49	6.59	4.10	8.93	10.20

NS: Non-Significant



Fig. 1. Overview of field experiment

3.1.3 Yield

With the application of 4.5 kg Zn ha⁻¹ together with the recommended dose of NPK (20-40-00 kg ha⁻¹) significantly increased seed yield (873 kg ha⁻¹) and stover yield (1378 kg ha⁻¹) (Table 2). "The higher yields with zinc application could be ascribed to accelerated nutrient uptake helped the plant to put optimum growth. As these growth and yield attributes as well as nutrients uptake showed significant increase in seed yield, evidently resulted in higher yields with zinc fertilization" [9]. Similarly, Pooja and Sarawad found that application of 4 kg Zn/ha increased seed yield (2065 kg/ha) and straw yield (1734 kg/ha) in chickpea. [11].

3.2 Effect of Thiourea

3.2.1 Growth attributes

In green gram, the number of branches plant⁻¹ is a crucial morphological characteristic that is closely associated to yield. With the application of foliar sprays of thiourea @ 500 ppm at 30 and 45 DAS along with the recommended dose of NPK (20-40-00 kg ha⁻¹), the maximum plant height (46.60 cm) and number of branches plant⁻¹ (7.75) were recorded (Table 2). This favourable changes in the crop may be attributed to improved growth and development of the crop with thiourea treatment action potentially targeting the meristematic activity of apical tissues with stimulatory effects on cell division which promotes growth in shoot length and cell number for improved leaf area (mainly by higher sulphur and nitrogen nutrition). Thiourea boosted net photosynthetic rates as well as total chlorophyll and starch concentrations in the leaves, resulting in maximum plant height and branches [12]. Singh and Singh also reported that application of thiourea @ 500 ppm at pre-

flowering and pod initiation stages significantly increased plant height and branches than the control treatments [13].

3.2.2 Yield attributes

Plant yield is directly correlated with yield-attributing variables such pods plant⁻¹, seeds pod⁻¹ and seed index. The application of foliar sprays of thiourea @ 500 ppm at 30 and 45 DAS, combined with the recommended dose of NPK (20-40-00 kg ha⁻¹), resulted in the highest pods plant⁻¹ (19.82), seeds pod⁻¹ (8.26), and seed index (4.48) (Table 2). This could be related to the crop's accelerated nitrogen metabolism and extended moisture retention, particularly during the moisture stress stage, which may have contributed to a higher number of pods plant⁻¹ and seeds pod⁻¹ at harvest. Sarita et al. also observed that "application of thiourea @ 500 ppm at flower initiation recorded significantly higher number of pods plant⁻¹, seeds pod⁻¹ and test weight over rest of treatments" [14].

3.2.3 Yield

Significantly higher seed yield (868 kg ha⁻¹) and stover yield (1438 kg ha⁻¹) were recorded with application of foliar sprays of thiourea @ 500 ppm at 30 and 45 DAS along with recommended dose of NPK (20-40-00 kg ha⁻¹) (Table 2). The increased photosynthesis and effective transport of photosynthates towards the sink, which ultimately results in more seed yield, may be the cause of the improved yield through thiourea. The increase in yield observed in this study could be a reflection of thiourea's impact on growth and development. The increased number of branches plant⁻¹ resulted in more pods and consequently more seeds. Additionally, the rise in photosynthetic activity could have increased photosynthesis, which would have increased the

amount of assimilates transferred to the seeds and increased their weight. Devi et al. also observed that “application of thiourea @ 500 ppm increased the seed and stover yield of greengram by 58 and 84 per cent, respectively over control” [15].

4. CONCLUSION

In conclusion the results of experiment indicated that the application of 4.5 kg Zn ha⁻¹ and foliar sprays of thiourea @ 500 ppm at 30 and 45 DAS along with recommended dose of NPK (20-40-00 kg ha⁻¹) were significantly improved the growth and the yield of green gram.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mecarty JS, Dawson J, Lalrammawii C. Effect of micronutrients on growth and yield of green gram (*Vigna radiata* L.). The Pharma Innovation Journal. 2022;11(7): 2967-2969.
2. Cakmak I. Enrichment of fertilizers with zinc: An excellent investment for humanity and crop production in India. Journal of Trace Elements in Medicine and Biology. 2009;23:281-289.
3. Garg BK. Role of thiourea in improving the productivity of arid legumes. Journal Arid Legumes. 2007;4 (1):1-5.
4. Arora D. Effect of thiourea and zinc on growth, yield and quality of barley (*Hordeum vulgare* L.). Ph.D. Thesis, Rajasthan Agricultural University, Bikaner; 2004.
5. Garg BK, Burman U, Kathju S. Influence of thiourea on photosynthesis, nitrogen metabolism and yield of clusterbean (*Cyamopsis tetragonoloba* (L.) Taub.) under rainfed condition of Indian arid zone. Plant Growth Regulation. 2006;48(3):237-245.
6. Priyanka. Effect of zinc and thiourea application on growth and yield of green gram (*Vigna radiata* (L.) Wilczek). M.Sc. (Agri.) Thesis. Sri Karan Narendra Agriculture University, Jobner; 2017.
7. Cochran WG, Cox GM. Experimental design. 2nd edition Wiley, New York; 1957.
8. Kuniya N, Patel BB, Malav JK, Chaudhary N, Pavaya RP, Patel JK, Kumar S, Jat JR, Patel DM, Chaudhary MG, Patel BT, Patel VR. Yield and nutrient content and uptake by clusterbean (*Cyamopsis tetragonoloba* L.) as influenced by different levels of sulphur and zinc application under light textured soil. Journal of Pharmacognosy and Phytochemistry. 2019;8(3):2160-2163.
9. Kareti Siva S, Krishna S, Georg PJ. Effect of levels of phosphorus and zinc on growth and yield of Kabuli chickpea (*Cicer kabulium* L.). Journal of Pharmacognosy and Phytochemistry. 2017;6(4):1013-1016.
10. Ram S, Katiyar TPS. Effect of sulphur and zinc on the seed yield and protein content of summer mungbean under arid climate. International Journal of Science and Nature. 2013;4(3):563-566.
11. Pooja C, Sarawad IM. Influence of iron and zinc on yield, quality of chickpea and status of iron and zinc in postharvest soil. Agricultural Science Digest. 2019;39(1):31-35.
12. Yadav LR. Effect of bioregulators on productivity of wheat (*Triticum aestivum* L.) varieties under normal and late sown conditions. PhD Thesis, Rajasthan Agricultural University, Bikaner; 2005.
13. Singh RP, Singh D. Response of lentil to thiourea application under rain fed conditions of central India. International Journal of Current Microbiology and Applied Sciences. 2017;6(3):2556-2560.
14. Sarita, Sharma OP, Shukla UN, Yadav SK, Choudhary H. Effect of fertility levels and stress mitigating chemicals on yield attributes and yield of mungbean (*Vigna radiata* (L.) Wilczek). International Journal of Chemical Studies. 2019;7(2):1421-1424.
15. Devi S, Patel PT, Chaudhary KM. Effect of application of SH- compounds on yield, protein and economics of summer greengram under moisture stress in north Gujarat condition. Legume Research- An International Journal. 2015;38(4):542-545.

© 2023 Patel et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/103036>