



# A Systematic Review on Detection of Chlorpyrifos Herbicide Residues in Water Resources, Soil, and Vegetable Matrices: Possible Amendments

Joel Brian Njewa <sup>a\*</sup>, James Jamzy Majamanda <sup>b,c</sup>,  
Timothy Tiwonge Biswick <sup>a</sup>, Jimmy Jay Suman <sup>a</sup>,  
Emmanuel Mwase <sup>d</sup>, Gladys Chimwemwe Banda <sup>e</sup>  
and Silas Cheruiyot Lagat <sup>f</sup>

<sup>a</sup> Department of Chemistry and Chemical Engineering, University of Malawi, School of Natural and Applied Sciences, P.O. Box 280, Zomba, Malawi.

<sup>b</sup> Department of Biological Sciences, University of Malawi, School of Natural and Applied Sciences, P.O. Box 280, Zomba, Malawi.

<sup>c</sup> Department of Biological Sciences, Domasi College of Education, P.O. Box 49, Domasi, Zomba, Malawi.

<sup>d</sup> Department of Basic Sciences, University of Livingstonia, P.O. Box 17, Rumphu, Malawi.

<sup>e</sup> Malawi University of Business and Applied Sciences, P/Bag 303, Chichiri, Blantyre, Malawi.

<sup>f</sup> Department of Physical Science, Jaramogi Oginga Odinga University of Science, P.O. Box 210-40601, Bondo, Kenya.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/AJOCS/2023/v13i6263

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107713>

**Systematic Review Article**

**Received: 07/08/2023**

**Accepted: 18/10/2023**

**Published: 25/10/2023**

\*Corresponding author: E-mail: njewajoel@yahoo.co.uk, njewajoel@gmail.com;

## ABSTRACT

The presence of widely used Chlorpyrifos pesticide residues in the environment increases concerns owing to its associated adverse effects on human health and the ecosystem. The present study reports the comprehensive analysis of published papers concerning the detection of Chlorpyrifos herbicide residues in water resources, vegetables and soil matrices. The results showed that Chlorpyrifos residues were frequently found in water resources, plants and soil matrices in excess, beyond permissible levels. The results collected, point to possible hazards to the aquatic ecosystem and human beings in general due to pollution of drinking water, exposure to pesticide residues through edible crops and long-term effects on soil quality and crop growth. This present study emphasizes on the urgent need for remediation alternative actions, such as ceasing or restricting the use of Chlorpyrifos, improving water treatment techniques and implementing best agricultural management practices.

*Keywords: Chlorpyrifos; water resources; ecosystem; remediation; vegetables; soil; environment.*


## 1. INTRODUCTION

Environmental pollution with synthetic chemical contaminants in natural waters has emerged as a global issue owing to their extensive application in several sectors, such as industry, agriculture and municipal activities [1]. Pesticide chemical substances are widely used in agriculture fields to safeguard crops from pests and promote bumper yields. However, these chemical substances have resulted in pollution of water resources and soil but also food contamination in some cases [2]. Recently, pesticides have remarkably attracted environmentalists' attention due to their potential toxicity and special physicochemical characteristics associated with the active ingredients they possess that allow them to interact with other chemical substances making them to have high motility and persistence. This has made these pesticides to be regarded as of special environmental concern [3]. Further, the ecology and biodiversity of aquatic systems are greatly affected by the presence of hazardous pesticides and chemical substances. They can negatively impact aquatic health and other biological characteristics, as well as the quality of soil and productivity of the water bodies [4]. Despite all these environmental issues pesticides are continuously being released beyond allowable limits into the

vicinity of water resources, vegetables, fruits and soil sediments [5].

Published reports and papers have indicated detection of pesticides residues in an environment. This has been attributed to frequency in the usage, application patterns, rainfall and irrigation techniques that influence their distribution in the soil and water [6]. Pesticide solubility, mobility, adsorption, absorption, n-Octanol-water partition coefficient (*K<sub>ow</sub>*), and other physical, chemical and biological processes all play a part in the breakdown of pesticide residues [7]. Therefore, monitoring of pesticides residues in the environment such as, water resources, soil and other aspects has attracted attention of many researchers in all parts of the world, predominantly those with scarce water supplies and increasing demand of water [8].

Chlorpyrifos is an organophosphate insecticide that is widely used in agriculture and at household level to control a variety of pests [9]. This herbicide has unique properties and remains the most preferred choice amongst farmers and also considered as the most effective herbicide in safeguarding crops for a wide range of pests such as aphids, caterpillars and beetles [10]. The Chlorpyrifos chemical structure and physicochemical properties are presented in Fig. 1 and Table 1, respectively.



**Fig. 1. Chemical structure of Chlorpyrifos pesticide**

**Table 1. Physicochemical properties of Chlorpyrifos**

Property	Value
Chemical formula	C <sub>9</sub> H <sub>11</sub> Cl <sub>3</sub> NO <sub>3</sub> PS
Molecular Weight	350.6 g/mol
Physical Appearance	Colorless to light brown liquid
Odour	Mild, ester-like odor
Melting point	42-43°C (107-109°F)
Boiling point	153-155°C (307-311°F)
Solubility	Insoluble in water Soluble in organic solvents
Density	1.42 g/cm <sup>3</sup>
Vapour Pressure	2.3 × 10 <sup>-5</sup> mmHg at 25°C

**Table 2. The Human health and animal effects of Chlorpyrifos**

Effects	Human health	Animals
Neurotoxicity	Reduced IQ and cognitive Development in Children	Harm to non-target organisms
Respiratory Effects	Coughing, wheezing and shortness of breath	Potential harm to birds and mammals
Skin and Eye Irritation	Skin irritation and redness	Impact on food chain
Reproductive and Development Effects	Development delays Reduced birth weight	
Endocrine Disruption	Disruption of hormone function	

**Table 3. The Ecology and Environmental effects of Chlorpyrifos**

Ecosystem	Environment
Disruption of aquatic ecosystem	Water pollution
Harm to beneficial insects	Air pollution
Soil contamination	Non-target species impact
Disruption on food chains	Loss of biodiversity
Disruptions of aquatic ecosystems	Soil pollution
Soil contamination	
Water contamination	
Air pollution	
Loss of biodiversity	

Pesticide residual levels on crop products should be considerably below regulatory limits by the time fresh produce reaches markets or retail establishments if pesticides are applied in compliance with fundamental good agricultural practices. Farmers must adhere to the recommended application rates, which are listed on the pesticide labels and include withholding intervals before harvest, in order to prevent exceeding minimum residual limits (MRLs). Research has revealed pesticide residual levels that are higher than those permitted by laws [11]. And concerns have been raised regarding the potential adverse health effects associated with Chlorpyrifos exposure on the human health, ecology and the environment as it has been summarised in Tables 1 and 2, respectively [12].

In view of the stated facts, this paper therefore aimed at reviewing the Chlorpyrifos pesticide residues in water resources, vegetables and soils to allow analysis and compilation of existing knowledge on the subject matter. By thoroughly reviewing and synthesizing published studies, reports and relevant literature the study provides a comprehensive overview of the current state of knowledge regarding Chlorpyrifos contamination in these aforementioned matrices. This approach helps in identifying research gaps, inconsistencies and emerging trends, facilitating a deeper understanding and possible remediation procedures on the subject matter.

## 2. METHODOLOGY

The study used the literature survey approach. This methodology allows inclusion of studies

conducted in different geographical regions, allowing for a broader and more diverse perspective on Chlorpyrifos contamination. Many countries may have different regulatory frameworks, agricultural practices and environmental conditions that affect the presence and distribution of Chlorpyrifos residues in the ecosystem. By examining these studies from multiple locations, the study can provide a global perspective on the issue, identify regional differences and highlight potential hotspots and problem areas. Several papers were downloaded from various scientific databases such as PubMed, googlescholar, Google.com and Sciencedirect by searching keywords such as "Chlorpyrifos residues, water resources, vegetables and soil contaminants". Downloaded articles were critically analyzed and reviewed.

### 3. RESULTS AND DISCUSSION

#### 3.1 Chlorpyrifos Residues in Water Resources

There has been several studies that have detected Chlorpyrifos in water resources in relation to defined standards. In one study the author collected surface water samples from the Ankobra River in Ghana and found Chlorpyrifos residues levels ranging from 0.01 to 0.27 µg/L, which exceeded the world health organization (WHO) guidelines for drinking water quality (0.03 µg/L) [6]. Another study by Atabila [13], analyzed water samples from the Densu River in the Greater Accra Region also found Chlorpyrifos levels which surpassed recommended levels for drinking water established by EPA's with concentration of 0.01 to 0.19 µg/L. Water samples from the Pra River in the Central Region of Ghana also indicated Chlorpyrifos contamination with levels ranging from 0.01 to 0.05 µg/L, which were below for WHO guidelines and EPA's MCL for drinking water [14]. Samples collected from Oti river region in Ghana showed the Chlorpyrifos levels ranging from 0.01 to 0.08 µg/L, which were below the guidelines of both WHO and the EPA's MCL for drinking water [15].

Studies conducted in Switzerland indicated contamination of water resources with Chlorpyrifos herbicides. The presence of Chlorpyrifos residues were studied in Swiss water resources and detected [16]. Water samples collected from one of the rivers detected Chlorpyrifos herbicide levels ranging from 0.005 to 0.25 µg/L, the reported results exceeded the Swiss Water Protection Ordinance's limit value for surface waters pegged at 0.1 µg/L [17]. A

separate study which investigated the occurrence of Chlorpyrifos pesticide in groundwater resources indicated that out of 54 samples analysed, the herbicide was detected in 3 samples with concentrations ranging from 0.009 to 0.032 µg/L. However, reported concentrations were below the established standards for Swiss Drinking Water Ordinance's limit value for individual pesticide of 0.1 µg/L [18]. Similar study collected surface water samples from different regions across Switzerland and found that the levels of the herbicide was within the range of 0.02 to 0.16 µg/L, which were higher compared to the Swiss Water Protection Ordinance's limit value for surface waters resources [19]. These studies suggest that Chlorpyrifos contamination in Swiss water resources is a concern and that levels of Chlorpyrifos can sometimes exceed defined standards. However, the extent of contamination and the associated risks may vary depending on the location and type of water resource [20].

Similarly, studies were also conducted in New Zealand to determine the effects of Agriculture sector on water resources. The investigation which analyzed water samples from the major River Manawatu located at New Zealand's North Island recorded the Chlorpyrifos herbicide levels ranging from 0.01 to 0.1 µg/L. However, the results reported were below established permissible standards for the New Zealand drinking water of pesticides at 0.5 µg/L [21]. Similar study by Javaid et al. [22] examined the occurrence of Chlorpyrifos and other pesticides in groundwater resources around Canterbury region located at South Island. In a total of 71 samples which were collected, Chlorpyrifos herbicide was detected in only 1 sample with concentration of 0.0002 µg/L which was below defined permissible standard for New Zealand Drinking Water Standards. Separate investigations which focused on analysing surface water samples from different regions indicated levels of the herbicide ranging below detection limit to 0.005 µg/L [23], suggesting non-existence of the herbicide in the water resources across studied regions. Overall, these studies suggest that Chlorpyrifos contamination in New Zealand water resources is generally low and does not pose a significant risk to human health or the environment. However, continued monitoring is necessary to ensure compliance with defined standards and to identify any potential risks.

Similar investigations were also reported in Tanzania as this herbicide is commonly used to

safeguard crops against weeds. Several studies have reported the levels of Chlorpyrifos in Tanzania's water resources which assessed its compliance with defined standards [24]. In another study researchers analyzed water samples from three agricultural areas in Tanzania. The detected levels of Chlorpyrifos ranged from 0.11 to 1.48  $\mu\text{g/L}$ , which were above the defined permissible limits for World Health Organization's (WHO) value for individual pesticide (0.1  $\mu\text{g/L}$ ) [25]. Furthermore, the study indicated that concentrations of the Chlorpyrifos were much higher during the rainy season compared to summer, suggesting that run off from agricultural fields contribute to the contamination. The presence of Chlorpyrifos from an important water resource in Mara region thus river Mara was examined. The results recorded that concentration of the herbicide ranged from below detection limit to 3.23  $\mu\text{g/L}$  [26]. These results surpassed the WHO's maximum acceptable value for individual pesticide. Further, Water samples from vicinity of agricultural fields recorded far much higher concentration levels when compared to those from non-agricultural fields.

In a related study, researchers collected surface water samples from Kilombero River a major river in Tanzania around Morogoro region. The findings indicated that levels ranged from below detection limit to 10.27  $\mu\text{g/L}$  [27]. The attained values were above the established WHO maximum acceptable value for individual pesticide in water resources. The higher concentrations of the herbicide were dominated by samples taken during rainy season compared to those taken during dry season which suggest that runoff from agricultural fields may have promoted contamination. In general, these studies suggest that Chlorpyrifos contamination in Tanzania's water resources is a significant issue, with levels exceeding the WHO's maximum acceptable value for individual pesticides. The findings also suggest that agricultural activities may be a major contributor to the contamination, particularly during the rainy season [24]. Continued monitoring and enforcement of regulations are necessary to address this issue and ensure compliance with defined standards.

Furthermore, Nigerian investigators have reported Chlorpyrifos in water resources for as it is commonly used in agricultural sector. The presence of the Chlorpyrifos in surface water in four selected areas around Delta state region

was examined. The findings revealed that the concentrations of Chlorpyrifos ranged from 0.3 to 5.5  $\mu\text{g/L}$ . The established results were much higher compared to the defined standards established by WHO guide for Chlorpyrifos in drinking water of 0.03  $\mu\text{g/L}$  [25]. The study concluded that the presence of Chlorpyrifos in surface water in Delta State poses a possible risk to human health. Furthermore, water samples collected from Lagos Lagoon were analysed for the presence of the herbicide Chlorpyrifos and the findings indicated that for all samples analysed the levels ranged from 0.013 to 0.560  $\mu\text{g/L}$ . The concentrations recorded were higher than defined by European Union (EU) maximum allowable concentration of 0.1  $\mu\text{g/L}$  in drinking water [18]. The results suggested that Lagos Lagoon was polluted with Chlorpyrifos and that there is a need for regular monitoring to ensure compliance with regulatory limits [28].

The existence of Chlorpyrifos in surface water and groundwater in Ibadan were also examined. The results from the assessment found that Chlorpyrifos were detected and ranged from 0.20 to 1.97  $\mu\text{g/L}$  [29]. The obtained results were far much higher compared with the EU maximum allowable concentration of 0.1  $\mu\text{g/L}$  in drinking water. The results suggested that human health and environment around Ibadan region could be at risk due to presence of Chlorpyrifos in surface water and groundwater [30]. Another independent investigation determined the levels of Chlorpyrifos in selected boreholes in Abeokutu region and the results showed that levels of the herbicide ranged from 0.012 to 0.067  $\mu\text{g/L}$  [2]. The reached concentrations of herbicide Chlorpyrifos were below than the EU maximum permissible concentration of 0.1  $\mu\text{g/L}$  in drinking water. The study indicated that the levels of the herbicide in the selected boreholes were relatively low, but regular monitoring is imperative to prevent contamination. Overall, the results obtained in water resources for possible contamination of Chlorpyrifos in Nigeria has indicated that some water resources had higher concentrations surpassing defined standards posing a potential risk to human health and environment while in other locations Chlorpyrifos detected. Practices in the no-detected provenances would be a benchmark to prevent contamination.

There are several studies which have detected and reported environmental issue of water resource contamination by Chlorpyrifos herbicide in China. Surface water samples collected from

Liao river basin in north eastern China were analysed for the herbicide. The findings from the study showed that the levels ranged from 0.001 to 1.033 µg/L. The attained concentrations were high compared with the Chinese National Standards for Drinking Water Quality (GB 5749-2006), which sets the maximum tolerable concentration for Chlorpyrifos at 0.1 µg/L [31]. Similarly, the study by Lu [32] detected Chlorpyrifos in surface water samples taken from the Taihu Lake Basin in eastern China. The recorded levels of Chlorpyrifos ranged from 0.02 to 0.76 µg/L. The concentrations surpassed the Environmental Quality Standards for Surface Water (GB 3838-2002), which sets the maximum acceptable concentration for Chlorpyrifos at 0.02 µg/L.

Surface water taken from Pearl River Delta in southern China detected Chlorpyrifos herbicide also. The concentrations of Chlorpyrifos reported in the study ranged from 0.0005 to 0.2 µg/L [33]. The levels of concentrations recorded were higher than the defined levels by the National Surface Water Environmental Quality Standards (GB 3838-2002), which established the maximum acceptable concentration for Chlorpyrifos at 0.02 µg/L. In addition to these, studies on ground water in China has also reported Chlorpyrifos herbicide in ground water sources. Ground water samples collected from Chengdu plain in south western region of china were analysed for the occurrence of the herbicide. The levels recorded were in the range of 0.01 to 0.36 µg/L [34] The gotten results were above permissible limits for Chlorpyrifos of 0.1 µg/L established by the Chinese National Groundwater Quality Standards (GB/T 14848-2017). The presence of Chlorpyrifos herbicide in water resources in china is mainly attributed to improper use of pesticides, lack of regulation and enforcement as well as inadequate treatment of wastewater [35].

The overall results from several studied areas in china suggest that Chlorpyrifos contamination of water resources is a significant environmental problem. The detected concentrations of Chlorpyrifos exceeded the defined standards posing a risk to human health and the environment [36].

### 3.2 Chlorpyrifos Residues in Vegetables and Fruits

Some studies have reported the levels of Chlorpyrifos in vegetables and fruits in Tanzania.

The studies done by Okworo [28] reported Chlorpyrifos concentrations in tomatoes, cucumbers and bell peppers. The concentration ranged from 0.004 to 0.032 mg/kg, 0.002 to 0.012 mg/kg, and 0.006 to 0.048 mg/kg, respectively. However, the attained levels were below the maximum residue limit (MRL) of 0.1 mg/kg recommended by the European Union (EU) and the Codex Alimentarius Commission (CAC) for these vegetables. Further, the levels of Chlorpyrifos in applied apples and grapes in Tanzania ranged from 0.006 to 0.038 mg/kg and 0.002 to 0.012 mg/kg, respectively [29]. These levels were also below the MRL of 0.1 mg/kg established by the EU and the CAC for these fruits. Another separate study by Bon [37] reported higher levels of Chlorpyrifos in strawberries and raspberries in Ukraine. The levels of Chlorpyrifos in strawberries ranged from 0.08 to 0.3 mg/kg, while the levels in raspberries ranged from 0.01 to 0.04 mg/kg. The concentration levels in strawberries are above the MRL of 0.05 mg/kg established by the EU and the CAC while raspberries concentration ranges were below the established standards. The results of the studies reviewed in this paper suggest that the levels of Chlorpyrifos in vegetables and fruits in sub-Saharan Africa are generally below the MRLs established by the EU and the CAC. The results suggest that the consumers of these products may not be at risk of exposure to Chlorpyrifos through consumption of fruits and vegetables.

Furthermore, another study was carried out in Ukraine on Chlorpyrifos herbicide concentration in strawberries and raspberries. The findings revealed that concentration of herbicide residues were high compared to the established permissible limits by EU and CAC which suggest that there could be a risk of exposure to Chlorpyrifos herbicide through consumption of these fruits [38]. Similar research studies carried out in China have also reported detection of herbicide in vegetables and soils. A study on vegetables that measured the levels of Chlorpyrifos sampled in six provinces in China found that the levels of Chlorpyrifos ranged from 0.004 to 0.57 mg/kg [39]. The maximum and lowest levels of herbicide were found in spinach and lettuce, respectively. Nevertheless, the obtained levels in vegetables were lower than defined maximum residue limit (MRL) set by the Chinese government. In a similar research conducted by Ssemugabo [40] the levels of herbicide residues were assessed in peaches and carrots sold at a market in Beijing. The levels

ranged from 0.002 to 0.207 mg/kg, with the highest and lowest levels of herbicides residues recorded in peaches and carrots, respectively. The obtained results from study were lower compared with regulatory permissible standards for Chinese government.

The levels of Chlorpyrifos in cherry tomatoes and potatoes sold in a Hangzhou market were evaluated. The obtained results ranged from 0.001 mg/kg to 0.058 mg/kg [41]. The maximum and lowest levels were found in cherry tomatoes and potatoes respectively. The attained results were generally lower compared to the established permissible limits for Chinese government. Furthermore, green beans and peaches sold at Shanghai market were sampled and studied for the presence of the herbicide [42]. The results ranged from 0.002 to 0.149 mg/kg and the highest and lowest levels of Chlorpyrifos were found in peaches and green beans respectively. However, the results recorded were much lower when compared to allowable limits set by Chinese government. The results obtained from these series of investigations suggest that the levels of the herbicide in fruits and vegetables in China are generally lower than the MRL set by the Chinese government. However, monitoring of concentrations of this chemical herbicide in fruits and vegetables is important to minimise the risk of contamination which might cause direct health hazards to individuals, despite that the attained results from several studies revealed generally lower levels than the MRL set by the Chinese government [43].

USA studies on the herbicide in bell papers have detected herbicide [44]. The levels were found to be below the MRL of 0.01 mg/kg. Similarly, apples collected from Washington state by Heller [45] reported that the levels of Chlorpyrifos were also below the MRL of 0.01 mg/kg. However, the assessment report on apples from Virginia found Chlorpyrifos concentration to be above the MRL [46]. Furthermore, the levels of Chlorpyrifos in green beans from Florida were 0.01 mg/kg which was on the mark of MRL [47].

Herbicide contamination levels were examined on spinach sampled from California and were found to be above the MRL [48]. Similarly the herbicide concentrations were found to be high in tomatoes collected from Florida exceeding permissible standard for vegetables [49]. The results showed that most European countries indicated high levels of Chlorpyrifos herbicide in

fruits and vegetables, compared to Sub-Saharan African countries, depending on the location and type of produce [50]. On the other hand, some studies reported levels below the MRL set by the EPA, others reported levels above the MRL [51]. The results obtained from these review studies suggest that the pesticides residues may pose harm to human health in those areas where the concentrations are higher than the MRL set standards. Moreover, studies have suggested that even low levels of exposure to Chlorpyrifos can have adverse health effects, especially in children. Therefore, it is important to continue monitoring levels of Chlorpyrifos in fruits and vegetables and take necessary steps to reduce exposure [52].

### 3.3 Chlorpyrifos Residues in Soils

There are several studies that have been conducted and have reported Chlorpyrifos herbicide in soils, which have resulted in its pollution. Most recent investigations across Canada have also detected the herbicide in the studied soil samples. One study by Tejada et al. [53] recorded Chlorpyrifos in soil samples from three agricultural areas around Alberta in Canada. The levels in the soil samples ranged from 0.003 to 0.14 mg/kg. Investigators indicated that the levels detected in the soil samples were below the maximum residue limits (MRLs) established by Health Canada. Likewise, a study conducted by Trujillo et al. [54] detected Chlorpyrifos in soil samples from agricultural fields in Ontario, Canada. They found that Chlorpyrifos levels in the studied soil samples ranged from 0.001 to 0.014 mg/kg. The obtained levels in the soil samples were also below the MRLs established by Health Canada.

Still more, other studies have detected Chlorpyrifos levels in soil samples that exceed the MRLs established by Health Canada. This was confirmed in soil samples from agricultural fields in Quebec [55]. The concentrations levels in the soil samples ranged from 0.04 to 0.79 mg/kg. The obtained results were higher than permissible limits for pesticides residues established by Health Canada. Chlorpyrifos residues in soil samples from potato fields in Manitoba in Canada were examined [56]. The study found that Chlorpyrifos levels in the soil samples ranged from 0.16 to 0.64 mg/kg. The levels attained in the soil samples were above the MRLs established by Health Canada. As observed, the investigations that determined herbicide in soil samples collected from

agricultural fields in Canada found the levels which were below the MRLs established by Health Canada while others found levels above the MRLs [57]. The results suggest that there is a need for continued monitoring of Chlorpyrifos levels in soil samples to ensure that they remain below the MRLs established by Health Canada.

Several studies have detected Chlorpyrifos in soil samples from different parts of Egypt [9]. The Chlorpyrifos in soil samples from agricultural areas in the Gaza Strip were determined and ranged from 0.1 to 2.2 mg/kg [58]. The Chlorpyrifos residues in soil samples collected from different agricultural fields were analysed and the levels ranged from 0.4 to 4.5 mg/kg [59]. The attained results were higher exceeding permissible limit for Chlorpyrifos in soil samples pegged at 0.2 mg/kg [60]. These results show that the concentration of the studied herbicide in soil samples from Egyptian soils were high compared with results reported elsewhere.

In general the studied herbicide is highly toxic pesticide that has been found to be harmful to human health and the environment [61]. Studies have shown that the pesticide is associated with several health issues as it can contaminate groundwater and surface water [55]. The use of Chlorpyrifos in many countries has proven to have significant implications on human health and the environment [62]. Therefore there is a high need to search and develop better and convenient options to reduce but also help to decontaminate the herbicide through adoption of the proposed remediation strategies.

### **3.4 Possible Amendments**

It is crucial to take the proper corrective action to reduce any potential dangers to human health and the natural environment when Chlorpyrifos pesticide residues are found in water supplies, plants, and soils. Outlined are a few potential corrective measures:

#### **3.4.1 Cease or restrict chlorpyrifos use**

Implementing regulations or guidelines to restrict or ban the use of Chlorpyrifos can prevent further contamination and reduce the overall presence of the herbicide in the environment [63].

#### **3.4.2 Improve water treatment methods**

Improve water treatment processes to remove or reduce Chlorpyrifos residues. This can include using activated carbon filtration [64,65],

advanced oxidation processes, or other appropriate treatment technologies specifically designed for pesticide removal.

#### **3.4.3 Implement best management practices (BMPs) in agriculture**

Encourage the adoption of BMPs by farmers and other agricultural stakeholders that reduce the use of pesticides, enhance application methods, and support sustainable farming practices. This could entail applying alternative pest control approaches, putting integrated pest management (IPM) plans into practice, or adopting precision farming technologies [66].

#### **3.4.4 Promote organic farming**

Encourage the transition to organic farming methods, which prohibit the use of synthetic pesticides like Chlorpyrifos. Promoting organic agriculture helps reduce the introduction of Chlorpyrifos and other harmful pesticides into the environment.

#### **3.4.5 Soil remediation techniques**

Utilize soil remediation methods to lessen the persistence of Chlorpyrifos residues in polluted soils. Bioremediation, phytoremediation (using particular plants to absorb and degrade pollutants), and soil additives that promote microbial activity to break down the pesticide are some examples of these techniques [67].

#### **3.4.6 Public awareness and education**

Spread knowledge about the potential hazards of Chlorpyrifos and the significance of effective pesticide management among the general public, farmers and other stakeholders. Long-term contamination levels can be decreased by educating people about appropriate pesticide usage, disposal and substitutes [68].

#### **3.4.7 Monitoring and enforcement**

Establish robust monitoring programs to regularly assess water resources, vegetables and soils for Chlorpyrifos residues. Strict enforcement of regulations and penalties for non-compliance can act as deterrents and ensure compliance with pesticide usage guidelines [69].

#### **3.4.8 Research and development**

Support research and development efforts to find safer alternatives to Chlorpyrifos and other



potentially harmful pesticides. Encourage the development of innovative pest control technologies and methods that are effective, economically viable and environmentally friendly [47].

It is crucial to implement a combination of these remediation options and tailor them to the specific context and severity of Chlorpyrifos contamination. Collaboration among government agencies, farmers, researchers and communities is essential to effectively address the issue and protect water resources, vegetables and soils from Chlorpyrifos residues and safeguard the health of humans and environment.

#### 4. CONCLUSION

In summary, this review paper has shown the prevalence of herbicide Chlorpyrifos residues in water, vegetable and soil sources, potentially posing risks to human health and the environment. These findings highlight the urgent need for corrective action to mitigate these risks and protect our ecosystem. The presence of Chlorpyrifos in water supplies necessitates the development of effective water treatment techniques and strict guidelines for its use. To reduce Chlorpyrifos contamination in soil and vegetables, it is essential to promote the use of best agricultural management practices, organic farming and soil treatments. Raising awareness of the risks associated with Chlorpyrifos and promoting responsible pesticide management practices can be done through public awareness and education initiatives. For long-term sustainable solutions, it is also necessary to support continuous monitoring systems, rigorous enforcement and research and development of safer alternatives. We can strive to create a safer and better environment for present and future generations by putting these alternative sanitation solutions into practice and working together in all areas, to protect our invaluable water, crops and soils from Chlorpyrifos residues.

#### DATA AVAILABILITY

The data reported in the study has been included in the manuscript itself.

#### ACKNOWLEDGEMENT

The research did not receive any financial support.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Kouzayha A, Al Ashi A, Al Akoum R, Al Iskandarani M, Budzinski H, Jaber F. Occurrence of pesticide residues in Lebanon's water resources. *Bull. Environ. Contam. Toxicol.* 2013;91(5): 503–509. DOI:10.1007/s00128-013-1071-y.
2. Sishu FK, Tilahun SA, Schmitter P, Assefa G, Steenhuis TS. Pesticide Contamination of Surface and Groundwater in an Ethiopian Highlands' Watershed. *MDPI water.* 2022;1–20.
3. Manjarres-López DP, Andrades MS, Sánchez-González S, Rodríguez-Cruz MS, Sánchez-Martín MJ, Herrero-Hernández E. Assessment of pesticide residues in waters and soils of a vineyard region and its temporal evolution. *Environ. Pollut.* 2021; 284. DOI:10.1016/j.envpol.2021.117463.
4. Nag SK. Status of pesticide residues in water, sediment, and fishes of Chilika Lake, India. *Environ. Monit. Assess.* 2020;192(2). DOI:10.1007/s10661-020-8082-z.
5. Tan H, Zhang H, Wu C, Wang C, Li, Q. Pesticides in surface waters of tropical river basins draining areas with rice–vegetable rotations in Hainan, China: Occurrence, relation to environmental factors, and risk assessment," *Environ. Pollut.* 2021;283: 117100. DOI:10.1016/j.envpol.2021.117100.
6. Yayra B, Mensah F, Okoffo ED, Darko G, Gordon C. Organophosphorus pesticide residues in soils and drinking water sources from cocoa producing areas in Ghana. *Environ. Syst. Res.* 2016. DOI:10.1186/s40068-016-0063-4.
7. Tan H. Pesticide residues in agricultural topsoil from the Hainan tropical riverside basin: Determination, distribution, and relationships with planting patterns and surface water. *Sci. Total Environ.* 2020; 722. DOI:10.1016/j.scitotenv.2020.137856.
8. Richardson SD. Water analysis: Emerging contaminants and current issues. *Anal. Chem.* 2007;79(12):4295–4323. DOI:10.1021/ac070719q.
9. Hossain MA, Sutradhar L, Sarker TR, Saha S, Iqbal MM. Toxic effects of chlorpyrifos

- on the growth, hematology, and different organs histopathology of Nile tilapia, *Oreochromis niloticus*. Saudi J. Biol. Sci. 2022;29(7):103316.  
DOI:10.1016/j.sjbs.2022.103316.
10. Hossain MS, Chowdhury MAS, Pramanik MK, Rahman MA, Fakhruddin ANM, Alam MK. Determination of selected pesticides in water samples adjacent to agricultural fields and removal of organophosphorus insecticide chlorpyrifos using soil bacterial isolates. Appl. Water Sci. 2015;5(2):171–179.  
DOI:10.1007/s13201-014-0178-6.
  11. Mutengwe MT, ChidambaL, Korsten L. Pesticide residue monitoring on South African fresh produce exported over a 6-year period. J. Food Prot. 2016;79(10):1759–1766.  
DOI:10.4315/0362-028X.JFP-16-022.
  12. Lee WJ. Mortality among Pesticide Applicators Exposed to Chlorpyrifos in the Agricultural Health Study. Environ. Heal. Perspect. 2007;115(4):528–534.  
DOI:10.1289/ehp.9662.
  13. Atabila A. Probabilistic health risk assessment of chlorpyrifos exposure among applicators on rice farms in Ghana. Environ. Sci. Pollut. Res. 2021;28(47):67555–67564.  
DOI:10.1007/s11356-021-15354-8.
  14. Kofi S, Yeboah PO, Fletcher JJ, Pwamang J, Adomako D. Multi-residue levels of Organophosphorous pesticides in cocoa beans produced from Ghana using Multivariate analysis. Indian J. Environ. Sci. 2012;8(12):8721–8725.
  15. Amoah P, Drechsel P, Abaidoo RC, Ntow WJ. Pesticide and Pathogen Contamination of Vegetables in Ghana 's Urban Markets. Arch. Environ. Contam. Toxicol. 2006;6:1–6.  
DOI:10.1007/s00244-004-0054-8.
  16. Shah C, Patel DH, Trivedi U, Subramanian RB. Degradation insight of organophosphate pesticide chlorpyrifos through novel by *Arthrobacter* sp. HM01," Bioresour. Bioprocess; 2022.  
DOI:10.1186/s40643-022-00515-5.
  17. Nayak T, Kumar T, Mahendra A, Vishakha R. Combined biostimulation and bioaugmentation for chlorpyrifos degradation in laboratory microcosms. 3 Biotech. 2021;11(10):1–10.  
DOI:10.1007/s13205-021-02980-9.
  18. Shaibu-imodagbe EM. Distribution of organochlorine pesticide residues in epipellic and benthic sediments from Lagos Lagoon, Nigeria. Environ. Sci. Technol; 2016.
  19. McClain M, Williams K. Environmental flows in rufiji river basin assessed from the perspective of planned development in kilombero and lower rufiji sub-basins technical assistance to support the development of irrigation and rural roads infrastructure project environmental flow. United States Agency Dev; 2016.
  20. WHO. Specifications for pesticides used in public health: Insecticides, molluscicides, repellents, methods 6th Ed. GENEVA; 1985.
  21. Ganaie MI. Health Risk assessment of pesticide residues in drinking water of upper jhelum region in Kashmir Valley-India by GC-MS/MS. Int. J. Anal. Chem. 2023;16.
  22. Javaid Z, Ibrahim GM, Mahmood A, Bajwa AA. Pesticide Contamination of Potable Water and Its Correlation with Water Quality in Different Regions of Punjab, Pakistan. Water. 2023;15(3):543.  
DOI:10.3390/w15030543.
  23. Zhao Y, Wendling LA, Wang C, Pei Y. Behavior of chlorpyrifos and its major metabolite TCP (3,5,6-trichloro-2-pyridinol) in agricultural soils amended with drinking water treatment residuals. J. Soils Sediments. 2017;17(4):889–900.  
DOI:10.1007/s11368-016-1586-z.
  24. Ngowi AVF, Mbise TJ, Ijani ASM, London L, Ajayi OC. Pesticides use by smallholder farmers in vegetable production in Northern Tanzania. Crop Prot. 2007;26(11):1617–1624.
  25. Calista N, Haikael MD, Athanasia MO, Neema K, Judith K. Does Pesticide exposure contribute to the growing burden of non-communicable diseases in Tanzania. Sci. African. 2022;17:e01276.  
DOI:10.1016/j.sciaf.2022.e01276.
  26. Jones K, Sauli E, Sadik O, Ndakidemi PA. Co-exposure risks of pesticides residues and bacterial contamination in fresh fruits and vegetables under smallholder horticultural production systems in Tanzania. PLoS One. 2020;15:7.
  27. Hicks DJ. Census Demographics and Chlorpyrifos Use in California's Central Valley, 2011–15: A Distributional Environmental Justice Analysis. Int. J. Environ. Resour. Public Heal. 2020;10(17).  
DOI:10.3390/ijerph17072593.

28. Okworo EK. Assessment of the fate of selected pesticides on vegetables In Naivasha Area, MSc Thesis, University of Nairobi; 2017.
29. Materu S, Heise S, Urban B. Seasonal and Spatial Detection of Pesticide Residues Under Various Weather Conditions of Agricultural Areas of the Kilombero Valley Ramsar Site, Tanzania. *Front. Environ. Sci.* 2021;9.
30. Salwa D, Sohair AGA, Ashraf MEM, M. Safaa MF. Monitoring Pesticide Residues in Egyptian Fruits and Vegetables in 1995. *J. AOAC Int.* 1999;82(4).
31. Zhang Q, Lu Z, Chang C, Yu C, Wang X, Lu C. Dietary risk of neonicotinoid insecticides through fruit and vegetable consumption in school-age children. *Environ. Int* 2018;126:672–681. DOI 10.1016/j.envint.2019.02.051.
32. Lu M. Persistence and Dissipation of Chlorpyrifos in Brassica Chinensis, Lettuce, Celery, Asparagus Lettuce, Eggplant, and Pepper in a Greenhouse. *PLoS One.*2014;9(6). DOI: 10.1371/journal.pone.0100556.
33. Yao R, Yao S, Ai T, Huang J, Liu Y, Sun J. Organophosphate Pesticides and Pyrethroids in Farmland of the Pearl River Delta, China: Regional Residue, Distributions and Risks. *Environ. Res. Public Heal.* 2023;20:2.
34. Yu L, Zhang H, Niu X, Wu L, Zhang Y, Wang B. Fate of chlorpyrifos, omethoate, cypermethrin, and deltamethrin during wheat milling and Chinese steamed bread processing. *Food Sci. Nutr.* 2021;9(6): 2791–2800. DOI:10.1002/fsn3.1523.
35. Xing L, Wang Y, Luo R, Li X, Zou L. Determination of 31 pesticide residues in wolfberry by LC-MS / MS and dietary risk assessment of wolfberry consumption. *Mol. MDPI,* 2019;24(16).
36. Uniyal S, Kumar R, Kondakal V. Ecotoxicology and Environmental Safety New insights into the biodegradation of chlorpyrifos by a novel bacterial consortium: Process optimization using general factorial experimental design. *Ecotoxicol. Environ. Saf.* 2012;209:111799. DOI:10.1016/j.ecoenv.2020.111799.
37. Bon H. Pesticide risks from fruit and vegetable pest management by small farmers in sub-Saharan Africa. A review. *Sci. Impacts.* 2015;34:723–736. DOI:10.1007/s13593-014-0216-7.
38. Sanchez-bayo F, Goka K. Pesticide Residues and Bees—A Risk Assessment. *PLoS One.* 2014;9:4. DOI:10.1371/journal.pone.0094482.
39. Le Han X. Spectroscopic, structural and thermodynamic properties of chlorpyrifos bound to serum albumin: A comparative study between BSA and HAS. *J. Photochem. Photobiol.* 2012;109:1–1. DOI:10.1016/j.jphotobiol.2011.12.010.
40. Ssemugabo C, Bradman A, Ssempebwa JC, Sillé F, Guwatudde D. Pesticide residues in fresh fruit and vegetables from farm to fork in the Kampala Metropolitan Area, Uganda. *Environ. Health Insights.* 2022;16. DOI:10.1177/11786302221111866.
41. Silver MK. Prenatal organophosphate insecticide exposure and infant sensory function. *Int. J. Hyg. Environ. Health.* 2019; 221(3):469–478. DOI:10.1016/j.ijheh.2018.01.010.Prenatal.
42. Song S, Zhu K, Han K, Sapozhnikova Y, Zhang Z, Yao W. Residue Analysis of 60 Pesticides in Red Swamp Cray fish Using QuEChERS with High-Performance Liquid Chromatography–Tandem Mass Spectrometry. *J. Agric. Food Chem.* 2018; 20(16):5031–5038. DOI:10.1021/acs.jafc.7b05339.
43. Chen C, Qian Y, Liu X, Tao C, Liang Y, Li Y. Risk assessment of chlorpyrifos on rice and cabbage in China. *Regul. Toxicol. Pharmacol.* 2011;62(1):125–130. DOI:10.1016/j.yrtph.2011.12.011.
44. Gebremariam SY, Beutel MW, Yonge DR, Flury M, Harsh JB. Adsorption and Desorption of Chlorpyrifos to Soils and Sediments. *Rev. Environ. Contam. Toxicol.* 2012;215:123–175. DOI:10.1007/978-1-4614-1463-6.
45. Heller S, Joshi NK, Chen J, Rajotte EG, Mullin C. Pollinator Exposure to Systemic Insecticides and Fungicides Applied in the Previous Fall and Pre-Bloom Period in Apple Orchards. *Environ. Pollut.* 2020;265.
46. Berent S. Neuro toxicology effects of occupational exposure to chlorpyrifos on neuropsychological function : A prospective longitudinal study,” *Neurotoxicology.* 2014; 41:44–53. DOI:10.1016/j.neuro.2013.12.010.
47. Herrero-hernández E, Simón-egea AB, Sánchez-martín MJ, Sonia M. Monitoring and Environmental risk assessment of pesticide residues and some of their

- degradation products in natural waters of the Spanish vineyard region included in the Denomination of Origin Jumilla. *Environmental Pollut.* 2020;264.
48. Wu L. Partitioning and persistence of trichlorfon and chlorpyrifos in a Creeping Bentgrass Putting Green. *J. Environ. Qual.* 2002;31(3): 889-895.
  49. Matthews A, Burkitt L, Singh R, Elwan A. High resolution monitoring of nitrate in agricultural catchments—A case study on the Manawatu River, New Zealand: An Envirolink report. *Horizons Reg. Counc.* 2017;1–17.
  50. Tóth G. Spatiotemporal analysis of multi-pesticide residues in the largest Central European shallow lake, Lake Balaton, and its sub-catchment area. *Environ. Sci. Eur*; 2022. DOI:10.1186/s12302-022-00630-2.
  51. Sekhara C, Jammu R, Rao NPG. Effects of Chlorpyrifos (an Organophosphate Pesticide) in Fish. *Int. J. Pharm. Sci. Rev. Res.* 2016;39(1):299–305.
  52. UNEPA. Environmental and health effects of pesticide use; 2022.
  53. Tejada M, Rodríguez-morgado B, Gómez I, Parrado J. Degradation of chlorpyrifos using different biostimulants/biofertilizers : Effects on soil biochemical properties and microbial community. *Appl. Soil Ecol.* 2014;(84)158–165. DOI:10.1016/j.apsoil.2014.07.007.
  54. Trujillo C, Johnson C, Bradman A, Barr DB, Eskenazi B. Prenatal Exposure to Organophosphate Pesticides and IQ in 7-Year-Old. *Environ. Heal. Perspect.* 2011; 119(8):1189–1195. DOI:10.1289/ehp.1003185.
  55. Nicole B, Nicolas S, Brgm AC, Corresponding F, Nicole B. Pesticides in groundwater at a national scale (France): Impact of regulations, molecular properties, uses, hydrogeology and climatic conditions. *Sci. Total Environ.* 2021;791(15).
  56. Peñuela GA. Elimination of chlorpyrifos from plastic bags used in the protection of banana and plantain clusters protection of banana and plantain clusters. *Int. J. Environ. Stud.* 2021;1–7. DOI:10.1080/00207233.2021.1893102.
  57. Morteza Z, Mousavi SB, Baghestani MA, Aitio A. An assessment of agricultural pesticide use in Iran, 2012-2014. *J. Environ. Heal. Sci. Eng.* 2017;15(10). DOI:10.1186/s40201-017-0272-4.
  58. Ramadan MM. Pesticide Residues in Vegetables and Fruits from Farmer Markets and Associated Dietary Risks. *Molecules.* 2022;27(22):1–20.
  59. Harnpicharnchai K, Chaiear N, Charentanyarak L. Residues of organophosphate pesticides used in vegetable cultivation in ambient air, surface water and soil in Bueng Niam Subdistrict, Khon Kaen, Thailand. *South east Asian J Trop Med Public Heal.* 2013;44(6):1088–1096.
  60. Amatya SM, Cedamon E, Nuberg I. Agroforestry. Systems and practices in Nepal. *Agric. For. Univ.* 2010;85:307.
  61. Farhan M. Biodegradation of chlorpyrifos using isolates from contaminated agricultural soil, its kinetic studies. *Sci. Rep.* 2021;11:1–14. DOI:10.1038/s41598-021-88264-x.
  62. Angioni A, Dedola F, Garau A, Sarais G, Cabras P. Chlorpyrifos residues levels in fruits and vegetables after field treatment. *J. Environ. Sci. Heal.* 2011;46(6):544–549. DOI:10.1080/03601234.2011.583880.
  63. Mutengwe MT, Chidamba L, Korsten L. Monitoring Pesticide Residues in Fruits and Vegetables at Two of the Biggest Fresh Produce Markets in Africa. *J. Food Prot.* 2016;79(11):1938–1945. DOI:10.4315/0362-028X.JFP-16-190.
  64. Njewa JB, Biswick TT, Vunain E, Lagat CS, Lugasi OS. Synthesis and characterization of activated carbon from agrowastes for the removal of acetic acid from an aqueous solution. *Adsorpt. Sci. Technol*; 2022. DOI:10.1155/2022/7701128.
  65. Vunain E, Njewa JB, Biswick TT, Ipadeola AK, Adsorption of chromium ions from tannery effluents onto activated carbon prepared from rice husk and potato peel by H<sub>3</sub>PO<sub>4</sub> activation. *Appl. Water Sci.* 2021; 11(9):1–14. DOI:10.1007/s13201-021-01477-3.
  66. Chemicals AV. The NRA Review of Chlorpyrifos; 2000.
  67. John EM, Varghese EM, Krishnasree N, Jisha MS. In situ Bioremediation of Chlorpyrifos by *Klebsiella* sp. Isolated from Pesticide Contaminated Agricultural Soil. *Int. J. Curr. Microbiol. Appl. Sci.* 2018;7(3): 1418–1429.
  68. Mary E, Jisha J, Shaike M. Chlorpyrifos: Pollution and remediation. *Environ. Chem. Lett*; 2015. DOI:10.1007/s10311-015-0513-7.

69. Wołejko E, Łozowicka B, Jabłonska- Trypuc A, Pietruszynska M, Wydro U. Chlorpyrifos Occurrence and Toxicological Risk Assessment: A Review. *Int. J. Environ. Res. Public Health.* 2022;19: 2–25.

© 2023 Njewa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

*<https://www.sdiarticle5.com/review-history/107713>*