



Evaluation of Broad Bean (*Vicia faba* L) Genotypes for Genetic Variability, Heritability and Genetic Advance

**Sunil Kumar ^{a*}, Deepa H. Dwivedi ^{a++}, Putan Yadav ^b,
Virendra Kumar ^{b#} and Rameshwar Dayal ^b**

^a Department of Horticulture, Babasaheb Bhimrao Ambedkar University, Lucknow, Uttar Pradesh 226025 (U.P.), India.

^b Department of Vegetable Science, Acharya Narendra Deva of University, Agriculture & Technology, Kumarganj, Ayodhya, 224229 (U.P.), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i113304

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/107953>

Original Research Article

Received: 11/08/2023

Accepted: 18/10/2023

Published: 25/10/2023

ABSTRACT

The study investigated the phenotypic and genotypic coefficient of variation (PCV and GCV) and heritability estimates for various agronomic and nutritional traits in Broad bean (*Vicia faba*). The analysis aimed to identify traits with optimum variability and heritability, as well as to explore the potential presence of additive genetic variance in specific characteristics. The 18 genotypes were grown in randomized block design with the 3 replication during *Rabi* season in the year 2021-22 at the Research Farm- 1, Department of Horticulture, Babasaheb Bhimrao Ambedkar University, Vidya-Vihar, Raebareli Road, Lucknow-226025 (U.P.). The results showed that number of seed per plant, number of pod per plant and seed yield per plant exhibited high estimates of both PCV and GCV, indicating the presence of adequate variability for these particular characters within the

⁺⁺ Professor;

[#] Ph.D Research Scholar;

^{*}Corresponding author: E-mail: sunilkr09320@gmail.com;

population. Additionally, the heritability for the characters of 100 seed yield, number of seed per plant, titratable acidity (g/l), total soluble solids (TSS) and genetic advance for the number of seed per plant, number of pod per plant, seed yield per plant, 100 seed weight were found to be significant indicating that these characters were predominantly governed by additive gene action. The findings of this study hold significant implications for future breeding programs and genetic improvement strategies. The high heritability and genetic advance for the above-mentioned characters suggest that these characteristics can be effectively manipulated through selective breeding to develop improved and high-yielding plant varieties.

Keywords: Broad bean; GCV; PCV; heritability; genetic advance.

1. INTRODUCTION

The broad bean (*Vicia faba* L.) $2n = 2x = 12, 14$) is a tropical to temperate grain legume vegetable. It is commercially grown for green pod and dried seeds. It is locally known as Broad bean, faba bean, field bean, bell bean, tic bean, kala matar and bakala are two terms used in India (Singh et al., [1]). The broad bean belongs to the family Fabaceae. It is originated in Mediterranean region. It's also cultivated in Australia, South America, Mexico, Brazil, and Andes. The faba bean is a underutilized legume vegetable in India. In Indian condition, it is cultivated under rainfed conditions in Jammu and Kashmir, Himachal Pradesh, Uttar Pradesh, and Uttarakhand, as well as in other parts of South and North India. Pulses are famous as a health food that provides nutritional security to the millions of malnourished people throughout the world, particularly in South Asia and Africa.

Broad bean seeds contain relatively high proteins, carbohydrates, vitamins B, antioxidants and minerals. Protein content in different varieties varies from 26% to 41% [2]. Carbohydrate contents varies from 51% to 68%, of which major proportion is contributed by starch (41–53%) [3]. The antioxidant properties of phenolic compounds may provide excellent dietary source for natural antioxidants for prevention of chronic diseases and health promotion [4]. Pulses are an essential source of nutritional protein in vegan diets, particularly in impoverished nations.

After soybean (*Glycine max* L.) and pea (*Pisum sativum* L.), it is the third most significant legume green vegetable in worldwide. According to Department of Agriculture and Farmers Welfare, Directorate of Economics and Statistics targeted to total pulse production in 2020-21 was estimated at 25.60 million tonnes but the production was 25.58 million tons. Anonymous, [5].

Winter season varieties have 4-6 stems/plant and the spring season varieties have 1-2 stems/plant. The root is typical tap root with secondary roots and nodules due to *Rhizobium leguminosarum v. viciae*. Stem growth is indeterminate. There are leaves upto 5th to 10th node followed by raceme of 2-12 flowers in the leaf axils. There are 2 leaflets/leaf at the bottom and 6-8 leaflets/leaf at the top. Flowers, 2-3 cm long at anthesis, have a typically papilionaceous structure. They can be completely white, brown or violet. In most cases they concentrate their colour on black or brown melanin spots on the wings. Pods are short and erected in minor and paucijuga types (3-4 ovules pod) and long and hanging in major types (8-12 ovules per pod). Seed colour can be yellow, green, brown, black or violet and the seed may sometimes carry punctuations, brown spots or stripes around the hilum. The hilum can be black Ram, 2019). The annual herb faba bean has coarse and erect stalks. The blooms are big, white with dark purple patterns and borne in bunches on short pedicels. The fruit is a large, leathery pod that starts off green and becomes blackish-brown as it ripens. Through biological nitrogen fixation, it can contribute to the long-term sustainability or augmentation of total soil nitrogen fertility [6].

2. MATERIALS AND METHODS

The present investigation was carried out during *Rabi* season in the year 2021-22 at the Research Farm- 1, Department of Horticulture, Babasaheb Bhimrao Ambedkar University, Vidya-Vihar, Raebareli Road, Lucknow-226025 (U.P.). Lucknow is situated in the Gangetic alluvial plain of eastern Uttar Pradesh located at 26° 50 North latitude and 80° 52 East longitude and an altitude of 111 meter above mean sea level (MSL). The topography of the experimental field is plain. This region comes under Agro climatic zone 5th of Uttar Pradesh. (ICAR). The 18 genotypes of broad bean (*Vicia faba* L.) were planted in a Randomized Block Design during *Rabi* season

year 2021-22 under Lucknow condition. There were 3 replication, each having 18 plots and each plot replace by has 1 germplasm. The spacing row to row were 45 cm and plant to plant spacing 15 cm. The observation were recorded to dermination percent, plant height(cm), number of branches per plant, days to first flowering, days to 50% flowering, days to first fruit set, number of pod per plant, number of seed per pod, pod length (cm), pod diameter (mm), pod volume (ml), specific gravity, titratable acidity (g/l), total soluble solids(%), number of seed per plant, days to maturity, 100 seed weight (gm), seed yield per plant (gm).

2.1 Statistical Analysis

The phenotypic (PCV) and genotypic coefficient of variation (GCV) were calculated following standard procedure proposed by Burton and DeVane [7]. The categories for genotypic (GCV) and phenotypic coefficients of variation (PCV) given by Subramanian and Menon (1973) as low (0-10%), moderate (10-20%), and high (20% and above) were adopted in the present investigation also. Heritability in a broad sense was calculated by using the formula suggested by Allard, [8]. The estimates of broad sense heritability ($h^2_{(b)}$) were also classified into three categories as suggested by Robinson *et al.* [9]. Genetic advance was calculated by using the formula suggested by Lush [10]. Genetic advance as percent mean was also classified into 3 categories viz., low (0- 10%), moderate (10-20%), and high (20% and above as suggested by Johnson et al. [11].

3. RESULTS AND DISCUSSION

The estimate of genotypic coefficient of variation is of prime importance to breeder over genetic variation alone. It does not allow discussion as to which characters were showing the highest degree of variability, therefore, accurate relative comparison can be made with the help of genotypic and phenotypic coefficient of variation. In general phenotypic coefficient of variability were higher than the genotypic coefficient of variability for all the characters under study which indicates that the environment plays vital role in expression of the characters.

3.1 Estimation of Phenotypic and Genotypic Variance

The table explained that phenotypic variance was higher as compare to genotypic variance for all traits during search experiment. The

maximum variability was reported for 100 seed weight (407.14) followed by number of seed per plant (192.33), germination percent (145.38), plant height (95.45), pod volume (69.51).

The phenotypic variance for all characters under investigation were recorded higher compared to genotypic variances as reported by Ojha *et al.* [12] and Singh *et al.*[13] in dolichos bean. The influences of environmental factors in expression of these characters played a significant contribution. The vast range of phenotypic and genotypic variance were recorded in all evaluated germplasm for all the traits. The high phenotypic and genotypic variance were showed by the characters viz., 100 seed weight, number of seed per plant, germination % , plant height and pod volume. These results also reported by Muluaem et al., [14] while low phenotypic variance was bserved for number of branches per plant, number of seed per pod, specific gravity and titratable acidity. However, these traits should be given importance during selection process.

3.2 PCV and GCV

3.2.1 Phenotypic coefficient of variation

The phenotypic coefficient of variation was high for the characters number of seed per plant(46.40), number of pod per plant (37.13), seed yield/plant (35.41), 100 seed weight(32.99), pod volume (28.37), number of seed per pod (19.97), specific gravity (18.86), titratable acidity (18.86), days to first fruit set (18.58), number of branches per plant (18.20), germination % (16.97), Total soluble solids (16.62), pod length (16.17), pod diameter (12.58), plant height (11.80), days to first flowering (11.11), days to 50% flowering(9.53) and days to maturity(5.40) exhibited low PCV. In the present findings phenotypic coefficient of variations were observed to be higher than corresponding genotypic coefficient of variations for all the characters studied. However, the differences were narrow which implied their relative resistance to environmental variation. It also described that genetic factors were predominantly responsible for expression of those attributes and selection could be made effectively on the basis of phenotypic performance. The findings were similar to Kaliya *et al.* [15] Fatih and Hamza. [16]. The magnitude of phenotypic and genotypic coefficient of variation ranged from 5.40 (%) to 46.40 (%) and

4.13(%) to 45.95 (%). The findings were similar to Afeta et al. [17] Mulualem et al. [14].

3.2.2 Genotypic coefficient of variation

It is revealed from Table that genotypic coefficient of variation ranged from 4.13 (Days to maturity) to 45.95 (Number of seed per plant) for different characters under study. High genotypic coefficient of variation was observed in number of seed per plant (45.95), number of pod per plant (35.97), seed yield per plant (34.34), 100 seed weight (32.77), pod volume (27.43), number of seed per pod (18.92), acidity (18.57) days to first fruit set (17.01), number of branches per plant (16.96), specific gravity (16.96) and germination percent (16.57), Total soluble solids (16.28), pod length (12.85), pod diameter (11.18) plant height (11.17), days to first flowering (8.34), days to 50% flowering (8.02) and days to maturity (4.13) exhibited low GCV %.The genotypic and phenotypic coefficient of variation was high for the characters number of seed per plant, number of pod per plant, seed yield per plant, 100 seed weight, pod volume, number of seed per pod, titratable acidity, days to first fruit set, number of branches, specific

gravity, germination, total soluble solids, pod length, pod diameter, plant height, days to first flowering, days to 50 % flowering and days to maturity. The finding were similar to Kalia et al., [15] Mulualem et al. [14] Lal et al., [18], Fatih and Hamza [16].

3.3 Heritability

Results indicated that the heritability estimates were observed very high for 100 seed weight (98.69) followed by number of seed per plant (98.06), titratable acidity (96.97), Total soluble solids (95.87), germination percent (95.43), seed yield per plant (94.04), specific gravity (93.91), number of pod per plant (93.84) and volume (93.47). Result indicated that the heritability estimate were observed high for number of seed per pod (89.79), plant height (89.48), number of branches per plant (86.84), days to first fruit set (84.27), pod diameter (78.97) and days to 50% flowering (70.90). Result indicated that the heritability estimate were observed moderate for pod length (63.14), days to maturity (58.40), and days to first flowering (56.30).The estimates of heritability in broad sense ($h^2 b$) and genetic




Fig. 1. GCV (%) and PCV (%) for different characters of broad bean (*Vicia faba L*)

Table 1. Mean range, genotypic and phenotypic variation, heritability, genetic advance percent mean, and genotypic and phenotypic coefficient of variation

Charecters	Mean	Min	Max	Genotypic variation	Phenotypic variation	Heritability (%)	GA% mean	GCV (%)	PCV (%)
Germination Percentage	71.07	44.58	93.75	138.74	145.38	95.43	33.35	16.57	16.97
Plant Height (cm)	82.77	67.23	101.73	85.41	95.45	89.48	21.76	11.17	11.80
Number of branches	5.12	3.55	6.33	0.75	0.87	86.84	32.56	16.96	18.20
Days to first flowering	58.82	48.66	68.22	24.04	42.71	56.30	12.88	8.34	11.11
Days to first fruit Set	19.96	13.67	25.00	11.53	13.68	84.27	32.16	17.01	18.53
Days to 50% Flowering	61.59	50.00	70.67	24.42	34.44	70.90	13.92	8.02	9.53
Number of pod per plant	11.34	5.00	20.44	16.64	17.73	93.84	71.77	35.97	37.13
Number of seed per pod	2.82	1.80	3.80	0.29	0.32	89.79	36.93	18.92	19.97
Pod length (cm)	6.72	5.57	9.30	0.75	1.18	63.14	21.03	12.85	16.17
Pod diameter(mm)	14.91	11.91	18.37	2.78	3.52	78.97	20.47	11.18	12.58
Pod volume (ml)	29.39	24.00	55.00	64.97	69.51	93.47	54.63	27.43	28.37
Specific gravity	0.87	0.66	1.21	0.02	0.02	93.91	33.87	16.96	17.51
Titrateable Acidity(g/l)	0.40	0.26	0.54	0.01	0.01	96.97	37.67	18.57	18.86
Total soluble solids	6.96	5.40	9.30	1.28	1.34	95.87	32.83	16.28	16.62
Number of seed per plant	29.89	11.93	60.15	188.60	192.33	98.06	93.74	45.95	46.40
Days to Maturity	122.65	112.33	132.33	25.60	43.84	58.40	6.49	4.13	5.40
100 Seed weight	61.17	38.62	110.20	401.81	407.14	98.69	67.06	32.77	32.99
Seed yield/ plant (gm)	16.89	6.53	29.51	33.63	35.76	94.04	68.60	34.34	35.41

advance in per cent of mean estimated for eighteen characters have been presented in table. Heritability ranged from 56.30 (days to first flowering) to 98.69 (100 seed weight). Very high estimates of broad sense heritability (>90%) was estimated for 100 seed yield (98.69), number of seed per plant (98.06), titrateable acidity (96.97), total soluble solids (95.87), germination percent (95.43), seed yield per plant (94.04), specific gravity (93.91), number of pod per plant (93.84) and pod volume (93.47). High estimates of heritability (70-90%) were observed for number of seed per pod (89.79), plant height (89.48), number of branches (86.84), days to first fruit set (84.27), pod diameter (78.97) and days to 50% flowering (70.90). The result were similar to the findings of Kaliya et al. [15]. The moderate

estimates of heritability (50-70%) for pod length (cm) (63.14), days to maturity (58.40), days to first flowering (56.30). The similar result were found by Mulualet et al. [14].

3.4 Genetic Advance

Genetic advance is defined as the difference between the mean genotypic value of the selected lines and the mean genotypic value of the parental population. In the present study, GAM (Genetic advance as per cent of mean) was observed to be high in number of seed per plant (93.74%), number of pod per plant (71.77%), seed yield per plant (68.60%), 100 seed weight (67.06%), pod volume (54.63%), titrateable acidity (37.67%). Number of seed per




Fig. 2. Heritability and GA (%) in different characters of broad bean (*Vicia faba* L)

pod (36.93%), specific gravity (33.87%), germination percent (33.35%), Total soluble solids (32.83%), number of branches per plant (32.56%), days to first fruit set (32.16%), plant height (21.76%), pod length (21.03%) and pod diameter (mm) (20.47%). Medium estimates of GA as percent of mean was recorded for days 50% flowering (13.92%), days to first flowering (12.88%) and the lowest GA was recorded in days to maturity (6.49%). The high estimates of genetic advance in per cent of mean (>20%) were recorded for number of seed per plant (93.74%), number of pod per plant (71.77%), seed yield per plant (68.60%), 100 seed weight (67.06%), pod volume (54.63%), titratable acidity (37.67%), number of seed per pod (36.93%), specific gravity (33.87%), germination percent (33.35%), total soluble solids (32.83%), number of branches per plant (32.56%), days to first fruit set (32.16%), plant height (21.76%), Pod length (cm) (21.03%) and Pod diameter (20.47%). Similar result were found by Fikreselassie and Seboka [19,20,21] Medium estimates of GA as percent of mean (>10-20%) was recorded for days 50% flowering (13.92%), days to first flowering (12.88%) and the lowest GA was recorded in days to maturity (6.49%). The similar result were reported by Afeta et al. [17].

4. CONCLUSION

It was concluded that the number of seed per plant, number of pod per plant and seed yield per

plant exhibited high estimates of both PCV and GCV, indicating the presence of adequate variability for these particular characters within the population. Additionally, the heritability for the characters of 100 seed yield, number of seed per plant, titratable acidity (g/l), total soluble solids (TSS) and genetic advance for the number of seed per plant, number of pod per plant, seed yield per plant, 100 seed weight per plant were found to be significant indicating that these characters were predominantly governed by additive gene action. The findings of this study hold significant implications for future breeding programs and genetic improvement strategies. The identified characters with high heritability and high genetic advance can be effectively manipulated through selective breeding to develop improved and high-yielding varieties.

ACKNOWLEDGEMENTS

The authors are grateful to Indian Institute of Vegetable Research (IIVR) Jakkhini, Varanasi, Uttar Pradesh 221305 for providing the seeds to conducting the research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Singh AK, Bharati KC, Manibhushan NC, Pedpati and Anitha. An assessment of

- faba bean (*Vicia faba* L.) current status and future prospect. Academic J. 2013,8(50): 6634 – 6641.
2. Picard J. Some Results Dealing with Breeding Protein Content in (*Vicia faba* L.) Protein Quality from Leguminous Crops, seminar held at Dijon, France, 3-5 November; 1976. (EUR, 5686 EN. 339347).
 3. Cerning J, Saposnik A, Guilbot A. Carbohydrate composition of horse beans (*Vicia faba*) of different origins. Cereal Chem. 1975;52:125–138.
 4. Oomah B, Tiger N, Olson M. and Balasubramanian P. Phenolics and antioxidative activities in narrow-leafed lupins (*Lupinus angustifolius* L.) Plant Food Hum. Nutr. 2006;61:91–97.
 5. Anonymous. Ministry of Agriculture and Farmers Welfare, Department of Agriculture, Cooperation and Farmers Welfare Directorate of economics and Statistics Third advance Estimation of Production of food grain for 2020-21 Available:(<http://agricoop.nic.in>)
 6. Lindemann WC, Glover CR. Nitrogen Fixation by Legumes. Guide A-129 Cooperative Extension Service. New Mexico State University; 2003.
 7. Burton GM, De Vane EM. Estimating heritability in tall Fescue from replication clonal material. Agron. J. 1953;45:478-481.
 8. Allard R.W. Principle of Plant Breeding. John Wiley and Sons Inc., New York, USA. 1960;485.
 9. Robinson HF, Comstock RE, Harvey PH. Estimation of heritability and the degree of dominance in corn. Agron. J. 1949;41:353-359.
 10. Lush JL, Heritability of quantitative characters in farm animals. Hereditas (suppl.).1949;35:256-261.
 11. Johnson HW, Robinson HF, Comstock RE. Genotypic, phenotypic correlations in soybean and their implication in selection. Agronomy Journal. 1955;47:477- 48
 12. Ojha, Sankar V, Nath S, Singh R. Genetic variability in chickpea (*Cicer arietinum* L.) .Progressive Res. 2010;5(2):275-276.
 13. Singh S, Singh PK, Singh DR, Panday VB, Srivastava RC. Genetic variability and character association study in dolichos bean Indian J. Hort. 2015;72(3):343-346
 14. Mulualet al. Correlation and path coefficient analysis of yield and its components in faba bean (*Vicia faba* L.) germplasm. International J. Biodiversity and Conservation. 2013;3(8):376-382
 15. Kalia P, Sood S, Singh. Genetic variability in faba bean (*Vicia faba* L.) for pod yield and its contributing traits. Indian J.Genet.and Plant Breed. 2003;63(3):261-262.
 16. Fatih EA, Hamza. Performance Assessment, Genetic Variability, Heritability, Genetic Advance and Correlation Coefficient Analysis for Yield and Some Agro -Morphological Traits in Faba Bean (*Vicia faba* L.) Genotypes in the Northern State, Sudan. Int. J. Curr. Microbiol. App. Sci. 2017;6(11): 1206-1214.
 17. Afeta T, Tesse B, Lule D, Kebede W. Study of genetic variability among released Faba Bean (*Vicia faba* L.) varieties in central and southern highlands of Ethiopia. 2020;2(4) :18-24/18.
 18. Lal k, Yadav CB, Nath S, Dwivedi DK. Correlation and Path Analysis in Faba bean (*Vicia faba* L.). Trends in Biosciences. 2019;12(11). Print:ISSN0974-8431,779-787.
 19. Fikreselassie M, Seboka H. Genetic variability on seed yield and related traits of elite Faba bean (*Vicia faba* L.) Genotypes. Pakistan J. of Bio. Sci. 2012;6(2):116-117.
 20. Mahalanobis PC. Studies on generalized distance in Statistics. Proc .Nat .Inst .Sci . India. 1936;2(1):49 -55.
 21. Ram H. Vegetable Breeding Principle and Practices; Third edition. 2019;701-704.

© 2023 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/107953>