



# **Occurrence of Potentially Pathogenic Bacteria in Commercially Sold Seafood from Misamis Occidental, Philippines**

**Nelfa Dumaloan-Canini <sup>a\*</sup>, Emily E. Cadelinia <sup>a</sup>  
and Jessie Jane Olarte-Bala <sup>a</sup>**

<sup>a</sup> *Department of Natural Sciences, College of Arts and Sciences, Misamis University, Ozamiz City, Philippines.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. Author NDC designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Author EEC and Author JJOB managed the analyses of the study. Author JJOB managed the literature search. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/EJNFS/2024/v16i21386

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112884>

**Original Research Article**

**Received: 07/12/2023**  
**Accepted: 11/02/2024**  
**Published: 15/02/2024**

## **ABSTRACT**

Seafood is a nutritious food that is an enviable component of a healthy diet. Nevertheless, there are health risks associated with the consumption of seafood. This study aimed to investigate the occurrence of potentially pathogenic *Vibrio* spp., *Staphylococcus aureus*, and *Salmonella* spp. in 200 seafood samples purchased from several retail outlets in Ozamiz City, Philippines. Different samples analyzed using standard microbiological techniques were clam, shrimp, squid, and fish (anchovy and oil sardines). The *Vibrio* species identified and isolated were *V. parahaemolyticus* (47%), *V. cholerae* (16%) and *V. alginolyticus* (9%). Thirty percent (30%) of the samples analyzed were positive for *S. aureus* and 9% of the samples were contaminated with *Salmonella* sp. The total aerobic plate count mean log<sub>10</sub> cfu/g of clams, shrimps, head/gills, intestine of fish (oil sardine, anchovy), and squids were considered as

\*Corresponding author: Email: nelfacanini9@gmail.com;

medium to high risk of transmitting pathogenic bacteria. Moreover, the clams and head/gills of oil sardines were considered to be at medium to high risk of transmitting pathogenic *V. parahaemolyticus*. The findings revealed that seafood samples analyzed microbiologically were loaded with potentially pathogenic bacteria. This result is of public health concern because these organisms are documented as causes of food-borne diseases.

**Keywords:** Pathogenic bacteria; food-born pathogen; shrimp; squid; marine environment.

## 1. INTRODUCTION

“Seafood a vital food component for a large portion of the world’s population” [1]. “However, there are health risks linked to the ingestion of seafood. One of the main risks involves the consumption of improperly cooked or raw seafood that might be contaminated by foodborne pathogens in the marine environment” [2]. “Various outbreaks of bacterial diseases associated with the consumption have been reported in both developed and developing countries” [3]. “From these seafood-borne bacteria, *Vibrio* spp. *S. aureus*, and *Salmonella* spp comprise an important group of pathogenic bacteria that frequently cause human disease and even death when the contaminated food is consumed raw or improperly cooked” [4].

“In the Asian region, *Vibrio* spp. has been recognized as the leading cause of foodborne outbreaks in various countries, including Japan, India, China, Korea, and Malaysia” [5,6]. “Investigations have shown that many outbreaks may lead to acute gastroenteritis, including diarrhea, headache, vomiting, nausea, and fever, and are caused by ingestion of infected seafood” [7]. “For food samples, temperature abuse may be instigated by improper storage or a long holding time on the exhibit rack at the retail level without proper temperature control” [8,9]. “Among more than 20 *Vibrio* species identified to be linked with human disease, *V. cholerae*, *V. parahaemolyticus*, and *V. vulnificus* are the pathogenic species of *Vibrio* that pose the greatest threat to human health” [10]. “Most of these *Vibrio* species secrete enterotoxins in water, foods, or in the gastrointestinal tract” [11].

“Aside from *Vibrio* spp., several bacteria in seafood may cause infection in humans. Examples of these bacteria that may cause foodborne illness are *Salmonella*, *Staphylococcus*, *Shigella*, and *Listeria*” [12]. “*S. aureus* is one of the most popular sources of gastroenteritis worldwide, which is caused by the ingestion of food that contains preformed toxins. Humans are common carriers of *S. aureus* in the

nose, throat, and skin infections. Therefore, the organism can be just transmitted to seafood during handling” [13].

“Seafood is not the usual habitat of *Salmonella*, but still, the occurrence of *Salmonella* in seafood is escalating. Hence, fishery products have been recognized as major spreaders of foodborne pathogens” [14,15]. “*Salmonella* is associated with fish and fishery products, thereby causing a public health problem. Various authors have reported the incidence of enterotoxigenic *S. aureus* and *Salmonella* in seafood” [16]. Infection with these bacteria may affect only the intestines or may spread from the intestinal tract to the blood vessels and then to other body sites.

Although several technological alternatives that could augment the quality of the food supply have been implemented, instances of food-borne diseases caused by consumption of seafood still emerge. In Iligan City, Philippines [17], an outbreak of cholera killed two people. Health authorities have begun a series of tests of water samples in the villages. Other reports said some of the infected individuals were associated with the consumption of improperly cooked seafood, particularly oysters.

In the provinces of Zamboanga del Sur, Lanao del Norte, and Misamis Occidental Philippines, massive mortality were observed among pond-cultured shrimp due to *Vibrio harveyi* which was isolated purely from the lymphoid organs and hepatopancreas of affected shrimp, and histopathological observations indicated a severe bacterial infection [18]. Disease problems have been recorded in shrimp-finfish polyculture in Misamis Occidental due to bacteria [19]. Nevertheless, studies on the presence of potentially pathogenic bacteria in seafood are scarce. This study aimed to examine the occurrence of potentially pathogenic *Vibrio* sp., *S. aureus*, and *Salmonella* sp. in fresh seafood sold from several retail outlets in the province of Misamis Occidental, Philippines, and to highlight that consumption of raw and ingestion of

improperly cooked sea foods may pose serious health hazards.

## 2. MATERIALS AND METHODS

### 2.1 Sample Collection and Processing

Simple random sampling of two hundred (200) samples of commercially sold seafood such as clams (40), shrimps (40), squids (20), and fish [anchovy (50) and oil sardines (50)] were purchased from 10 randomly selected retail outlets in the province of Misamis Occidental, Philippines. Samples were placed in sterile, labeled, sealed plastic bags and were transported to the laboratory under refrigerated conditions for analysis. The samples were aseptically removed from the container. After the skin ranging, sterile forceps and knife appropriate for opening the shell to remove the intestine and gills were used. For shrimp samples, the shell was peeled and detached from the fresh part and intestine. For fishes, samples of head/gills, intestines, and flesh were analyzed microbiologically, whereas only the flesh of squid was examined using standard microbiological techniques. For the clams, the flesh and intervalve water were used for bacterial analyses. Approximately 1g from each sample was weighed and dispensed into 9 ml of sterile distilled water and shaken vigorously to mix properly. Ten-fold dilutions were made, and dilutions ( $10^{-6}$  and  $10^{-7}$ ) were selected for microbial enumeration using the standard pour plate method by Fawole and Oso [20].

### 2.2 Enrichment Procedures

Five (5) grams of each shellfish flesh was incised using a sterile scalpel after removal of the carapace. These 5 g meat samples were homogenized in 45 ml of 3% NaCl containing 1% alkaline peptone water (APW, pH: 8.6) using a sterile blender. The shellfish homogenates were incubated at 37°C for 18 h [21].

### 2.3 Isolation Procedures and Identification of the Bacterial Colonies

Analysis of *Vibrio spp.* in seafood samples was performed in triplicate. Standard procedures of the U.S. Food and Drug Administration (FDA) was used to detect the contamination of potentially pathogenic *Vibrio* species [22] with some modifications. Following incubation, shellfish homogenates were inoculated using an inoculating loop on thiosulfate citrate bile salts

sucrose agar media (TCBS, Hi-Media, India) and incubated at 37°C for 18 h [23]. For each seafood sample, 25 g were homogenized in 225 ml sterile 0.1% peptone water using a sterile blender. A 10-fold serial dilution was prepared in sterile 0.1% peptone water as described in the Bacteriological Analytical Manual [24]. The spread plate method was carried out using 10 and ten dilutions on TCBS agar. The TCBS plates were incubated at 37°C for 18-24 h, and counts were made for each colony type. Discrete colonies were aseptically subcultured into fresh agar plates to obtain pure colonies.

Colonies recognized as discrete were carefully examined macroscopically for cultural characteristics. Pure isolated colonies were subjected to Gram staining and grown at different salt concentrations by placing the colonies into tubes containing peptone water and 0%, 3%, 6%, and 10% NaCl, and these tubes were incubated at 37°C for 24 h [25]. In addition, all bacterial colonies from different samples grown on TCBS plates were chosen to be streaked on the surface of trypticase soya agar slants (TSA; Oxoid, UK) added with 2% NaCl and incubated for 24 h at 37°C. Further identification of *Vibrio sp.* was performed using physiological, morphological, and different biochemical tests available in the laboratory. The isolates were identified by comparing their characteristics with those of recognized taxa, as illustrated and described by [26-29]. To isolate and identify *S. aureus*, 10 g of each seafood sample was homogenized with 90 ml of a sterile 0.1% peptone water solution. Serial dilutions of the homogenate were performed using sterile peptone water and plated in triplicates on Baird Parker agar (Merck 1.05406). After incubation at 35 °C for 48 h, the distinctive black colonies with apparent halo of *S. aureus* were subjected to Gram staining to clarify bacterial morphology. Staphylococci are spherical Gram-positive bacteria that occur in microscopic cluster. *S. aureus* forms a relatively large yellow colony on rich medium, whereas *S. epidermidis* has a comparatively small white colony [30].

For the detection of *Salmonella sp.*, the EN ISO 6579 reference method was used. This process currently involves non-selective pre-enrichment in buffered peptone water followed by selective enrichment in Rappaport–Vassiliadis soya broth and Muller–Kauffmann tetrathionate/novobiocin broth followed by selective plating of these on Salmonella–Shigella agar. Incubation was made at 35°C, and colonies resembling *Salmonella sp.*

or *Shigella sp.* *Salmonella* will produce hydrogen sulfide (H<sub>2</sub>S) gas but will not ferment lactose. The resulting bacterial colonies will appear colorless with black centers. Any suspect salmonella strains that are isolated are then confirmed by biochemical and serological test methods.

### 2.4 Bacterial Counts and Data Analysis

Following proper incubation, dilutions of 30–300 colonies were selected and counted. The number of colony-forming units per g (cfu/g) of seafood was computed by multiplying the number of bacteria by the dilution. Total aerobic plate count (TAPC), *Vibrio sp.*, *S. aureus*, and *Salmonella sp.* counts that were attained for cfu/g of food were converted into log<sub>10</sub> values. Seafood was classified as having a no-to-low risk of transmitting pathogenic bacteria if the TAPC was 5.0 log<sub>10</sub> cfu/g and if the counts of *Vibrio sp.*, *S. aureus*, and *Salmonella sp.* were 3.0 log<sub>10</sub> cfu/g. Furthermore, if the value is at least 5.0 log<sub>10</sub> cfu/g

for the total bacterial counts and at least 3.0 log<sub>10</sub> cfu/g for *Vibrio sp.*, *S. aureus*, and *Salmonella sp.*, the seafood is classified as medium-to-high risk of transmitting pathogen [31,32].

## 3. RESULTS AND DISCUSSION

### 3.1 The Occurrence of *Vibrio sp.*

Seafood samples analyzed microbiologically in this study showed varying degrees of bacterial contamination. The results on the contamination of *Vibrio sp.* are summarized in Tables 1a and 1b. Of the 200 seafood samples tested, 65.5% were contaminated with potentially pathogenic *Vibrio sp.* The high incidence of *Vibrio sp.* could be due to the nature of these bacteria, which are known as halophilic waterborne and commonly dwell in environmental water sources worldwide. brackish water, marine environments, and freshwater may support the growth of these organisms, which are pathogenic to humans [33].

**Table 1a. Distribution and frequency of occurrence of *Vibrio sp.* isolates from fresh seafood samples**

Samples	No. Tested (%)	No. Positive for <i>Vibrio sp.</i>
FISH		
Anchovy	50 (25)	39 (78)
Sardines	50 (25)	42 (84)
SHRIMP	40 (20)	18 (45)
CLAM	40 (20)	22 (55)
SQUID	20 (10)	10 (50)
Total	200 (100)	131 (65.5)

**Table 1b. Distribution and frequency of occurrence of *Vibrio sp.* isolates from fresh seafood samples**

Seafood Samples	Number (%) of positive samples			Total Positive for <i>Vibrio</i> (%)
	<i>V. parahaemolyticus</i>	<i>V. cholerae</i>	<i>V. alginolyticus</i>	
FISH				
Anchovy				
Gills	14 (28)	06 (12)	04 (08)	24 (48)
Intestine	05 (16)	04 (08)	02 (04)	11 (22)
Meat	04 (08)	00 (00)	00 (00)	04 (08)
SARDINES				
Gills	16 (40)	04 (08)	02 (04)	22 (44)
Intestine	8 (20)	02 (04)	02 (04)	12 (24)
Meat	06 (12)	02 (04)	00 (00)	08(16)
SHRIMP	10 (25)	06 (15)	02 (05)	18(45)
CLAM	12 (30)	06(15)	04(10)	22(55)
SQUID	06 (30)	02 (10)	02 (10)	10(50)
Total number (%) of positive samples	81 (40.5) /200	32 (16)/200	18 (09) /200	131(65.5)

The samples that harbored most of the *Vibrio* spp. were the fish in which the gills obtained most of the isolates. This could be due to the gill physiology, which serves as a sieve that filters water that comes into the fish from their water ecosystem. Fish could be predisposed to gill disease because the gill is used for the exchange of gasses, and the fish will be hassled because of the intricacy in the exchange due to the heavy load of microbes in water, causing the gill to be infected [34].

Among the *Vibrio* species isolated, *V. parahaemolyticus* was the predominant species (40.5%), followed by *V. cholera* (16%) and *V. alginolyticus* (09%). The findings of this study agree favorably with the results of several authors in a similar study conducted in Brazil, China, India, and Bursa province, Turkey [35,36]. On the other hand, varying rates have been reported in some other studies. In Nigeria, *V. cholerae* was the most frequently isolated *Vibrio* sp. followed by *V. parahaemolyticus* [37].

The predominant *V. parahaemolyticus* is salt tolerant but very sensitive to heat and is destroyed by cooking. However, gastroenteritis is a common cause of foodborne diseases in many Asian countries such as China, Japan, and Taiwan and is recognized in the United States as the leading cause of human gastroenteritis [38]. This species is associated with the consumption of insufficiently or raw cooked seafood, poor handling of seafood during preparation, and improper post-harvest storage conditions [39]. This pathogen is commonly isolated from a variety of seafood including oyster, crab, scallop, lobster, shrimp, and fish [40].

In this study, 16% of the samples tested were positive for *V. cholera*. This bacterium favors alkaline surroundings to thrive well [41]. Perhaps the presence of this bacterium was linked to the low salinity in various locations due to the influx of freshwater into the ocean. This species can cause cholera and is typically a waterborne illness. Food-borne cases, primarily fish outbreaks, are also known carriers of this microorganism.

In this study, the prevalence of *V. alginolyticus* in seafood was 9%. This finding was close to the result of Baffone et al. (2006), who reported 8.16% in clams in Italy. In contrast, Hassanin (2007) reported higher infection rates for *V. alginolyticus* in shrimps in Egypt. This bacterium is one of the most popular *Vibrio* species

occurring in marine environments and seafood (Neetoo et al. 2022). *V. alginolyticus* is an opportunistic pathogen, and its pathogenicity is thought to be similar to that of *V. parahaemolyticus* [42]. In India and Taiwan, this *Vibrio* species is linked with a white spot in shrimp while the zoonotic risk of this pathogen has been implicated in soft tissue, wound and ear infections in humans [43]. *Vibrio* spp. and several pathogenic species are commonly associated with outbreaks of *Vibrio* infections due to ingestion of raw or improperly cooked seafood, water contaminated with human feces or sewage, inappropriate post-harvest storage conditions, or poor handling of seafood during preparation.

### 3.2 Occurrence of *S. aureus* and *Salmonella* sp.

The incidence of *S. aureus* in this study was found to be higher than that of *Salmonella* sp. *S. aureus* was detected in 30% of the 200 raw samples of seafood, whereas the incidence of *Salmonella* sp was 9%. The head/gills of fish (oil sardines & anchovy) were heavily loaded with *S. aureus*. Four isolates of *S. aureus* were obtained from shrimp samples, three from clams, and two from squids (Table 2).

It is well documented that freshly caught seafood is free from *S. aureus*, and contamination occurs upon handling [44]. Other conditions such as poor personal hygiene, inadequate refrigeration, and delay in processing are associated with staphylococcal contamination. *S. aureus* is a major cause of food poisoning and a range of extraintestinal infections [45]. Staphylococcal food poisoning is caused by a preformed enterotoxin produced by the growth of bacteria in food that is ingested [46].

The overall prevalence of *Salmonella* sp. in 100 seafood samples tested was in agreement with that reported elsewhere. The findings of the present study showed that the organism was not recovered from the gills and flesh of the fish but was isolated from their intestine only. This result is in agreement with the work of Green [47], in which the organism was never recorded from the flesh of the fish but was documented from the viscera and epithelium. Several scientists believe that fish are possible carriers of *Salmonella*, which accumulate in their intestines for somewhat short periods of time, and some believe that fish become actively infected by *Salmonella* [48,49]. Furthermore, the mollusk

and crustacean samples also harbored colonies of *Salmonella*.

The possible cause of *Salmonella* contamination in seafood might be farm runoff and fecal contamination from nearby domesticated or wild animals and poor water quality. In addition to poor distribution, retail marketing, handling, and high stocking, preparation practices, densities, and high water temperature may be responsible for increased *Salmonella* contamination in shrimp [50]. Other investigations of live bivalve mollusks confirmed that this bacterium could endure coastal water salinity. The bacteria may be found in 0%–2.5% of live shellfish, and intake of these foods may cause salmonellosis, characterized by enteric fever along with gastroenteritis and diarrhea. Similarly, the United States Food and Drug Administration data revealed that *Salmonella* was the most common contaminant of fish and fishery products [51]. Approximately 12% of the foodborne outbreaks related to the consumption of fish are due to bacteria, including *Salmonella*.

### 3.3 Bacterial Counts in Seafood

The mean log<sub>10</sub> cfu/g of *Vibrio* spp., *S. aureus*, and *Salmonella* sp. and total aerobic plate counts (TAPC) values are presented in Table 3. The clam and shrimp samples obtained the highest TAPC values of 7.61 and 7.48, respectively. These molluscan bivalves and decapod crustaceans are filter feeders, and they may be apt to hoard microbes in nearby waters, which may also contain these microorganisms. They

are typically grown and collected in near-shore; shallow estuarine waters [52] and are therefore expected to harbor high concentrations of pathogenic organisms. If this seafood is eaten after a very mild heat treatment or raw, it poses a significant health risk to consumers. The seafood that may be linked with human infection caused by *V. parahaemolyticus* are bivalve mollusks, crustaceans, and finfish, especially those served raw or superficially cooked and exposed to temperature abuse [53]. Of the fishes, the head/gills of oil sardines and anchovy harbored considerable TAPC values of 7.42 and 6.78, respectively.

The TAPC value suggests that samples of shrimps, clams, head/gills, and intestines of fishes as well as squids examined in the study were considered as medium to high risk of transmitting pathogenic bacteria (not within acceptable limits, i.e. <5.0 log<sub>10</sub> cfu/g for total counts). Moreover, the clams and head/gills of oil sardines were also considered to be at medium to high risk of transmitting pathogenic *V. parahaemolyticus*. From the zoonotic point of view, contamination of potentially pathogenic bacteria in all samples of seafood tested suggests a possible risk to the health of people consuming raw or improperly cooked seafood. Naturally, fish and other seafood have immunity to infections, but their immunity decreases when the concentration of microorganisms is on the increase [54]. Bacteria in seafood improperly cooked could be transmitted to humans as they establish themselves in the intestine, particularly those potentially pathogenic, causing problems,

**Table 2. Distribution and frequency of *Salmonella* spp. and *S. aureus* isolates from fresh seafood samples**

Seafood Samples	Number (%) of positive samples/number of samples tested	
	<i>Salmonella</i> sp.	<i>S. aureus</i>
FISH		
Anchovy		
Gills	00 (00) /50	08 (16)/50
Intestine	02 (04) /50	06 (12)/50
Meat	00 (00) /50	04 (08)/50
Sardines		
Gills	00 (00) /50	12(24)/50
Intestine	04 (08) /50	04/(08)50
Meat	00 (00) /50	08(16)/50
SHRIMP	04 (10) /40	06(15)/40
CLAM	06(15)/40	08(20)/40
SQUID	02 (10) /20	04(20)/20
Total number (%) of positive samples/total number of samples tested	18 (09) /200	60 (30) /200

**Table 3. Total aerobic plate counts, *Vibrio* spp., *S. aureus*, and *Salmonella* spp. counts mean log<sub>10</sub> cfu/g (ml) ± standard deviation of seafood in Ozamiz City**

Seafood Item	n	TAPC	<i>Vibrio</i> isolates of <i>S. aureus</i> <i>Salmonella</i>				
			<i>V. parahaemolyticus</i>	<i>V. cholerae</i>	<i>V. alginolyticus</i>		
Fish							
Anchovy	25						
Head/gills		6.78±1.7	2.87±1.0	1.23±0.7	1.02±0.6	1.19±0.4	00
Intestine		5.60±0.5	1.14±0.9	1.70±0.9	0.37±0.6	1.10±0.5	1.66±1.2
Meat		3.64±2.0	1.05±1.1	00	00	1.20±0.1	00
Oil Sardines	25						
Head/gills		7.42±2.1	3.04±0.4	1.21±1.0	1.03±0.1	2.19±0.5	00
Intestine		5.90±1.9	1.06±0.5	1.07±1.6	1.87±0.6	1.19±0.5	1.15±0.3
Meat		3.51±1.0	1.04±0.7	1.02±0.8	00	1.05±0.5	00
Shrimp	20	7.48±1.8	2.69±0.7	1.10±0.7	0.87±0.8	1.73±1.1	1.06±1.8
Clam	20	7.61±1.5	3.14±0.5	1.05±0.4	0.89±0.4	1.93±0.5	1.59±0.2
Squid	10	5.55±0.9	1.20±0.2	1.02±0.2	1.04±0.9	1.14±1.3	1.01±0.9

including diseases of various kinds. These bacteria enter the body and then discharge endotoxins that irritate the stomach and bowels. This toxin might lead to stomach cramps, diarrhea, and fever and is called foodborne infection.

#### 4. CONCLUSION

Seafood is considered an essential element in the diet and plays a vital role in the development of foodborne diseases. This study revealed that seafood samples analyzed microbiologically could be a source of foodborne bacterial pathogens such as *Vibrio* sp. *S. aureus* and *Salmonella*. The contamination of these bacteria could be attributed to cross-contamination from the surroundings and handling by the sellers and thus constitute a potential health hazard to the public.

Nevertheless, fish and other seafood have to be properly cooked before consumption, and proper quality control measures have to be adopted in the culturing, processing, harvesting, and consumption of seafood. Therefore, monitoring the contamination of potentially pathogenic bacteria in harvested seafood is vital for the sustenance of public health. Further studies have to be conducted to include other seafood products in the microbial analyses and to examine other pathogens probably in seafood.

#### ACKNOWLEDGEMENTS

The authors are grateful to the Misamis University for funding this study and to the Natural Science Department Laboratory

personnel especially to Jendelou Buhian and Mark Anthony Manapsal for the assistance during the conduct of the study. Worth to mention is Noel U. Canini for the support during the sampling period.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Gopal TS. Foods from the ocean for nutrition, health, and wellness. In Nutritional and Health Aspects of Food in South Asian Countries Academic Press. 2020;113-122.
- Song X, Zang J, Yu W, Shi X, Wu Y. Occurrence and identification of pathogenic *Vibrio* contaminants in common seafood available in a Chinese traditional market in Qingdao, Shandong Province. *Frontiers in Microbiology*. 2020;11(1488).
- Ndraha N, Wong HC, Hsiao HI. Managing the risk of *Vibrio parahaemolyticus* infections associated with oyster consumption: A review. *Comprehensive Reviews in Food Science and Food Safety*. 2020;19(3):1187-1217.
- Gautam RK, Venugopal V. Electron beam irradiation to control biohazards in seafood. *Food Control*. 2021;130:108320.
- Tunung R, Margaret SP, Jeyaletchumi P, Chai LC, Zainazor TC, Ghazali FM, et al. Prevalence and quantification of *Vibrio* in raw salad vegetables at retail *Vibrio parahaemolyticus* from fresh, low-

- temperature preserved, dried and salted seafood products; 2021.
6. Loo KY, Letchumanan V, Law JWF, Pusparajah P, Goh BH, Ab Mutalib NS, Lee LH. Incidence of antibiotic resistance in *Vibrio spp.* Reviews in Aquaculture. 2020;12(4):2590-2608.
  7. Gourama H. Foodborne pathogens. Food Safety Engineering. 2020;25-49.
  8. Canini ND, Bala JJO, Maraginot EN, Mediana BCB. Evaluation of Street Food Vending in Ozamiz City. Journal of Multidisciplinary Studies. 2013;1(1).
  9. Zhou B, Luo Y, Huang L, Fonseca JM, Yan H, Huang J. Determining effects of temperature abuse timing on shelf life of RTE baby spinach through microbial growth models and its association with sensory quality. Food Control. 2022;133:108639.
  10. Mok JS, Ryu A, Kwon JY, Park K, Shim KB. Abundance, antimicrobial resistance, and virulence of pathogenic *Vibrio strains* from molluscan shellfish farms along the Korean coast. Marine Pollution Bulletin. 2019;149:110559.
  11. Bonnin-Jusserand M, Copin S, Le Bris C, Brauge T, Gay M, Brisabois A, Midelet-Bourdin G. *Vibrio* species involved in seafood-borne outbreaks (*Vibrio cholerae*, *V. parahaemolyticus* and *V. vulnificus*): Review of microbiological versus recent molecular detection methods in seafood products. Critical Reviews in Food Science and Nutrition. 2019;59(4):597-610.
  12. Owusu-Apenten R, Vieira E. Microbial Foodborne Disease Outbreaks. In Elementary Food Science. Cham: Springer International Publishing. 2022; 171-196.
  13. Dai J, Huang J, Wu S, Zhang F, Li Y, Rong D, Wu Q. Occurrence, Antibiotic Susceptibility, Biofilm Formation and Molecular Characterization of *Staphylococcus aureus* Isolated from Raw Shrimp in China. Foods. 2023;12(14):2651.
  14. Olgunoğlu İA. *Salmonella* in Fish and Fishery Products. University of Adiyaman Vocational School of Kahta Turkey; 2012. Available:www.intechopen.com ISBN. 978-953-307-782-6.
  15. Parlapani FF, Boziaris IS, DeWitt CAM. Pathogens and their sources in freshwater fish, sea finfish, shellfish, and algae. In Present Knowledge in Food Safety. Academic Press. 2023;471-492.
  16. Sorsa M, Mamo G, Abera L. Major fish-borne bacterial and parasitic zoonoses in Ethiopia: A review. Int. J. Fauna Biol. Studies. 2019;6:50-58.
  17. Man D. On cholera outbreak. The Mindanao Examiner Regional Newspaper; 2008.
  18. Pena LDDL, Lavilla-Pitogo CR. Mortality in pond-cultured shrimp *Penaeus monodon* in the Philippines associated with *Vibrio harveyi* and white spot syndrome virus. 2003;38(2):59-61.
  19. Fitzsimmons K, Bolivar R, JunRey Sugue. Survey of Tilapia-Shrimp Polycultures in the Philippines (10NSR3E). Final Report. 2003;14-15.
  20. Fawole MO, Oso BA. Laboratory manual of Microbiology: Revised edition pectrum books Ltd. Ibadan. 2001;127.
  21. Wafaa MKB, Walaa AH, Amani FA. Detection of *Salmonella* and *Vibrio* species in some seafood in Alexandria. J. American Science. 2011;7(9):663-668.
  22. Nordstrom JL, Kaysner CA, Blackstone GM, Vickery MCL, Bowers JC, De Paola A. Effect of intertidal exposure on *Vibrio parahaemolyticus* levels in Pacific Northwest oysters. Journal of Food Protection. 2004;67(10):2178-2182.
  23. Wafaa MKB, Walaa AH, Amani FA. Detection of *Salmonella* and *Vibrio* species in some seafood in Alexandria. J. American Science. 2011;7(9):663-668.
  24. BAM, Bacteriological Analytical Manuel 8 Ed, Garthersbury, M.D U.S.A; 1998.
  25. Lahfi SK, Kuhne M. Occurrence of *Vibrio* spp. in blue mussels (*Mytilus edulis*) from the German Wadden Sea. International Journal of Food Microbiology. 2007;116:297-300.
  26. Jolt JG, Krieg NR, Sneath PHA, Stanley JT, Williams ST. Bergey's manual of systematic bacteriology, 9 ed. Williams Wilkins Co. Baltimore, Maryland, pp: 786. Journal of Food Microbiology. 1994;126:71-75.
  27. Cheesbrough M. District laboratory practice in tropical countries. Cambridge University Press. 2006;434.
  28. FAO/WHO.FAO/WHO guidance to Governments on the application of HACCP in small and/or less-developed food businesses. 2007;24:2045-2054.
  29. Oyeleke SB, Manga SB. Essentials of Laboratory Practicals in Microbiology.



- Tobest publisher, Minna, Nigeria. 2008;36-75.
30. Todor K. Online Textbook of Bacteriology. 2012;1.  
Available:www.textbookofbacteriology.net
  31. Mensah P, Manu DY, Darko KO, Ablordey A. Street foods in Accra, Ghana: how safe are they? Bulletin of the World Health Organization. 2002;80(7):546–554.
  32. Jans C, Sarno E, Collineau L, Meile L, Stärk KD, Stephan R. Consumer exposure to antimicrobial resistant bacteria from food at Swiss retail level. Frontiers in Microbiology. 2018;9:362.
  33. Harrison JP, Hoellein TJ, Sapp M, Tagg AS, Ju-Nam Y, Ojeda JJ. Microplastic-associated biofilms: A comparison of freshwater and marine environments. Freshwater microplastics: Emerging Environmental Contaminants? 2018;181-201.
  34. Megarajan S, Ranjan R, Xavier B, Ghosh S. Training Manual on Cage Culture of Marine Finfishes. 2016;7-12.
  35. Dash HR, Mangwani N, Chakraborty J, Kumari S, Das S. Marine bacteria: Potential candidates for enhanced bioremediation. Applied Microbiology and Biotechnology. 2013;97(2):561-571.
  36. Mus, Tulay Elal, Figen Cetinkaya, Umut C. Occurrence of *Vibrio*, *Salmonella* and *Staphylococcus aureus* in retail fresh fish, mussel and shrimp. ACTA VET. BRNO. 2014;83:75–78.  
DOI: 10.2754/avb201483020075
  37. Bonnin-Jusserand M, Copin S, Le Bris C, Brauge T, Gay M, Brisabois A, Midelet-Bourdin G. *Vibrio* species involved in seafood-borne outbreaks (*Vibrio cholerae*, *V. parahaemolyticus* and *V. vulnificus*): Review of microbiological versus recent molecular detection methods in seafood products. Critical Reviews in Food Science and Nutrition 2019;59(4):597-610.
  38. Letchumanan V, Loo KY, Law JWF, Wong SH, Goh BH, Ab Mutalib NS, Lee LH. *Vibrio parahaemolyticus*: The protagonist of foodborne diseases. Progress In Microbes & Molecular Biology. 2019;2(1).
  39. Bedane TD, Agga GE, Gutema FD. Hygienic assessment of fish handling practices along production and supply chain and its public health implications in Central Oromia, Ethiopia. Scientific Reports. 2022;12(1):13910.
  40. Roobab U, Fidalgo LG, Arshad RN, Khan AW, Zeng XA, Bhat ZF, Aadil RM. High-pressure processing of fish and shellfish products: Safety, quality and research prospects. Comprehensive Reviews in Food Science and Food Safety 2022;21(4):3297-3325.
  41. Kostiuk B, Becker ME, Churaman CN, Black JJ, Payne SM, Pukatzki S, Koestler BJ. *Vibrio cholerae* Alkalizes Its Environment via Citrate Metabolism to Inhibit Enteric Growth *In-vitro*. Microbiology Spectrum. 2023;11(2):e04917-22.
  42. Cai H, Yu J, Li Q, Zhang Y, Huang L. Research Progress on Virulence Factors of *Vibrio alginolyticus*: A Key Pathogenic Bacteria of Sepsis; 2022.
  43. Wodajo, W. Review on Zoonotic Bacterial Diseases of Fish. ARC Journal of Animal and Veterinary Sciences. 2020;6(1):1-9.
  44. Stratev, D, Stoyanchev, T, Bangieva, D. Occurrence of *Vibrio parahaemolyticus* and *Staphylococcus aureus* in seafood. Italian Journal of Food Safety. 2021;10(4).
  45. Zhang P, Miao X, Zhou L, Cui B, Zhang J, Xu X, Wang X. Characterization of oxacillin-susceptible *mecA*-*Staphylococcus aureus* from food poisoning outbreaks and retail foods in China. Foodborne Pathogens and Disease. 2020;17(11):728-734.
  46. Pal M, Ketchakmadze D, Durglishvili N, Ketchakmadze K. *Staphylococcus aureus*: A major pathogen of food poisoning: A rare research report. Nutrition Food Process. 2022;5(1):1-3.
  47. Green BW. Effect of channel catfish stocking rate on yield and water quality in an intensive, mixed suspended-growth production system. North American Journal of Aquaculture. 2010;72(2):97-106.
  48. Mol S, Cosansu S, Alakavuk DU, Ozturan S. Survival of *Salmonella enteritidis* during salting and drying of horse mackerel (*Trachurus trachurus*) fillets. International Journal of Food Microbiology. 2010; 139:36–40.
  49. Olgunoğlu İA. *Salmonella* in Fish and Fishery Products. University of Adiyaman Vocational School of Kahta Turkey. Available:www.intechopen.com. ISBN 2012;978-953-307- 782-6.
  50. Loest D, Uhland FC, Young KM, Li X.Z, Mulvey MR, Reid-Smith R, Carson CA. Carbapenem-resistant *Escherichia coli* from shrimp and salmon available for purchase by consumers in Canada: A risk profile using the Codex framework. Epidemiology & Infection. 2022;150:e148.

51. Shen AQ. Multistate Outbreak of Salmonella Thompson Infections Linked to Seafood Exposure—United States, 2021. MMWR. Morbidity and Mortality Weekly Report. 2023;72.
52. Bauer RT. Symbioses. In Shrimps: Their Diversity, Intriguing Adaptations and Varied Lifestyles. Cham: Springer International Publishing. 2023;515-581.
53. Adeyemo OK, Foyle L. Public Health, Zoonoses, and Seafood, Safety. Fundamentals of Aquatic Veterinary Medicine. 2021;181-204.
54. Rohani MF, Islam SM, Hossain MK, Ferdous Z, Siddik MA, Nuruzzaman M, Shahjahan M. Probiotics, prebiotics and symbiotic improved the functionality of aquafeed: Upgrading growth, reproduction, immunity and disease resistance in fish. Fish & Shellfish Immunology. 2022;120: 569-589.

---

© 2024 Dumaloan-Canini et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/112884>