



# Genetic Analysis of Yield Traits in Snake Gourd (*Trichosanthes anguina* L.) Genotypes

M Jayanth Kumar <sup>a+++\*</sup>, G Kranthi Rekha <sup>a#</sup>,  
C Venkata Ramana <sup>b†</sup>, A Rajani <sup>c‡</sup>  
and D R Salomi Suneetha <sup>d^</sup>

<sup>a</sup> Department of Vegetable Science, College of Horticulture, Dr. YSR Horticultural University, Venkataramannagudem-534101, West Godavari, Andhra Pradesh, India.

<sup>b</sup> Horticultural Research Station, Dr. YSR Horticultural University, Lam, Guntur, Andhra Pradesh, India.

<sup>c</sup> Horticultural Polytechnic College, Nuzivedu, Krishna, Andhra Pradesh, India.

<sup>d</sup> Dr. YSR Horticultural University, Venkataramannagudem-534101, West Godavari, Andhra Pradesh, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/JEAI/2024/v46i52370

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114261>

Original Research Article

Received: 09/01/2024

Accepted: 13/03/2024

Published: 16/03/2024

## ABSTRACT

The present investigation was conducted to understand these parameters, which form an integral part of a programme for making improvements in snake gourd yield and its contributing characters. The present investigation was carried out with 32 snake gourd genotypes during late

<sup>++</sup> PhD Scholar;

<sup>#</sup> Assistant Professor (Horticulture);

<sup>†</sup> Senior Scientist (Horticulture);

<sup>‡</sup> Associate professor (Horticulture);

<sup>^</sup> Professor (Biochemistry), Dean of Student Affairs;

\*Corresponding author: E-mail: jayanth.m.babblu@gmail.com;

kharif 2021-22. There was need for more genetic research because the genotypes for every character under study differed significantly. Average fruit weight, fruit yield per vine, number of seeds per fruit, node to first female flower, fruit length, fiber content, acidity content, and potassium content all showed high values of PCV with correspondingly high values of GCV, indicating a greater degree of variability that could be attributed to genotype. Selection was very successful for these characters because of their high heritability and high genetic advance as measured by the percentage of mean for the following traits: fruit set percentage, fruit length, average fruit weight, flesh thickness, number of seeds per fruit, number of fruits per vine, fruit yield per vine, acidity content, and potassium content. Heritability was primarily caused by additive gene effect.

**Keywords:** Snake gourd; genetic advance; heritability; variability.

## 1. INTRODUCTION

“Snake gourd (*Trichosanthes anguina* L.)  $2n=2x=22$ ) is an annual, day-neutral, climbing herbaceous vegetable crop belonging to the Cucurbitaceae family, sub-family Cucurbitoideae, and tribe Trichosantheae. Originating in the Indo-Malayan region, it is believed to have evolved from wild species of *Trichosanthes*. Its distribution spans widely across Asia, from Malaya to North Australia, and extends through China and Japan. Presently, snake gourd cultivation also occurs in Mauritius, as well as in Central and East Java. In India, it is predominantly cultivated in South India and is also grown in regions including Punjab, Delhi, Uttar Pradesh, Bihar, Gujarat, and other parts of the country” [1].

“It serves as a valuable reservoir of minerals, fiber, and nutrients, contributing to the overall wholesomeness and healthfulness of food” [2]. “Notably, it contains significant proportions of protein (0.5%), fat (0.3%), minerals (0.5%), fiber (0.5%), and carbohydrates (3.3%). Its medicinal significance is attributed to its rich content of flavonoids, carotenoids, phenolic acids, and other compounds. Among its mineral constituents, potassium (121.6 mg/100 g), phosphorus (135 mg/100 g), sodium, magnesium, and zinc are prominently present” [3]. “The plant exhibits pharmacological and therapeutic properties, including anti-diabetic, hepatoprotective, cytotoxic, and anti-inflammatory effects, making it integral to Ayurveda and Siddha medical practices” [4].

Typically, snake gourd exhibits a higher yield per unit area, however, its average yield in India remains comparatively low compared to neighbouring countries, with production limited to a mere 3-4 months annually. Despite its economic and medicinal significance, there has been insufficient emphasis on implementing a

tailored crop improvement program, and little effort has been made to enhance the productivity and acceptance of this crop. Varietal improvement holds promise for significantly enhancing vegetable productivity. “The focus of improvement efforts should centre on selecting genotypes that offer improved yield and superior quality. Snake gourd, being a monoecious and highly cross-pollinated crop, boasts numerous cultivars showcasing extensive variation in fruit size, shape, and colour within India. Beyond its nutritional value, there exists a substantial market demand for snake gourd. Given its nutritional and medicinal significance, there is a critical need for its improvement. The identification of an improved variety boasting high yield and superior quality traits, coupled with broader adaptability, would greatly benefit farmers” [5].

“Understanding the extent of variability within a population is crucial as it forms the foundation for successful selection processes. A comprehensive grasp of genetic variability and its components is essential for planning breeding programs. Typically, the genotypic coefficient of variability (GCV) and phenotypic coefficient of variability (PCV) are assessed to analyse variability. Given that population variability results from both genotypic and environmental influences, understanding the nature and extent of genetic variation contributing to gains under selection is vital” (Singh *et al.*, 1986).

The success of improvement of characters through selection depends on the heritability coupled with its genetic advance. Once the relative amount of variability in population is assessed, it becomes necessary to partition the overall variability into heritable and non-heritable components. Magnitude of heritability indicates the effectiveness with which selection of genotypes can be based on phenotypic performance. Burton [6] suggested that

heritability along with GCV would provide a clear idea about the amount of genetic advance expected through selection. High value of heritability indicates that phenotype of the trait strongly reflects the genotype and suggests the major role of genotypic constitution in the expression of the character. Such traits are considered dependable from breeding point of view. Therefore, the present investigation was conducted to understand these parameters, which form an integral part of a programme for making improvements in snake gourd yield and its contributing characters.

## 2. MATERIALS AND METHODS

The experiment was conducted at College of Horticulture, Dr. Y.S.R. Horticultural University, Venkataramannagudem, West Godavari District. It was conducted during late *kharif* season 2021-22 and laid out in Randomized Block design replicated twice. Total thirty-two snake gourd genotypes were evaluated for growth yield and quality traits. Genotypes were collected from NBPGR regional station Thrissur. "The experimental site was well prepared, cultural practices include training, pruning, weeding, irrigation, fertilizer application and plant protection measures were followed for the healthy growth of crop" [7]. "Observations on growth parameters were recorded upto 4 months of planting. Data on yield and yield attributes were collected at appropriate stages" [8].

## 3. RESULTS AND DISCUSSION

The analysis of variance showed that the genotypes differed significantly among themselves for all the characters indicating the presence of adequate variability (Table 1). In the present investigation, phenotypic coefficient of variation was higher than corresponding genotypic coefficient of variation indicating the influence of environmental factors in the expression of these characters [9]. Results were estimated and presented in Table 2.

High PCV and GCV values were recorded for average fruit weight (32.34%, 30.88%), fruit yield per vine (42.24%, 40.54%), number of seeds per fruit (43.33%, 42.94%), node to first female flower (27.24%, 21.94%), fruit length (27.66%, 25.88%), fibre content (36.82%, 28.59%), acidity content (52.29%, 51.41%) and potassium content (27.96%, 27.90%) in all genotypes. This indicating the existence of more variability among all the traits recorded and ample scope for improvement of these characters through

selection. These results are in confirmation with the findings of Rahman et al. [10], Padmaraja [1] and Rana and Pandit [9] in snake gourd, Choudhary et al. [11] and Sravani et al. [12] in ridge gourd, Akter et al. [13] in pumpkin and Chakraborty et al. [14] in bitter gourd.

High PCV and moderate GCV were observed for node to first male flower (22.14%, 14.14%), flesh thickness (22.56%, 18.35%) and number of fruits per vine (21.75%, 19.91%). Moderate PCV and moderate GCV were observed for days to first male flower opening (14.26%, 13.66%), days to first female flower opening (17.15%, 16.53%), fruit set percentage (17.24%, 15.41%), vitamin C content (12.73%, 10.69%) and TSS (12.47%, 10.48%) indicating the existence of variability. These results are confirmation with the findings of Rana and Pandit [9] and Deepa and Mariappan [15] in snake gourd, Choudhary et al. [11] in ridge gourd, Tomar et al. [16] and Reddy et al. [17] in muskmelon and Chakraborty et al. [14] in bitter gourd.

Moderate PCV and low GCV were recorded for sex ratio (11.22%, 7.14%) and fruit girth (11.95%, 8.85%). Similar results are reported by Sundaram et al. [18] in watermelon and Kumar et al. [19] in cucumber. Low PCV and GCV for number of male flowers per vine (1.04%, 0.87%) and number of female flowers per vine (9.70%, 7.80%). This indicated the low variability for these characters which is the constraint for genetic improvement through selection. These results are contradictory to those reported by Rana and Pandit [9] in snake gourd and Karthick et al. [20] in cucumber where the values of PCV (31.27%) and GCV (22.86 %) were recorded to be high.

However, the PCV recorded in the present experiment was higher than GCV for all the characters, though closeness between PCV and GCV were recorded for some characters. These results show that, there is an abundant scope of improvement through selection as it has been indicated that phenotypic expression of all genotypes is mostly under genetic control and environment has slight to moderate influence which are in conformity with the finding were reported by Rana and Pandit [9].

In the present experiment high heritability coupled with high genetic advance as per cent of mean was recorded in days to first male flower opening (91.00%, 26.95%), days to first female flower opening (92.00%, 32.83%), node to first

**Table 1. Analysis of variance for different characters in 32 snake gourd genotypes**

S. No	Characters	Mean sum of squares		
		Replications Df= 1	Treatments Df= 31	Error Df= 31
1	Days to first male flower opening	26.21	1788.58**	77.27
2	Days to first female flower opening	66.54	6377.18**	234.35
3	Node to first male flower	1.24	51.69*	21.73
4	Node to first female flower	9.09	431.97**	92.00
5	Number of male flowers per vine	0.76	1069.65**	193.25
6	Number of female flowers per vine	0.52	340.33**	72.70
7	Sex ratio (%)	0.95	147.72*	62.62
8	Fruit set (%)	10.92	9151.98**	1019.70
9	Fruit length (cm)	75.97	7469.11**	494.41
10	Fruit girth (cm)	0.27	127.50**	37.15
11	Average fruit weight (g)	9713.08	465958.19**	21489.76
12	Flesh thickness (cm)	0.03	0.54**	0.11
13	Number of seeds per fruit	5.64	18580.25**	169.10
14	Number of fruits per vine	4.17	1044.15**	91.67
15	Fruit yield per vine (kg)	1.06	306.06**	12.56
16	Vitamin C content (mg/100 g)	0.03	14.36**	2.48
17	Acidity content (per cent)	0.0003	0.42**	0.007
18	Potassium (mg/100 g)	56.25	243215.93**	466.75
19	TSS (°Brix)	0.03	9.40**	1.61
20	Fibre content (g/100 g)	0.0006	1.29**	0.31

\*Significant at 5 % level of significance \*\* Significant at 1 % level of significance

**Table 2. Estimates of genetic parameters in snake gourd genotypes**

<b>Character</b>	<b>Range</b>	<b>Mean</b>	<b>GCV</b>	<b>PCV</b>	<b>h<sup>2</sup></b>	<b>GA</b>	<b>GAM (%)</b>
Days to first male flower opening	28.43-46.63	38.44	13.66	14.26	91.00	10.36	26.95
Days to first female flower opening	44.30-77.63	60.19	16.53	17.15	92.00	19.76	32.83
Node to first male flower	3.00-7.00	4.91	14.14	22.14	40.00	0.91	18.61
Node to first female flower	7.25-14.67	10.67	21.94	27.24	64.00	3.88	36.42
Number of male flowers per vine	423.90-443.13	431.80	0.87	1.04	69.00	6.45	1.49
Number of female flowers per vine	24.00-34.18	26.61	7.80	9.70	64.00	3.44	12.94
Sex ratio (%)	12.38-19.10	16.41	7.14	11.22	40.00	1.53	9.35
Fruit set (%)	51.50-95.83	74.29	15.41	17.24	79.00	21.09	28.39
Fruit length (cm)	26.40-78.70	40.97	25.88	27.66	87.00	20.44	49.90
Fruit girth (cm)	10.93-17.15	13.62	8.85	11.95	54.00	1.84	13.51
Average fruit weight (g)	141.25-459.50	274.12	30.88	32.34	91.00	166.55	60.75
Flesh thickness (cm)	0.30-0.70	0.46	18.35	22.56	66.00	0.14	30.76
Number of seeds per fruit	14.00-78.25	40.13	42.94	43.33	98.00	35.17	87.66
Number of fruits per vine	13.48-32.65	19.68	19.91	21.75	83.00	7.39	37.57
Fruit yield per vine (kg)	2.82-13.11	5.37	40.54	42.24	92.00	4.30	80.15
Vitamin C content (mg/100 g)	3.35-4.89	4.09	10.69	12.73	70.00	0.75	18.49
Acidity content (per cent)	0.11-0.18	0.16	51.41	52.29	96.00	0.16	104.11
Potassium (mg/100 g)	120.50-350.50	224.22	27.90	27.96	99.00	128.65	57.37
TSS (°Brix)	2.58-4.15	3.38	10.48	12.47	70.00	0.61	18.16
Fibre content (g/100 g)	0.15-0.75	0.44	28.59	36.82	60.00	0.20	45.75

**GCV** Genotypic coefficients of variation **PCV** Phenotypic coefficients of variation **h<sup>2</sup>** Heritability  
**GA** Genetic advance **GAM (%)** Genetic advance as per cent of mean

female flower (64.00%, 36.42%), fruit set percentage (79.00%, 28.39%), fruit length (87.00%, 49.90%), average fruit weight (91.00%, 60.75%), flesh thickness (66.00%, 30.76%), number of seeds per fruit (98.00%, 87.66%), number of fruits per vine (83.00%, 37.57%), fruit yield per vine (92.00%, 80.15%), acidity content (96.00%, 104.11%) and potassium content (99.00%, 57.37%) indicating heritability was mainly due to additive gene effect and hence selection was highly effective for these characters. These results are in line with the findings of and Rahman et al. [10], Rana and Pandit [9] and Deepa and Mariappan [15] in snake gourd, Choudhary et al. [11] in ridge gourd, Chakraborty et al. [14] in bitter gourd, Reddy et al. [17] in muskmelon and Shah et al. [21] in cucumber.

According to Panse and Sukhatme [22], "the characters with high heritability coupled with high genetic advance as per cent of mean were controlled by additive gene action and therefore amenable to improvement through selection. So, the selection of phenotypically superior plants with respect to the discussed characters will result in significant improvement in the next generation".

High heritability along with moderate genetic advance was observed for number of female flowers per vine (64.00%, 12.94%), vitamin C content (70.00%, 18.49%) and TSS (70.00%, 18.16%) indicated the presence of additive gene action and selection may be effective for this character. High heritability along with low genetic advance was observed for number of male flowers per vine (69.00%, 1.49%). Similar results were obtained by Rana and Pandit [9] in snake gourd and Reddy et al. [17] in muskmelon.

Moderate heritability and high genetic advance were observed for fibre content (60.00%, 45.75%) indicated that these character was influenced by environmental factors. Moderate heritability and moderate genetic advance were observed for node to first male flower (40.00%, 18.61%) and fruit girth (54.00%, 13.51%). These results are in conformity with the findings of Rana and Pandit [9] in snake gourd, Veena et al. [23] in cucumber and Pathak et al. [24] in bitter gourd. The trait sex ratio (40.00%, 9.35%) exhibited moderate heritability and low genetic advance which indicates that it was influenced by environment and governed by non-additive gene

action. Similar results were reported by Sundaram et al. [18] in watermelon.

#### 4. CONCLUSION

Significant values of PCV along with corresponding high GCV were observed across various parameters including average fruit weight, fruit yield per vine, number of seeds per fruit, node to first female flower, fruit length, fiber content, acidity content, and potassium content. These findings suggest a substantial degree of variability attributable to genotype, indicating that these traits are less influenced by environmental factors and are amenable to improvement through selective breeding. In particular, efforts to enhance fruit yield per vine in snake gourd should focus on traits such as days to first male and female flower opening, node to first female flower, fruit set percentage, fruit length, average fruit weight, flesh thickness, number of seeds per fruit, number of fruits per vine, acidity content, and potassium content, as these traits exhibit high heritability and substantial genetic advance as a percentage of the mean. Breeding programs should aim to leverage both additive and non-additive gene effects, employing suitable techniques to facilitate varietal development and contribute to the advancement of snake gourd breeding initiatives.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Padmaraja SR. Genetic variability studies in snake gourd (*Trichosanthes cucurmarina* L.) M.Sc. (Hort.) Thesis. University of Agricultural Sciences, Bengaluru (India); 2011.
2. Ahmed MS, Rasul MG, Bashir MK, Mian ASM. Variability and heterosis in snake gourd (*Trichosanthes anguina* L.). Bangladesh Journal of Plant Breeding and Genetics. 2000;13(2):27-32.
3. Ojiako OA, Igwe CU. The nutritive, anti-nutritive and hepatotoxic properties of (*Trichosanthes anguina* L.) (snake tomato) fruits from Nigeria. Pakistan Journal of Nutrition. 2008;7(1):85–89.
4. Warrior PK, Nambiar VPK, Ramankutty C. Indian medicinal plants compendium of

- 500 species. Orient Longman private limited Chennai. 1993:320-22.
5. Sivabodh B. Divergence and character association studies in snake gourd (*Trichosanthes cucumerina* L.) M.Sc. (Hort.) Thesis. University of horticultural sciences, Bagalkot (India); 2018.
  6. Burton GW, Devane EM. Estimating heritability from replicated clonal material. *Agronomy Journals*. 1952;45:478-81.
  7. Thamburaj S, Singh N. Vegetables, tubercrops and spices. Indian Council of Agricultural Research, Krishi Anusandhan Bhawan, Pusa, New Delhi; 2004.
  8. Kumar MJ, Rekha GK, Ramana CV, Rajani A, Suneetha DS. Performance of snake gourd (*Trichosanthes anguina* L.) genotypes for growth, yield and quality traits in coastal Andhra Pradesh. *The Pharma Innovation*. 2022;11(8):232-5.
  9. Rana NP, Pandit MK. Studies on genetic variability, character association and path analysis in snake gourd (*Trichosanthes anguina* L.) genotypes. *Journal of Crop and Weed*. 2011;7(2):91-96.
  10. Rahman MA, Hossain MD, Islam MS, Biswas DK, Ahiduzzaman M. Genetic variability, heritability and path analysis in snake gourd (*Trichosanthes anguina* L.). *Pakistan Journal of Biological Sciences*. 2002;5(3):284-86.
  11. Choudhary BR, Fageria MS, Dhaka RS. Correlation and path coefficient analysis in muskmelon (*Cucumis melo* L.). *Indian Journal of Horticulture*. 2011;61(2): 258-62.
  12. Sravani Y, Kranthi RG, Venkata RC, Naram NL, Salomi SDR. Studies on genetic variability, heritability and genetic advance in F<sub>2</sub> generation of ridge gourd. *The Pharma Innovation Journal*. 2021;10(7):927-30.
  13. Akter S, Rasul MG, Aminul Islam AKM, Hossain MM. Genetic variability, correlation and path coefficient analysis of yield and quality traits in pumpkin (*Cucurbita moschata*). *Bangladesh Journal of Plant Breeding and Genetics*. 2013; 26(1):25-33.
  14. Chakraborty L, Acharya Raychaudari S. Diversity analysis of *Momordica charantia* L. accessions from Eastern and Northeastern India based on morphological, yield related traits and molecular marker. *Proceedings of International Symposium on Quality Management of Fruits and Vegetables for Human Health*. 2013; 179-93.
  15. Deepa D, Mariappan S. Genetic variability, heritability and genetic advance for yield and its components snake gourd (*Trichosanthes anguina* L.). *African Journal of Agricultural Resource*. 2013;8(42): 5221-25.
  16. Tomar RS, Kulkarni GU, Kakade DK. Genetic analysis in Muskmelon. *Journal of Horticultural Science*. 2008; 3(2): 112-18.
  17. Reddy BK, Begum H, Sunil N, Reddy MT. Variance component analysis of quantitative traits in musk melon (*Cucumis melo* L.). *Trakia Journal of Science*. 2013; 2:118-24.
  18. Sundaram MS, Kanthaswamy V, Ashok KG. Studies on variability, heritability, genetic advance and character association in watermelon (*Citrullus lunatus* (Thunb.) Matsam and Nakai). *Progressive Horticulture*. 2011;43(1):20-24.
  19. Kumar R, Ameta KD, Dubey RB, Sunil P. Genetic variability, correlation and path analysis in sponge gourd (*Luffa cylindrica*). *African Journal of Biotechnology*. 2013; 12(6):539-43.
  20. Karthick K, Arumugam T, Rajasree V, Ganesan KN, Karthikeyan M. Evaluation and assessment of genetic variability of cucumber (*Cucumis sativus* L.) genotypes. *The Pharma Innovation Journal*. 2019; 8(11):156-60.
  21. Shah KN, Rana DK, Singh V. Genetic evaluation of cucumber (*Cucumis sativus* L.) strains for different growth yield, quality and seed parameters. *International Journal of Agricultural Invention*. 2017; 2(2):130-35.
  22. Panse VG, Sukhatme PV. *Statistical Methods for Agricultural Workers*. 2nd Edition ICAR, New Delhi. 1967;361.
  23. Veena R, Amrik SS, Pitchaimuthu M, Souravi K. Genetic evaluation of Cucumber (*Cucumis sativus* L.) genotypes for some yield and related traits. *Electronic Journal of Plant Breeding*. 2012; 3(3): 945-48.

24. Pathak M, Manpreet, Pahwa, K. Genetic variability, correlation and path coefficient analysis in bittergourd (*Momordica charantia* L.). International Journal of Agricultural Research. 2014; 2(8): 179-84.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/114261>*