



Genomic Evolution and Dynamics of Drug Resistance in *Mycobacterium Tuberculosis* across West Africa- A Review

**Oluwabusola Oluwakorede Asenuga ^a,
James Ponman Sargwak ^b, Emmanuel Chimeh Ezeako ^c,
Matthew Babalola ^d, Faith Omosigho ^e,
Eberechukwu Osinachi Azubuiké ^f
and Chibuzo Valentine Nwokafor ^{f,g*}**

^a Department of Zoology and Environmental Biology, Lagos State University, Ojo, Lagos State, Nigeria.

^b Department of Biochemistry, Plateau State University, Bokkos, Plateau State, Nigeria.

^c Department of Biochemistry, University of Nigeria, Nsukka, Enugu State, Nigeria.

^d Department of Animal Science, University of Abuja, Abuja, Nigeria.

^e Department of Pharmacology and Therapeutics, Delta State University, Abraka, Nigeria.

^f Department of Microbiology, Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria.

^g Department of Biotechnology, University of the West of Scotland, Scotland, United Kingdom.

Authors' contributions

This work was carried out in collaboration among all authors. The final manuscript was written, edited, read and approved by all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/mrji/2024/v34i111504>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124356>

*Corresponding author: E-mail: chibuzonwokafor@gmail.com;

ABSTRACT

Background: The continuous evolution of drug-resistant Tuberculosis strains around the globe, particularly in West Africa, continues to be a major public health problem and poses serious threats to the actualization of the "End Tuberculosis Strategy" projected by 2030. Hence, a comprehensive knowledge on genetic variations, lineage distributions and evolutionary adaptations implicated in *M. tuberculosis* drug-resistance could strengthen research efforts in TB control.

Objectives: This review summarizes the genetic mechanisms of drug resistance in *Mycobacterium tuberculosis* and the extent to which they pose threats to public health. It also gives recommendations on managing drug resistant TB in West Africa.

Literature Search: This review assesses available literature relating to the genomic evolution and dynamics of drug resistance in *Mycobacterium tuberculosis* across west Africa. Search for relevant articles in databases such as Google Scholar, PubMed, Scopus and Web of Science was conducted using keywords such as "Genomics", "Evolution", "Drug Resistance", "*Mycobacterium tuberculosis*" and "West Africa ". Sixty articles were found in this search and forty-six articles were selected and reviewed.

Results: Multidrug-resistant tuberculosis (MDR-TB) and extensively drug-resistant tuberculosis (XDR-TB), which are common in West Africa, have been linked to the MTBC Lineages 5 and 6 also known as *M. africanum*. The devastating challenges these resistant strains exert on public health justified the urgency for exploring novel therapeutic avenues, improved diagnostic protocol, and robust healthcare systems to curb the disease.

Conclusion: This review maintained a strong advocacy for proper public health education, installation of adequate surveillance systems, and the adoption of alternative therapeutic modalities to tackle drug-resistant *M. tuberculosis*, effectively in West Africa and mitigate the public health burden it poses globally.

Keywords: Genomics; evolution; drug resistance; *Mycobacterium tuberculosis*; West Africa.

1. INTRODUCTION

One of the most difficult and oldest known infectious diseases still in existence, with a history dating back up to a century and a half, is tuberculosis (TB), which is caused by the *Mycobacterium tuberculosis* (Barberis et al. 2017). As a communicable disease TB pose a severe threat to public health and represent major concerns that unevenly affect and decimate vulnerable populations of developing countries (Wu et al. 2023). Vulnerable populations are at alarmingly high risk for TB owing to their marginalized or disadvantaged socioeconomic status, which limit their access to adequate healthcare facilities. Although substantial scientific efforts have been leveraged in the combat of TB, the disease continues to rank among the top ten worldwide causes of death (Litvinjenko et al. 2023). In 2022, 10.6 million new cases of TB and 1.6 million deaths

related to the disease were reported, according to a recent World Health Organization (WHO) report (WHO 2023). This puts TB as the 13th most deleterious disease worldwide, ahead of both HIV and AIDS (WHO 2023). Africa, with 15.19% of the world's population, accounts for approximately 24% of the worldwide tuberculosis (TB) burden (Otchere et al. 2024). West Africa is home to all six main MTBC lineages, in contrast with other regions that only have one or a few of these lineages (Otchere et al. 2024). Furthermore, the global burden of TB is unevenly distributed, with developing countries and middle-income regions particularly in Asia and Africa having the highest burden. In fact, socioeconomic and health-related issues, including HIV co-infection and inadequate healthcare systems, as well as poverty and hunger contribute to this inequality (Bai et al. 2024). In addition, the recent appearance and reemergence of different types of tuberculosis,

such as extensively drug-resistant TB (XDR-TB) and multidrug-resistant TB (MDR-TB), which continue to counter scientific efforts to curtail the disease, presenting a major threat to public health (Mancuso et al. 2023).

Alarming high rates of tuberculosis cases are reported in West Africa, where together Nigeria, Ghana, Liberia, Senegal, and Mali account for a sizable share of the world's tuberculosis cases (Adebisi et al. 2019). Notably, the entire West Africa region is faced with substantial challenges in controlling the disease, however, the rate of TB incidence in this region differs across countries. Socioeconomic determinants, like poverty, migration, inadequate reach to healthcare infrastructures, poor disease diagnosis and treatment, particularly in rural areas contribute to a large extent to the spread of TB (Msoka et al. 2021). Delayed diagnosis and stigma associated with patients constitute the impediments truncating efforts to control TB. Moreover, the high rates of HIV recorded in this region worsens the TB epidemic, and as a co-infection, HIV distorts immune functions and increases risk of TB (Kraef et al. 2021).

More worrisome is the alarming rates of the prevalence of multidrug resistant strains of the bacteria in West Africa, where inadequate diagnostics and surveillance systems often translates to underreporting of cases. Besides, the porosity of the region's borders and inadequate control of cross-border migrations also facilitate TB transmission, including drug-resistant strains across neighboring countries (Otchere et al. 2024). The majority of drug resistant tuberculosis cases are caused by human intervention. The first documented case of drug resistance in *M. tuberculosis* was linked to the use of streptomycin (S) shortly after its discovery. Combined resistance of Streptomycin with other drugs such as isoniazid, pyrazinamide, ethambutol and rifampin also emerged (Islam et al. 2017).

In order to stop the worldwide tuberculosis (TB) epidemic, the World Health Organization (WHO) has implemented the "End TB Strategy", which would lower global TB incidence and mortality rates by 90% and 95%, respectively, by 2035 compared to 2015 (WHO 2023).

The expected number of new TB cases and deaths has continued to fall, a trend that started in the African Region in 2005. TB and TB/HIV therapies are expected to have prevented 16 million deaths in the region between 2000 and

2021. With a 22% decrease in new cases from 2015 to 2021, the African Region of the "End TB Strategy" achieved its 2020 milestone. Additionally, eight nations have achieved a 35% decrease from 2015 (WHO 2023).

In this review, we discussed the mechanisms of drug resistance in *Mycobacterium tuberculosis*, along with the challenges and implications of the disease for public health. We also highlighted strategies for detecting and controlling drug-resistant tuberculosis in West Africa.

2. LITERATURE SEARCH

This review assesses available literature relating to the genomic evolution and dynamics of drug resistance in *Mycobacterium tuberculosis* across the west Africa region. Search for relevant articles in databases such as Google Scholar, PubMed, Scopus and Web of Science was conducted using keywords such as "Genomics", "Evolution", "Drug Resistance", "*Mycobacterium tuberculosis*" and "West Africa". Paper selections were gotten by reviewing abstracts and titles. Sixty articles were found in this search and forty six articles, including reports were selected and reviewed. The review discussed studies from 2014 to present.

2.1 Genetic Diversity of *Mycobacterium tuberculosis* in West Africa

Mycobacterium tuberculosis (MTB) is the causative agent of tuberculosis in the lungs of humans. It is composed of seven distinct lineages that are related to geographical regions based on phylogeny (Osei-Wusu et al. 2022). Lineages one through four (1-4) and seven (7) are generally known as *Mycobacterium tuberculosis sensu stricto*, and are widely distributed. On the other hand, lineages five and six (5 and 6), *Mycobacterium africanum* is exclusive to West Africa (Asante-Poku et al. 2015).

Genomic diversity of MTB in West Africa is characterized by the prevalence of lineages 5 and 6 (Coscolla 2014). The lineage five (5) is most common in the eastern part of West Africa, which encompasses Nigeria, Cameroon, the Benin Republic, and Ghana (Gehre et al. 2016) while the western areas in West Africa, which includes Ghana, Senegal Guinea Bissau, Sierra Leone, and Gambia, Lineage six (6) is more prevalent (De et al. 2010). *M. africanum* is usually known for its slow progression of the

tuberculosis disease compared to the progression in other lineages. It has also been established that areas with higher prevalence of *M. africanum* will experience different transmission dynamics and higher disease severity than regions with other lineages (Coscolla 2014). Evolutionary dynamics of tuberculosis in these regions are usually influenced by different factors, including environmental pressures and genetic mutations (Gagneux 2018). The evolution of tuberculosis is driven by genetic mutations that contribute to its adaptation. Mutations linked to drug resistance in tuberculosis have been found in genes such as *katG*, *inhA* and *rpoB* (Bentley et al. 2012).

2.2 Genetic Mechanisms of Drug Resistance

There are different types of drug-resistant tuberculosis disease.

Resistance to just one tuberculosis treatment drug is known as Mono-resistant TB disease. Poly-resistant TB disease is caused by tuberculosis bacteria that are resistant to at least two TB drugs (other than isoniazid and rifampin) (Bentley et al. 2015).

The resistance to more than one drug is defined as Multi-drug resistance and in tuberculosis, it is explained as the minimum resistance to both isoniazid and rifampin (Hamed et al. 2021).

Pre-extensively drug-resistant TB (pre-XDR TB) disease is a type of multidrug resistant

tuberculosis caused by TB bacteria that are resistant to isoniazid, rifampin (MDR-TB), a fluoroquinolone and at least one of the three second-line injectables (amikacin, capreomycin, and kanamycin) (Seung et al. 2015).

Rifampicin resistance (RR) is the resistance to rifampicin detected using phenotypic or genotypic methods, with or without resistance to other anti-TB drugs. It includes any resistance to rifampicin, in the form of mono-resistance, poly-resistance, MDR or XDR (Seung et al. 2015).

Mycobacterium tuberculosis resistance is caused by genetic mutations which are usually single nucleotide polymorphisms (SNPs), small deletions or insertions, and in a few instances larger deletions or inversions. These mutations can appear suddenly, usually spread by replication inside of the host and subsequent transfer between bacteria resistant hosts (Nimmo et al. 2022). Three primary ways by which *Mycobacterium tuberculosis* can develop resistance to drugs include efflux pump modulations, enzyme inactivation and target-based mutations. Efflux pumps expel drugs from bacterial cells reducing drug concentrations and contribute to resistance while target-based mutations are scenarios in which the drug target undergoes mutation preventing binding of drugs (Nimmo et al. 2022). Other drug resistance mechanisms include, prevention of drug penetration, target overproduction, target mimicry (Nina et al. 2006).




Fig. 1. A Cartogram describing the global TB burden using *M. tuberculosis* lineage (Koleske et al. 2023)




Fig. 2. Methods of drug resistance mechanisms (Nina et al. 2006)




Fig. 3 The 2016 WHO estimates of multidrug resistant tuberculosis compared with reports from nine West African countries (WHO 2023)

2.3 Mechanisms of First Line Drug Resistance

Isoniazid: When treating tuberculosis, isoniazid is frequently used as a medication. The catalase-oxidase enzyme, which is encoded by the *katG* gene, activates it. Isoniazid resistance is caused by the most prevalent mutation, the *katG* mutation, which primarily occurs at codon 315. Further mutations that result in resistance can occur within the *inhA* gene and promoter region (Dookie et al. 2018). 7.1% and 7.9% of newly

diagnosed and previously treated TB patients were identified worldwide, according to global data on isoniazid resistance (World Health Organization 2018).

Rifampicin: Rifampicin, another first-line drug, RNA polymerase encoded by *rPOB* gene which reduces the rifampicin binding and leads to bacterial resistance. The codons 516 and 529 are the sites of the most frequent mutations (Dookie et al. 2018) saw a global prevalence of rifampicin resistance in 3.4% of newly diagnosed

cases and 18% of patients of tuberculosis that have already received treatment (Dean et al. 2020).

The frequency of multidrug resistance (MDR) was examined in isolates from eight West African nations: Gambia, Nigeria, Mali, Togo, Ghana, Guinea-Bissau, Senegal and Burkina-Faso. The study was conducted by Gehre et al. 2016. According to the study, 39% of the patients who underwent testing had firstline medication resistance. Nigeria has the greatest prevalence, with 30% in Ibadan and 59% in Lagos. Ghana followed closely with MDR prevalence of 26%. The percentages of MDR prevalence in Guinea Bissau and Gambia were low at 7%. In previously treated cases, Mali's prevalence was at 59% while Nigeria (Lagos and Ibadan) had 66% and 39% respectively. Multidrug resistance TB cases were found to be more prevalent in these three locations. With 13%, Gambia had the lowest.

2.4 Geographic Distribution and Epidemiology of Drug-resistant TB in West Africa

Very little information is known about drug resistant tuberculosis in West Africa (Saleri et al. 2010). A rising public health challenge for many West African Countries is drug-resistant tuberculosis, which consists of multidrug resistant and extensively drug resistant strains with varying frequency over time (World Health Organization 2023, N'Guessan et al. 2018). According to WANETAM (West African Network for Tuberculosis, AIDS and Malaria), collaborative surveillance on drug-resistant tuberculosis in nine West African countries was reported. Fig. 3 shows the estimates provided by the World Health Organization (WHO) often fall short of the actual incidence (World Health Organization 2023). For instance, the study conducted by WANETAM reported that 39% of patients were resistant to at least one specific tuberculosis drug, but the WHO indicated that the prevalence of multidrug resistant tuberculosis was 2% in new cases and 17% in cases that had previously received treatment. Highly noted was the presence of pre-extensive drug resistant tuberculosis cases in all the eight countries that participated (Gehre et al. 2016). The World Health Organization recently released data on tuberculosis cases which showed that the prevalence of multidrug resistance/rifampicin resistant tuberculosis varied widely among West

African nations, ranging from 54% in Guinea-Conakry to 0.67% in Togo. This is summarized in Fig. 4 (World Health Organization 2023).

2.5 Public Health Implications and Challenges

To effectively treat drug-resistant tuberculosis, there is a need for correct and timely diagnosis. It is possible to detect diseases using a variety of diagnostic techniques that combine both DNA and culture-based (phenotypic) testing (Yusoof et al. 2022). The disadvantages of drug susceptibility testing (DST) procedures based on culture include the need for specialized laboratories, a high level of expertise, and the fact that results can take weeks to obtain. In contrast, DNA based techniques, like polymerase chain reaction (PCR), give a more rapid alternative for testing (Boehme et al. 2010). Treating drug resistant tuberculosis is challenging as it demands longer use of drugs lasting between 6 to 9 months. This long period can lead to several difficulties. Firstly, side effects such as liver and gastrointestinal problems may develop as a result of combining drugs and cause treatment failures (Yang et al. 2017). Though the treatment rate was among the highest in West Africa, it still fell below the WHO recommended target of 90% registration. Limited access to these essential medications can cause inadequate treatment plans, which may undermine treatment results and lead to the progression of drug resistance further (World Health Organization 2020).

2.6 Managing Drug Resistant TB in West Africa Public Health Sector

In West Africa, managing DR-TB has proven to be a complex issue that needs careful solution. Firstly, the inadequate access diagnosis which then causes delayed diagnosis and less effective therapy needs to be tendered to and controlled (African Union 2019). Secondly, the delayed diagnosis of this drug-resistant TB then results in an extended infectious period which then increases the risk of further transmission to others that did not have the disease (Atre et al. 2022). This solidifies the need for prompt diagnosis and appropriate treatment as these are very important for improving treatment results and stopping transmission (Kaaffah et al. 2023).




Fig. 4. The global tuberculosis report on the estimates of multidrug resistant/rifampicin resistant tuberculosis among newly diagnosed and recurrent cases of tuberculosis in West Africa as of 2022 (World Health Organization 2023)

3. FUTURE DIRECTIONS AND RECOMMENDATIONS

Early detection of DR-TB is dependent on surveillance systems, and it is necessary that they should be improved. By improving better surveillance systems, healthcare professionals will be able to identify and respond to cases of emerging resistance faster (Stop TB Partnership 2022). The Government also has a role to play in improving health outcomes. Governments in countries in West Africa should reduce their dependence on foreign aid and should focus on building healthcare systems in the country to be more responsible and resilient (Global Fund 2021, Nathanson et al. 2010).

Contact tracing and monitoring of individuals that have been exposed to DR-TB patients should be effective. This, in addition with regular testing of patients will facilitate the detection of new cases early, it will prevent secondary transmission and improve treatment outcomes in the country (Velen et al. 2021, Koura et al. 2022).

There should be more research on TB in West Africa to advance our understanding and also aid in the creation of more effective control measures. There are several medicinal plants in West Africa, for treating different diseases

including tuberculosis. Investing in this plant-based field is necessary as it contains promising potential leads for new anti-tuberculosis drugs (Nguta et al. 2016). Plant species such as *Zanthoxylum leprieurii*, *Lantana camara*, *Cryptolepis sanguinolenta* have been reported to exhibit antimycobacterial action against drug-resistant strains of *Mycobacterium* TB (uyiringire et al. 2022).

Finally, educating the public is the first and most important step in addressing the growing threat of drug-resistant TB in these regions. Even healthcare workers are not excluded in education as a good number of them are ignorant on causes, symptoms and management of drug-resistant tuberculosis. Targeted educational programs can help address misconceptions about the disease and increase early diagnosis and treatment (Hassan et al. 2017). A combination of different communication platforms such as social media, mass media, print materials e.t.c should be used to reach a wide audience with consistent and accurate information (World Health Organization 2020).

4. CONCLUSION

This review demonstrated that the genomic diversity and emergence of DR-TB in *Mycobacterium tuberculosis* in West Africa

instigate tremendous threats to public health and contributes to the alarmingly high rate of tuberculosis cases in this region, caused by *M. africanum* lineages (5 and 6). The advent of MDR-TB strains endangers public health, necessitating the active search for innovative therapeutic modalities against drug-resistant tuberculosis, as well as the implementation of enhanced diagnostic paradigm, effective treatment regimen, and resilient healthcare facilities. To address the several challenges of drug-resistance in TB, we recommend the design of multidimensional and comprehensive strategies for the propagation of public health education against disease management, enhancement of surveillance systems, and the adoption of alternative remedies such as phytotherapeutics against TB and DR-TB. It also behooves the collaboration of governments agencies and other relevant stakeholders to strengthen the healthcare systems and promote timely and proper TB diagnosis, prevention and management, to minimize the spread and lethal impacts of tuberculosis in the developing countries. In addition, appropriating public funds to improve public health education programs and the installation of adequate healthcare facilities in rural and remote areas is quintessential to improving TB control and realizing better health outcomes among vulnerable populations in West Africa.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

ACKNOWLEDGEMENTS

The Authors wish to thank Chijamz Academy's Collaborative Academic Research Team (CART) for their help and support toward the preparation and drafting of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

Adebisi, Y. A., Agumage, I., Sylvanus, T. D., Nawaila, I. J., Ekwere, W. A., et al. (2019). Burden of tuberculosis and challenges

- facing its eradication in West Africa. *International Journal of Infectious Diseases*, 6(3), e92250. <https://doi.org/10.5812/iji.92250>
- African Union. (2019). *Africa Continental 2019 Africa Continental TB Scorecard*. WHO Press. <https://www.africaunion.org/en/africa-tb-scorecard>
- Asante-Poku, A., Yeboah-Manu, D., Otchere, I. D., Aboagye, S. Y., Stucki, D., Hattendorf, J., Borrell, S., Feldmann, J., Danso, E., Gagneux, S. (2015). Mycobacterium africanum is associated with patient ethnicity in Ghana. *PLoS Neglected Tropical Diseases*, 9(1), e3370.
- Atre, S. R., Jagtap, J. D., Faqih, M. I., Dumbare, Y. K., Sawant, T. U., Ambike, S. L., et al. (2022). Tuberculosis pathways to care and transmission of multidrug resistance in India. *American Journal of Respiratory and Critical Care Medicine*, 205, 233-241. <https://doi.org/10.1164/rccm.202012-4333OC>
- Bai, W., & Ameyaw, E. K. (2024). Global, regional, and national trends in tuberculosis incidence and main risk factors: A study using data from 2000 to 2021. *BMC Public Health*, 24, 12. <https://doi.org/10.1186/s12889-023-17495-6>
- Barberis, I., Bragazzi, N. L., Galluzzo, L., & Martini, M. (2017). The history of tuberculosis: From the first historical records to the isolation of Koch's bacillus. *Journal of Preventive Medicine and Hygiene*, 58(1), E9–E12.
- Bentley, S. D., Comas, I., Bryant, J. M., Walker, D., Smith, N. H., Harris, S. R., Thurston, S., Gagneux, S., Wood, J., Antonio, M., & Quail, M. A. (2012). The genome of Mycobacterium africanum West African 2 reveals a lineage-specific locus and genome erosion common to the M. tuberculosis complex. *PLoS Neglected Tropical Diseases*, 6(2), e1552.
- Boehme, C. C., Nabeta, P., Hillemann, D., Nicol, M. P., Shenai, S., Krapp, F., et al. (2010). Rapid molecular detection of tuberculosis and rifampin resistance. *New England Journal of Medicine*, 363, 1005-1015. <https://doi.org/10.1056/NEJMoa0907847>
- Coscolla, M., & Gagneux, S. (2014). Consequences of genomic diversity in Mycobacterium tuberculosis. In *Seminars in Immunology*, 26(6), 431–444. Academic Press.

- De Jong, B. C., Antonio, M., & Gagneux, S. (2010). Mycobacterium africanum—Review of an important cause of human tuberculosis in West Africa. *PLoS Neglected Tropical Diseases*, 4(9), e744.
- Dean, A. S., Zignol, M., Cabibbe, A. M., Falzon, D., Glaziou, P., Cirillo, D. M., Köser, C. U., Gonzalez-Angulo, L. Y., Tosas-Auget, O., Ismail, N., & Tahseen, S. (2020, January 21). Prevalence and genetic profiles of isoniazid resistance in tuberculosis patients: A multicountry analysis of cross-sectional data. *PLoS Medicine*, 17(1), e1003008. <https://doi.org/10.1371/journal.pmed.1003008>
- Dookie, N., Rambaran, S., Padayatchi, N., Mahomed, S., & Naidoo, K. (2018, May 1). Evolution of drug resistance in *Mycobacterium tuberculosis*: A review on the molecular determinants of resistance and implications for personalized care. *Journal of Antimicrobial Chemotherapy*, 73(5), 1138-1151. <https://doi.org/10.1093/jac/dky021>
- Gagneux, S. (2018). Ecology and evolution of *Mycobacterium tuberculosis*. *Nature Reviews Microbiology*, 16(4), 202–213.
- Gehre, F., Otu, J., Kendall, L., Forson, A., Kwara, A., Kudzawu, S., et al. (2016). The emerging threat of pre-extensively drug-resistant tuberculosis in West Africa: Preparing for large-scale tuberculosis research and drug resistance surveillance. *BMC Medicine*, 14, 160. <https://doi.org/10.1186/s12916-016-0704-5>
- Global Fund. (2021). *Approval of the Global Fund Strategy Narrative 46th Board Meeting*. <https://www.theglobalfund.org/board-decisions/eb01-2021-dp3/> (accessed August 22, 2024)
- Hamed, Z., Mohajeri, P., Farahani, A., Shamseddin, J., Zandi, M., Izadi, B., Atashi, S., Dastranj, M. (2021, March 1). The frequency of point mutations associated with resistance to isoniazid and rifampin among clinical isolates of multidrug-resistant *Mycobacterium tuberculosis* in the west of Iran. *Gene Reports*, 22, 100981. <https://doi.org/10.1016/j.genrep.2021.100981>
- Hassan, A. O., Olukolade, R., Ogbuji, Q. C., Afolabi, S., Okwuonye, L. C., Kusimo, O. C., Osho, J. A., Osinowo, K. A., & Ladipo, O. A. (2017). Knowledge about tuberculosis: A precursor to effective TB control—Findings from a follow-up national KAP study on tuberculosis among Nigerians. *Tuberculosis Research and Treatment*, 2017, 6309092. <https://doi.org/10.1155/2017/6309092>
- Islam, M. M., Adnan Hameed, H. M., Mugweru, J., Chhotaray, C., Wang, C., Tan, C., Liu, J., Li, X., Tan, S., Ojima, I., Wai Yew, W., Nuermberger, E., Lamichhane, G., & Zhang, T. (2017). Drug resistance mechanisms and novel drug targets for tuberculosis therapy. *Journal of Genetics and Genomics*, 44(1), 21–37.
- Kaaffah, S., Kusuma, I. Y., Renaldi, F. S., Pratiwi, A. D. E., Bahar, M. A., Lestari, Y. E. (2023). Knowledge, attitudes, and perceptions of tuberculosis in Indonesia: A multi-center cross-sectional study. *Infectious Diseases and Drug Resistance*, 16, 1787-1800. <https://doi.org/10.2147/IDR.S404171>
- Koleske, B. N., Jacobs, W. R., & Bishai, W. R. (2023). The *Mycobacterium tuberculosis* genome at 25 years: Lessons and lingering questions. *The Journal of Clinical Investigation*, 133(19).
- Koura, K. G., Mbitikon, O. B., Fiogbé, A. A., Ouédraogo, A. R., Kuate Kuate, A., Magassouba, A. S., Soumana, A., Hermana, G., Gning, B., Dogo, M. F., & Andefa, M. (2022, December 30). Programmatic implementation of contact investigation in eight African countries. *Tropical Medicine and Infectious Disease*, 8(1), 29. <https://doi.org/10.3390/tropicalmed8010029>
- Kraef, C., Bentzon, A., Panteleev, A., et al. (2021). Delayed diagnosis of tuberculosis in persons living with HIV in Eastern Europe: Associated factors and effect on mortality—a multicentre prospective cohort study. *BMC Infectious Diseases*, 21, 1038. <https://doi.org/10.1186/s12879-021-06745-w>
- Litvinjenko, S., Magwood, O., Wu, S., & Wei, X. (2023). Burden of tuberculosis among vulnerable populations worldwide: An overview of systematic reviews. *The Lancet Infectious Diseases*, 23(12), 1395–1407. [https://doi.org/10.1016/S1473-3099\(23\)00372-9](https://doi.org/10.1016/S1473-3099(23)00372-9)
- Mancuso, G., Midiri, A., De Gaetano, S., Ponzo, E., & Biondo, C. (2023). Tackling drug-resistant tuberculosis: New challenges from the old pathogen *Mycobacterium*

- tuberculosis. *Microorganisms*, 11(9), 2277. <https://doi.org/10.3390/microorganisms11092277>
- Msoka, E. F., Orina, F., Sanga, E. S., Miheso, B., Mwanyonga, S., Meme, H., Kiula, K., Liyoyo, A., Mwebaza, I. (2021). Qualitative assessment of the impact of socioeconomic and cultural barriers on uptake and utilisation of tuberculosis diagnostic and treatment tools in East Africa: A cross-section.
- N'Guessan, K., Ouassa, T., Dean, A. S., Alagna, R., Adagra, G. D., Ibode, V., et al. (2018). Multidrug-resistant tuberculosis in Côte d'Ivoire from 1995 to 2016: Results of national surveys. *European Journal of Microbiology & Immunology*, 8, 91-94. <https://doi.org/10.1556/1886.2018.00001>
- Nathanson, E., Nunn, P., Uplekar, M., Floyd, K., Jaramillo, E., Lönnroth, K., Weil, D., & Raviglione, M. (2010, September 9). MDR tuberculosis—critical steps for prevention and control. *New England Journal of Medicine*, 363(11), 1050-1058. <https://doi.org/10.1056/NEJMoa0907847>
- Nguta, J. M., Appiah-Opong, R., Nyarko, A. K., Yeboah-Manu, D., Addo, P. G., Otchere, I. D., Kissi-Twum, A. (2016, December 1). In vitro antimycobacterial activity and toxicity of eight medicinal plants against pathogenic and nonpathogenic mycobacterial strains. *International Journal of Mycobacteriology*, 5, S106-S107. <https://doi.org/10.1016/j.ijmyco.2016.06.119>
- Nimmo, C., Millard, J., Faulkner, V., Monteserin, J., Pugh, H., & Johnson, E. O. (2022, October 7). Evolution of *Mycobacterium tuberculosis* drug resistance in the genomic era. *Frontiers in Cellular and Infection Microbiology*, 12, 954074. <https://doi.org/10.3389/fcimb.2022.954074>
- Nina, P., Mark, S., Anh-Hue, T., Philip, L., & Brian, M. (2006, November 1). *OpenStax Microbiology*. <https://openstax.org/books/microbiology/pages/1-1-introduction-to-microbiology>
- Osei-Wusu, S., Otchere, I. D., Asare, P., Ntoumi, F., Zumla, A., Asante-Poku, A., & Yeboah-Manu, D. (2022). Relevance of genomic diversity of *Mycobacterium tuberculosis* complex in Africa. *International Journal of Infectious Diseases*, 124, S47–S49.
- Otchere, I. D., Asante-Poku, A., Akpadja, K. F., Diallo, A. B., Sanou, A., Asare, P., Osei-Wusu, S., Onyejebu, N., Diarra, B., Dagnra, Y. A., Kehinde, A., Antonio, M., & Yeboah-Manu, D. (2024). Opinion review of drug-resistant tuberculosis in West Africa: Tackling the challenges for effective control. *Frontiers in Public Health*, 12, 1374703. <https://doi.org/10.3389/fpubh.2024.1374703>
- Salari, N., Badoum, G., Ouedraogo, M., Dembélé, S. M., Nacanabo, R., Bonkougou, V., et al. (2010). Extensively drug-resistant tuberculosis, Burkina Faso. *Emerging Infectious Diseases*, 16(5), 840-842. <https://doi.org/10.3201/eid1605.091262>
- Seung, K. J., Keshavjee, S., & Rich, M. L. (2015, April 27). Multidrug-resistant tuberculosis and extensively drug-resistant tuberculosis. *Cold Spring Harbor Perspectives in Medicine*, 5(9), a017863. <https://doi.org/10.1101/cshperspect.a017863>
- Stop TB Partnership. (2022). *Digital TB Surveillance System Assessment Report*. <https://tbassessment.stoptb.org/> (accessed August 14, 2024)
- uyiringire, N., Mugisha, I. T., Tusubira, D., Munyampundu, J. P., Muvunyi, C. M., & Heyden, Y. V. (2022). In vitro antimycobacterial activity of medicinal plants *Lantana camara*, *Cryptolepis sanguinolenta*, and *Zanthoxylum leprieurii*. *Journal of Clinical Tuberculosis and Other Mycobacterial Diseases*, 27.
- Velen, K., Shingde, R. V., Ho, J., & Fox, G. J. (2021, December 1). The effectiveness of contact investigation among contacts of tuberculosis patients: A systematic review and meta-analysis. *European Respiratory Journal*, 58(6), 2100472. <https://doi.org/10.1183/13993003.00472-2021>
- World Health Organization (WHO). (2023). Tuberculosis deaths and disease increase during the COVID-19 pandemic. Retrieved August 16, 2024, from <https://www.who.int/news/item/27-10-2022-tuberculosis-deaths-and-disease-increase-during-the-covid-19-pandemic>
- World Health Organization (WHO). (2023). Tuberculosis in the WHO African Region: 2023 progress update. WHO African Region. Licence: BY-NC-SA 3.0 IGO.
- World Health Organization. (2018). *Global tuberculosis report 2018*. World Health Organization.

- https://www.who.int/tb/publications/global_report/en/
- World Health Organization. (2020). *Global tuberculosis report 2020*. World Health Organization.
https://www.who.int/tb/publications/global_report/en/
- World Health Organization. (2020). *WHO's Global Tuberculosis Programme*. Retrieved August 22, 2024, from <https://www.who.int/tb>
- World Health Organization. (2023). *Global tuberculosis report 2023*. World Health Organization.
https://www.who.int/tb/publications/global_report/en/
- Wu, S., Litvinjenko, S., Magwood, O., & Wei, X. (2023). Defining tuberculosis vulnerability based on an adapted social determinants of health framework: A narrative review. *Global Public Health*, 18(1), 2221729. <https://doi.org/10.1080/17441692.2023.2221729>
- Yang, T. W., Park, H. O., Jang, H. N., Yang, J. H., Kim, S. H., Moon, S. H., et al. (2017). Side effects associated with the treatment of multidrug-resistant tuberculosis at a tuberculosis referral hospital in South Korea. *Medicine*, 96, e7482. <https://doi.org/10.1097/MD.00000000000007482>
- Yusoof, K. A., García, J. I., Schami, A., Garcia-Vilanova, A., Kelley, H. V., Wang, S. H., et al. (2022). Tuberculosis phenotypic and genotypic drug susceptibility testing and immunodiagnostics: A review. *Frontiers in Immunology*, 13, 870768. <https://doi.org/10.3389/fimmu.2022.870768>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/124356>