



Quantitative Analysis of Total Carotenoids in Some Vegetables Consumed in Akwa Ibom State, Nigeria

Kufre Akpanikot^{1*}, Ofonimeh Udofia¹, Otobong D. Akan^{2,3} and Moses Uchechi¹

¹Department of Genetics and Biotechnology, Akwa Ibom State University, Mkpato Enin, Nigeria.

²Department of Microbiology, Akwa Ibom State University, Mkpato Enin, Nigeria.

³College of Food Science and Technology, Central South University of Forestry, Hunan, China.

Authors' contributions

This work was carried out in collaboration among all authors. Author KA designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors OU and ODA managed the analyses of the study. Author MU managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AFSJ/2019/v9i330010

Editor(s):

(1) Dr. Vodnar, Dan – Cristian, Assistant Professor, Department of Food Biotechnology, Food Chemistry and Food Microbiology, University of Agricultural Sciences and Veterinary Medicine, Cluj Napoca, Romania.

Reviewers:

(1) Marcelo Barcellos da Rosa, Federal University of Santa Maria, Brazil.

(2) Temitope Oyedepo, Adeleke University, Nigeria.

(3) Liamngee Kator, Benue State University, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48485>

Original Research Article

Received 02 March 2019

Accepted 11 May 2019

Published 17 May 2019

ABSTRACT

The quantitative analysis of total carotenoids in some vegetables consumed in Akwa Ibom State was evaluated using UV-Spectrophotometer and acetone as extraction solvent. Ten different vegetable samples i.e. Carrot (*Daucus carota*), purple egg plant (*Solanum melongena*), green egg plant (*Solanum melongena*), tomato (*Solanum lycopersicum*), cucumber (*Cucumis sativa*), fluted pumpkin (*Telfairia occidentalis*), spinach (*Spinacia oleracea*), red pepper (*Capsicum annum*), green pepper (*Capsicum annum*) and bitter leaf (*Vernonia amygdalina*) were purchased from local markets in Itam and Ikot Akpaden, both in Akwa Ibom State. The fresh samples were washed with tap water, followed by distilled water. The samples were homogenized separately using sterile mortar and pestle with 10 ml acetone as extraction solvent. The extract was centrifuged at 10,000rpm for 15 mins in 4°C and the supernatant analysed as soon as possible using spectrophotometer (UV-VIS 2500) at 470 nm. The quantification of carotenoids was carried out according to Harbone J B, 1973 and on the basis of Beer Lambert's equation: $A = \epsilon CL$ in mol/l and converted to mg/ml. From the result obtained, *Capsicum annum* (red pepper) had the highest amount of carotenoids (2.889

*Corresponding author: Email: kufreikot@gmail.com;

mg/ml), *Solanum lycopersicum* showed the lowest amount of carotenoid (0.260 mg/ml) the deep green vegetables also had high quantity of carotenoids. Carotenoids are important due to the health benefits associated with the pigment acting as provitamin A, a strong antioxidant and gives colouration to plants. It also helps in preventing photo-damage in plants and has some health benefits like improving sight, reducing cancer risk to animals too. Further studies analyzing the individual carotenoids is recommended and investigation on the effect of time on extraction.

Keywords: Carotenoids; vegetables; acetone; vitamin; antioxidant.

1. INTRODUCTION

Carotenoids are a class of phyto-nutrients (plant chemicals) found in the cells of a wide varieties of plants, bacteria and algae [1]. As a major plant pigment its distinct structures of biotechnological interest are natural colorants and human diseases' prevention [2]. These pigments play important roles in plant health and in extension, gives protective health benefits to consumers of carotenoids containing foods according to Szalay [1].

The total leaf pigment composition includes the chlorophyll-a, chlorophyll-b and carotenoids which are necessary for photosynthetic processes [3]. According to Sumanta et al. [3] some internal factors and environmental conditions can cause variations in the chlorophylls-carotenoids leaf pigment components [3]. The absorbance properties of these pigments facilitate their qualitative and quantitative analysis [3-4].

Carotenoids are usually represented by two carotenes (α - and β -) and five xanthophylls (luteins, zeaxanthin, violaxanthin, antheraxanthin and neoxanthin) which exhibit strong light absorption properties in the blue region of the spectrum and are non-uniformly distributed in photosynthetic and individual pigment protein complexes of chloroplast [5-8]. Carotenoids are synthesized in the plastid of plants and are the most diverse group of pigment found in nature. They are located in chromoplast and they give colouration to vegetables and fruits [3].

In human, carotenoids such as beta-carotene and lycopene are able to participate in free radical reactions where they help quench or prevent the formation of singlet oxygen through an efficient energy transfer process [1]. A recent study showed lycopene supplementation at the rate of 15 mg per day for 8 weeks to significantly decrease systolic blood pressures from the baseline value of 144mmHg to 134 mmHg in mildly hypertensive subjects [9-10]. In another

study a significant reduction in plasma lycopene was observed in the hypertensive patients compared to normal subjects [11]. When patients with liver cirrhosis, a condition closely associated with hypertension and disorders of the lymphatic circulation, were compared with matched controls a significant reduction in serum lycopene was observed along with other carotenoid antioxidants, retinol and vitamin E in the cirrhotic group [9,12].

In fact, dietary supplement of carotenoids shows a good relationship in reducing cancer risk by intake of various fruits and vegetables associated with as disease prevention [13]. To date, carotenoids have been associated with a vast range of diseases, especially degenerative diseases. Studies into carotenoids continue to demonstrate the importance of these phytochemicals. These include: oxidative stress, vitamin A deficiency, asthma and chronic obstructive pulmonary disease, Alzheimer's disease [14], cystic fibrosis, human immunodeficiency virus and many more [15].

Several attempts during the last decade have been undertaken to develop nondestructive techniques for carotenoid content assessment at both the leaf and canopy level [16-21]. For senescing leaves, reflectance indices sensitive to the molar concentration of carotenoids and chlorophylls ratio have been reported [16]. Over the years carotenoids have been known just as pro vitamin A with little known also of their quantity in some vegetables (especially those vegetable in Akwa Ibom State). This study therefore was designed to quantify carotenoids in some vegetables consumed in Akwa Ibom State.

2. MATERIALS AND METHODS

2.1 Sample Collection

Ten different fresh and matured vegetables (*Daucus carota*, *Solanum melongena* (purple),

Solanum melongena (green), *Solanum lycopersicum*, *Cucumis sativa*, *Telfairia occidentalis*, *Spinacia oleracea*, *Capsicum annuum* (red), *Capsicum annuum* (green) and *Vernonia amygdalina*) were purchased from local markets in Ikot Akpaden and Itam in Akwa Ibom State, Nigeria. The purchased vegetables were conveyed in sterile black nylon bags to the Genetics and Biotechnology laboratory, Akwa Ibom State University, Ikot Akpaden. The fresh samples were washed thoroughly with flowing tap water, followed by distilled water in the laboratory and further processed for the determination of total carotenoids content.

2.2 Carotenoid Determination

Ten (10) grams of each of the fresh vegetable sample was accurately weighed and homogenized for 10 mins, using a sterile mortar and pestle in 10 ml (10000 μ l) of extraction solvent (98% acetone). The homogenized sample was centrifuged at 10,000 rpm for 15minutes at 4°C. The supernatants (the pigment extracts) were separated and stored in ice in appropriately well labeled test tubes. Each extract was analyzed for total carotenoid content in triplicate using UV-VIS spectrophotometer (UV-2500) at a wavelength of 470 nm. Averages of triplicate absorbance readings at 470nm were recorded and the quantification of carotenoids carried out according to Harbone JB, 1973 and on the basis of the Beer-Lambert equation according to Lichtenthaler [8]:

$$A = \sum CL$$

Where;

A is the absorbance

\sum is the molar absorption coefficient for carotene (in acetone) at 470nm = 134×10^3 C is the concentration of the pigment (mol/l)

L is the path length of the light absorbing pigment (=1 cm).

2.3 Data Analysis

The triplicate result obtained was analyzed using ANOVA single factor which showed no significant difference, P- value of 0.99.

Table 1 was subjected to T-Test and it showed significant difference among the treatments with a P-value of 0.000097 is less than the P-value of the alpha level which is 0.05.

3. RESULTS AND DISCUSSION

Results for the extraction of carotenoids by acetone from different vegetable samples followed the sequence: *Capsicum annuum* (red) > *Vernonia amygdalin* > *Telfairia occidentalis* > *Spinacia oleracea* > *Solanum melongena* (purple) > *Solanum melongena* (green) > *Capsicum annuum* (green) > *Cucumis sativus* > *Daucus carota* > *Solanum lycopersicum*. The highest carotenoids content extraction was noted in *Capsicum annuum* (red) as compared to the nine other samples extracted from. Also, from the result obtained as shown in Table 1, coloured or deeply pigmented vegetables contained much more carotenoids when compared to their greenish counterparts as seen in *Solanum melongena* and *Capsicum annuum*. However, some deeply green vegetables like *Telfairia occidentalis* and *Vernonia amygdalina* with no deeply coloured counterpart contained high quantity of carotenoids too like the coloured vegetables with green counterparts. Results showed that deep green vegetables contains high quantity of carotenoids.

Carotenoid, chemically have high affinity towards polar solvents like the acetone and methanol [4,22], variations in the pigment concentrations is directly influenced by differences in environmental factors, seasons and even the species of the samples [3]. Carotenoid, therefore is important because of both its biotechnological potential [23], and its role in understanding the evolution of secondary metabolism [24].

Costache et al. [25] reported that carotenoids group and their derivatives consist of about 70 compounds that are present in most vegetables and fruits. Also, according to Vechetel and Ruppel [26], carotene pigments were the most important photosynthetic pigments and they prevented chlorophyll and thylakoid membrane from the damage of absorbed energy by peroxidation. Previous reports have considered carotenoid in bright coloured vegetables, since carotenoid contribute to their bright colouration. Times food reported that red Capsicum has more carotenoid than green Capsicum which is in line with the result of this research. However, this research has not only considered bright coloured vegetables with their green counterparts alone but has considered carotenoid in deep green vegetables which do not have bright coloured counterpart and the

Table 1. Spectrophotometric determination of absorbance and Quantification of carotenoids (mol/l) and (mg/ml)

S/N	Sample id	Absorbance	Quantity of carotenoids (mol/l)	Quantity of carotenoids (mg/ml) or mgg-1 fresh wt
1	<i>Spinacia oleracea</i>	0.543±0.010	4.052x10 ⁻⁶	2.175
2	<i>Solanum melongena</i> (green)	0.304±0.002	2.269x10 ⁻⁶	1.218
3	<i>Solanum melongena</i> (purple)	0.411±0.017	3.067x10 ⁻⁶	1.647
4	<i>Capsicum annuum</i> (red)	0.721±0.034	5.381x10 ⁻⁶	2.889
5	<i>Capsicum annuum</i> (green)	0.276±0.017	2.060x10 ⁻⁶	1.106
6	<i>Vernonia amygdalina</i>	0.709±0.004	5.291x10 ⁻⁶	2.841
7	<i>Daucus carota</i>	0.112±0.010	8.358x10 ⁻⁷	0.449
8	<i>Cucumis sativa</i>	0.146±0.003	1.089x10 ⁻⁶	0.585
9	<i>Solanum lycopersicum</i>	0.065±0.011	4.851x10 ⁻⁶	0.260
10	<i>Telfairia occidentalis</i>	0.635±0.018	4.739x10 ⁻⁶	2.544

Mean ±SD of mean in 3 Determinants

deep green vegetables without bright coloured counterpart has shown to contain high quality of carotenoid.

4. CONCLUSION

Having good vision and taking proactive measures in preventing the occurrence of certain diseases like cancer, cardiovascular disease is good for complete wellbeing of humans. Carotenoids which are present in vegetables act as pro-vitamin A and can serve these purposes. The study reveals the high amount of carotenoids in *Capsicum annuum* (red), *Telfairia occidentalis*, *Vernonia amygdalina*, *Solanum lycopersicum* (purple). These vegetables serve as a good source for high quantity of carotenoids. With health benefits in mind, people should be encouraged to incorporate carotenoids rich vegetables in their daily diets and in line with the result of the studies, *Capsicum annuum* (red), *Vernonia amygdalina*, *Solanum melongenta* (purple), *Telfairia occidentalis* being good sources with high amount of carotenoids are hereby recommended. Further studies on carotenoids analysis in vegetables is recommended to check the relationship between extraction and time alongside individual carotenoid concentrations.

ACKNOWLEDGEMENT

The authors are grateful to the Department of Biological Sciences Akwa Ibom State University for access to its laboratory and equipment during the research work. The authors are also grateful to the reviewers who diligently gave their time to scrutinize the research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Szalay J. What are Carotenoids? Livescience.com. 2015;52487. Available:https://amp.livescience.com/52487 [Accessed November 28, 2018]
- Klassen JL, Foght JM. Differences in carotenoid composition among hymenobacter and related strains support a tree-like model of carotenoid evolution. App Environ Microbiol. 2008;74:2016-2022.
- Sumanta N, Haque CI, Nishika J, Suprakash R. Spectrophotometric analysis of chlorophylls and carotenoids from commonly grown fern species by using various extracting solvents. Res J Chem Sci. 2014;4:63-69.
- James AO, Akaranta O. Inhibition of zinc in Hydrochloric acid solution by red onion skin acetone extract. Res J Chem Sci. 2011;1:31-37.
- Eldahshan OA, Singab AB. Carotenoids. J Pharmacog Phytochem. 2013;2:225-234.
- Demmig-Adams B, Gilmore AM, Adams III, WW. *In vivo* functions of carotenoids in higher plants. Fed Am Soc Exp Biol J. 1996;10: 403-412.
- Biswall B. Carotenoid catabolism during leaf senescence and its control by light. J Photochem Photobiol. 1995;30:3-14.

8. Lichtenthaler HK. Chlorophyll and carotenoids: Pigments of photosynthetic biomembranes. *Method Enzymol.* 1987; 148:350-382.
9. Paran E, Engelhard Y. Effect of Lyc-O-Mato, standardized tomato extract on blood pressure, serum lipoproteins plasma homocysteine and oxidative stress markers in grade 1 hypertensive patients. *Am Heart J.* 2001;151:100.
10. Paran E. Reducing hypertension with tomato lycopene. In: Rao, A. V., editor, *Tomatoes, lycopene and human health.* Scotland: Caledonian Sci Press. 2006; 169–82.
11. Moriel P, Sevanian A, Ajzen S, Zanella MT, Plavnik FL, Rubbo H, Abdalla DSP. Nitric oxide, cholesterol oxides and endothelium-dependent vasodilation in plasma of patients with essential hypertension. *Braz J Med Biol Res.* 2002; 35:1301-1309.
12. Rao AV, Mira MR, Rao LG. Lycopene. *Adv Food Nutri Res.* 2006;51:99-164.
13. Khachik F, Beecher GR, Goli MB. Separation, identification and quantification of carotenoids in fruits, vegetables and human plasma by high performance liquid chromatography. *Pure App Chem.* 1991;63:71-80.
14. Fleur B. The role of carotenoids as functional foods in disease prevention and treatment. *Nutrition.* 2009;3.
15. Braun L, Cohen M. *Herbs and natural supplements: An evidence based guide* (2nd Ed). Churchill Livingstone: Elsevier; 2007.
16. Gitelson AA, Zur Y, Chivokunova OB, Merzlyal MN. Assessing carotenoid content in plant leaves with reflectance spectroscopy. *Phytochem Phytobiol.* 2002; 75:272-281.
17. Blackburn GA. Relationship between spectral reflectance and pigment concentrations in stack of deciduous broadleaves. *Remote Sensing Environ.* 1999;70:224-237.
18. Gamon JA, Surfus JS. Assessing leaf pigment content and activity with a reflectancemeter. *New Phytologist J.* 1999;143:105-117.
19. Datt B. Remote sensing of chlorophyll a, chlorophyll b, chlorophyll a + b, and total carotenoid content in eucalyptus leaves. *Remote Sensing Environ.* 1998;66:111-121.
20. Blackburn GA. Quantifying chlorophylls and carotenoids at leaf and canopy scales: An evaluation of some hyperspectral approaches. *Remote Sensing Environ.* 1998;66: 273-285.
21. Chappelle EW, McMurtrey III JE. Ratio analysis of reflectance spectra (RARS): An alogarithm for the remote estimation of the concentrations of chlorophyll a, chlorophyll b, and carotenoids in soybean leaves. *Remote sensing Environ.* 1992;39:239-247.
22. Visca SI, Laslo V, Pantea S, Bandict GE. Chlorophyll and carotenoids pigments from Mistletoe (*Viscum album*) leaves using different solvents. *Facicula Biol.* 2010;2:213-218.
23. Sandmann G, Albrecht M, Schnurr G, Knörzer O, Böger P. The biotechnological potential and design of novel carotenoids by gene combination in *Escherichia coli*. *Trends Biotechnol.* 1999;17:233- 237.
24. Umeno D, Tobias AV, Arnold FH. Diversifying carotenoid biosynthetic pathways by directed evolution. *Microbiol Mol Bio Rev.* 2005;69:51-78.
25. Costache MA, Chappelle EW, Kim MS, McMurtrey III JE. Ratio analysis of reflectance spectra (RARS): An alogarithm for the remote estimation of the concentrations of chlorophyll a, chlorophyll b, and carotenoids in soybean leaves. *Remote sensing Environ.* 1992;39:239-247.
26. Vechetel BW, Ruppel HG. Lipid bodies in eremosphaeraviridis de vary (Chlorophyceae), *Plant Cell Physiol.* 1992; 31:41-48.

© 2019 Akpanikot et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
 The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/48485>